

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
27 September 2001 (27.09.2001)

PCT

(10) International Publication Number  
**WO 01/70816 A2**

(51) International Patent Classification<sup>7</sup>: **C07K 14/705**

MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,  
TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(21) International Application Number: PCT/US01/09050

(22) International Filing Date: 21 March 2001 (21.03.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/191,355 22 March 2000 (22.03.2000) US  
60/269,799 20 February 2001 (20.02.2001) US

(71) Applicant (for all designated States except US): **ROHM AND HAAS COMPANY** [US/US]; 100 Independence Mall West, Philadelphia, PA 19106-2399 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **PALLI, Subba, Reddy** [CA/US]; 832 Garfield Avenue, Lansdale, PA 19446 (US). **KAPITSKAYA, Marianna Zinovjevna** [RU/US]; 59 Wexford Road, North Wales, PA 19454 (US). **CRESS, Dean, Ervin** [US/US]; 104 Leslie Lane, Sounderton, PA 18964 (US).

(74) Agents: **MILLER, Suzanne, E.** et al.; Woodcock Washburn Kurtz Mackiewicz & Norris LLP, 46th floor, One Liberty Place, Philadelphia, PA 19103 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Declarations under Rule 4.17:**

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for all designations except US
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG)
- of inventorship (Rule 4.17(iv)) for US only
- of inventorship (Rule 4.17(iv)) for US only

**Published:**

- without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NOVEL ECDYSONE RECEPTOR-BASED INDUCIBLE GENE EXPRESSION SYSTEM

**GAL4Cre**

**GAL4DNAID**

**Cre**

**VP16RXR**

**VP16R**

**VP16R**

**pGAL4RELuc**

**GAL4RE**

**TATA**

(57) Abstract: This invention relates to the field of biotechnology or genetic engineering. Specifically, this invention relates to the field of gene expression. More specifically, this invention relates to a novel inducible gene expression system and methods of modulating gene expression in a host cell for applications such as gene therapy, large scale production of proteins and antibodies, cell-based high throughput screening assays, functional genomics and regulation of traits in transgenic plants and animals.

## NOVEL ECDYSONE RECEPTOR-BASED INDUCIBLE GENE EXPRESSION SYSTEM

This application claims priority to co-pending US provisional application Serial  
5 number 60/191,355, filed March 22, 2000 and to co-pending US provisional application Serial  
number 60/269,799, filed February 20, 2001.

### FIELD OF THE INVENTION

10 This invention relates to the field of biotechnology or genetic engineering. Specifically,  
this invention relates to the field of gene expression. More specifically, this invention relates to  
a novel ecdysone receptor-based inducible gene expression system and methods of modulating  
the expression of a gene within a host cell using this inducible gene expression system.

### 15 BACKGROUND OF THE INVENTION

In the field of genetic engineering, precise control of gene expression is a valuable tool  
for studying, manipulating, and controlling development and other physiological processes.  
Gene expression is a complex biological process involving a number of specific protein-protein  
20 interactions. In order for gene expression to be triggered, such that it produces the RNA  
necessary as the first step in protein synthesis, a transcriptional activator must be brought into  
proximity of a promoter that controls gene transcription. Typically, the transcriptional  
activator itself is associated with a protein that has at least one DNA binding domain that binds  
to DNA binding sites present in the promoter regions of genes. Thus, for gene expression to  
25 occur, a protein comprising a DNA binding domain and a transactivation domain located at an  
appropriate distance from the DNA binding domain must be brought into the correct position in  
the promoter region of the gene.

The traditional transgenic approach utilizes a cell-type specific promoter to drive the  
expression of a designed transgene. A DNA construct containing the transgene is first  
30 incorporated into a host genome. When triggered by a transcriptional activator, expression of  
the transgene occurs in a given cell type.

Another means to regulate expression of foreign genes in cells is through inducible  
promoters. Examples of the use of such inducible promoters include the PR1-a promoter,  
prokaryotic repressor-operator systems, immunosuppressive-immunophilin systems, and higher

eukaryotic transcription activation systems such as steroid hormone receptor systems and are described below.

The PR1-a promoter from tobacco is induced during the systemic acquired resistance response following pathogen attack. The use of PR1-a may be limited because it often  
5 responds to endogenous materials and external factors such as pathogens, UV-B radiation, and pollutants. Gene regulation systems based on promoters induced by heat shock, interferon and heavy metals have been described (Wurn et al., 1986, Proc. Natl. Acad. Sci. USA 83:5414-5418; Arnheiter et al., 1990 Cell 62:51-61; Filmus et al., 1992 Nucleic Acids Research 20:27550-27560). However, these systems have limitations due to their effect on expression of  
10 non-target genes. These systems are also leaky.

Prokaryotic repressor-operator systems utilize bacterial repressor proteins and the unique operator DNA sequences to which they bind. Both the tetracycline ("Tet") and lactose ("Lac") repressor-operator systems from the bacterium *Escherichia coli* have been used in plants and animals to control gene expression. In the Tet system, tetracycline binds to the TetR  
15 repressor protein, resulting in a conformational change which releases the repressor protein from the operator which as a result allows transcription to occur. In the Lac system, a lac operon is activated in response to the presence of lactose, or synthetic analogs such as isopropyl-b-D-thiogalactoside. Unfortunately, the use of such systems is restricted by unstable chemistry of the ligands, i.e. tetracycline and lactose, their toxicity, their natural presence, or  
20 the relatively high levels required for induction or repression. For similar reasons, utility of such systems in animals is limited.

Immunosuppressive molecules such as FK506, rapamycin and cyclosporine A can bind to immunophilins FKBP12, cyclophilin, etc. Using this information, a general strategy has been devised to bring together any two proteins simply by placing FK506 on each of the two  
25 proteins or by placing FK506 on one and cyclosporine A on another one. A synthetic homodimer of FK506 (FK1012) or a compound resulted from fusion of FK506-cyclosporine (FKCsA) can then be used to induce dimerization of these molecules (Spencer et al., 1993, *Science* 262:1019-24; Belshaw et al., 1996 *Proc Natl Acad Sci U S A* 93:4604-7). Gal4 DNA binding domain fused to FKBP12 and VP16 activator domain fused to cyclophilin, and FKCsA  
30 compound were used to show heterodimerization and activation of a reporter gene under the control of a promoter containing Gal4 binding sites. Unfortunately, this system includes immunosuppressants that can have unwanted side effects and therefore, limits its use for various mammalian gene switch applications.

Higher eukaryotic transcription activation systems such as steroid hormone receptor systems have also been employed. Steroid hormone receptors are members of the nuclear receptor superfamily and are found in vertebrate and invertebrate cells. Unfortunately, use of steroidal compounds that activate the receptors for the regulation of gene expression, particularly in plants and mammals, is limited due to their involvement in many other natural biological pathways in such organisms. In order to overcome such difficulties, an alternative system has been developed using insect ecdysone receptors (EcR).

- Growth, molting, and development in insects are regulated by the ecdysone steroid hormone (molting hormone) and the juvenile hormones (Dhadialla, et al., 1998. Annu. Rev. Entomol. 43: 545-569). The molecular target for ecdysone in insects consists of at least ecdysone receptor (EcR) and ultraspiracle protein (USP). EcR is a member of the nuclear steroid receptor super family that is characterized by signature DNA and ligand binding domains, and an activation domain (Koelle et al. 1991, Cell, 67:59-77). EcR receptors are responsive to a number of steroidal compounds such as ponasterone A and muristerone A. Recently, non-steroidal compounds with ecdysteroid agonist activity have been described, including the commercially available insecticides tebufenozide and methoxyfenozide that are marketed world wide by Rohm and Haas Company (see International Patent Application No. PCT/EP96/00686 and US Patent 5,530,028). Both analogs have exceptional safety profiles to other organisms.
- International Patent Application No. PCT/US97/05330 (WO 97/38117) discloses methods for modulating the expression of an exogenous gene in which a DNA construct comprising the exogenous gene and an ecdysone response element is activated by a second DNA construct comprising an ecdysone receptor that, in the presence of a ligand therefor, and optionally in the presence of a receptor capable of acting as a silent partner, binds to the ecdysone response element to induce gene expression. The ecdysone receptor of choice was isolated from *Drosophila melanogaster*. Typically, such systems require the presence of the silent partner, preferably retinoid X receptor (RXR), in order to provide optimum activation. In mammalian cells, insect ecdysone receptor (EcR) heterodimerizes with retinoid X receptor (RXR) and regulates expression of target genes in a ligand dependent manner. International Patent Application No. PCT/US98/14215 (WO 99/02683) discloses that the ecdysone receptor isolated from the silk moth *Bombyx mori* is functional in mammalian systems without the need for an exogenous dimer partner.

U.S. Patent No. 5,880,333 discloses a *Drosophila melanogaster* EcR and ultraspiracle

(USP) heterodimer system used in plants in which the transactivation domain and the DNA binding domain are positioned on two different hybrid proteins. Unfortunately, this system is not effective for inducing reporter gene expression in animal cells (for comparison, see Example 1.2, below).

5 In each of these cases, the transactivation domain and the DNA binding domain (either as native EcR as in International Patent Application No. PCT/US98/14215 or as modified EcR as in International Patent Application No. PCT/US97/05330) were incorporated into a single molecule and the other heterodimeric partners, either USP or RXR, were used in their native state.

10 Drawbacks of the above described EcR-based gene regulation systems include a considerable background activity in the absence of ligands and that these systems are not applicable for use in both plants and animals (see U.S. Patent No. 5,880,333). For most applications that rely on modulating gene expression, these EcR-based systems are undesirable. Therefore, a need exists in the art for improved systems to precisely modulate the expression of  
15 exogenous genes in both plants and animals. Such improved systems would be useful for applications such as gene therapy, large scale production of proteins and antibodies, cell-based high throughput screening assays, functional genomics and regulation of traits in transgenic animals. Improved systems that are simple, compact, and dependent on ligands that are relatively inexpensive, readily available, and of low toxicity to the host would prove useful for  
20 regulating biological systems.

Various publications are cited herein, the disclosures of which are incorporated by reference in their entireties. However, the citation of any reference herein should not be construed as an admission that such reference is available as "Prior Art" to the instant application.

25

### SUMMARY OF THE INVENTION

The present invention relates to a novel ecdysone receptor-based inducible gene expression system, novel receptor polynucleotides and polypeptides for use in the novel  
30 inducible gene expression system, and methods of modulating the expression of a gene within a host cell using this inducible gene expression system. In particular, Applicants' invention relates to an improved gene expression modulation system comprising a polynucleotide encoding a receptor polypeptide comprising a truncation mutation.

Specifically, the present invention relates to a gene expression modulation system comprising: a) a first gene expression cassette that is capable of being expressed in a host cell comprising a polynucleotide that encodes a first polypeptide comprising: i) a DNA-binding domain that recognizes a response element associated with a gene whose expression is to be modulated; and ii) a ligand binding domain comprising a ligand binding domain from a nuclear receptor; and b) a second gene expression cassette that is capable of being expressed in the host cell comprising a polynucleotide sequence that encodes a second polypeptide comprising: i) a transactivation domain; and ii) a ligand binding domain comprising a ligand binding domain from a nuclear receptor other than an ultraspiracle receptor; wherein the DNA binding domain and the transactivation domain are from a polypeptide other than an ecdysone receptor, a retinoid X receptor, or an ultraspiracle receptor; wherein the ligand binding domains from the first polypeptide and the second polypeptide are different and dimerize.

In a specific embodiment, the ligand binding domain of the first polypeptide comprises an ecdysone receptor (EcR) ligand binding domain

In another specific embodiment, the ligand binding domain of the second polypeptide comprises a retinoid X receptor (RXR) ligand binding domain.

In a preferred embodiment, the ligand binding domain of the first polypeptide comprises an ecdysone receptor ligand binding domain and the ligand binding domain of the second polypeptide comprises a retinoid X receptor ligand binding domain

The present invention also relates to a gene expression modulation system according to the invention further comprising c) a third gene expression cassette comprising: i) a response element to which the DNA-binding domain of the first polypeptide binds; ii) a promoter that is activated by the transactivation domain of the second polypeptide; and iii) the gene whose expression is to be modulated.

The present invention also relates to an isolated polynucleotide encoding a truncated EcR or a truncated RXR polypeptide, wherein the truncation mutation affects ligand binding activity or ligand sensitivity.

In particular, the present invention relates to an isolated polynucleotide encoding a truncated EcR or a truncated RXR polypeptide comprising a truncation mutation that reduces ligand binding activity or ligand sensitivity of said EcR or RXR polypeptide. In a specific embodiment, the present invention relates to an isolated polynucleotide encoding a truncated EcR or a truncated RXR polypeptide comprising a truncation mutation that reduces steroid binding activity or steroid sensitivity of said EcR or RXR polypeptide. In another specific

embodiment, the present invention relates to an isolated polynucleotide encoding a truncated EcR or a truncated RXR polypeptide comprising a truncation mutation that reduces non-steroid binding activity or non-steroid sensitivity of said EcR or RXR polypeptide.

The present invention also relates to an isolated polynucleotide encoding a truncated  
5 EcR or a truncated RXR polypeptide comprising a truncation mutation that enhances ligand binding activity or ligand sensitivity of said EcR or RXR polypeptide. In a specific embodiment, the present invention relates to an isolated polynucleotide encoding a truncated EcR or a truncated RXR polypeptide comprising a truncation mutation that enhances steroid binding activity or steroid sensitivity of said EcR or RXR polypeptide. In another specific  
10 embodiment, the present invention relates to an isolated polynucleotide encoding a truncated EcR or a truncated RXR polypeptide comprising a truncation mutation that enhances non-steroid binding activity or non-steroid sensitivity of said EcR or RXR polypeptide.

The present invention also relates to an isolated polynucleotide encoding a truncated RXR polypeptide comprising a truncation mutation that increases ligand sensitivity of a  
15 heterodimer comprising the truncated retinoid X receptor polypeptide and a dimerization partner. In a specific embodiment, the dimerization partner is an ecdysone receptor polypeptide.

The present invention also relates to an isolated polypeptide encoded by a polynucleotide according to Applicants' invention. In particular, the present invention relates  
20 to an isolated truncated EcR or truncated RXR polypeptide comprising a truncation mutation, wherein the EcR or RXR polypeptide is encoded by a polynucleotide according to the invention.

Thus, the present invention also relates to an isolated truncated EcR or truncated RXR polypeptide comprising a truncation mutation that affects ligand binding activity or ligand  
25 sensitivity of said EcR or RXR polypeptide.

Applicants' invention also relates to methods of modulating gene expression in a host cell using a gene expression modulation system according to the invention. Specifically, Applicants' invention provides a method of modulating the expression of a gene in a host cell comprising the gene to be modulated comprising the steps of: a) introducing into the host cell a  
30 gene expression modulation system according to the invention; and b) introducing into the host cell a ligand that independently combines with the ligand binding domains of the first polypeptide and the second polypeptide of the gene expression modulation system; wherein the gene to be expressed is a component of a chimeric gene comprising: i) a response element

comprising a domain to which the DNA binding domain from the first polypeptide binds; ii) a promoter that is activated by the transactivation domain of the second polypeptide; and iii) the gene whose expression is to be modulated, whereby a complex is formed comprising the ligand, the first polypeptide, and the second polypeptide, and whereby the complex modulates

5 expression of the gene in the host cell.

Applicants' invention also provides an isolated host cell comprising an inducible gene expression system according to the invention. The present invention also relates to an isolated host cell comprising a polynucleotide or polypeptide according to the invention. Accordingly, Applicants' invention also relates to a non-human organism comprising a host cell according to

10 the invention.



**BRIEF DESCRIPTION OF THE DRAWINGS**

**Figure 1:** An ecdysone receptor-based gene expression system comprising a first gene expression cassette encoding a Gal4DBD-CfEcRDEF chimeric polypeptide and a second gene expression cassette encoding a VP16AD-MmRXRDEF chimeric polypeptide; prepared as

5 described in Example 1 (switch 1.1).

**Figure 2:** An ecdysone receptor-based gene expression system comprising a first gene expression cassette encoding a Gal4DBD-CfEcRDEF chimeric polypeptide and a second gene expression cassette encoding a VP16AD-CfUSPDEF chimeric polypeptide; prepared as described in Example 1 (switch 1.2).

10 **Figure 3:** An ecdysone receptor-based gene expression system comprising a first gene expression cassette encoding a Gal4DBD-MmRXRDEF chimeric polypeptide and a second gene expression cassette encoding a VP16AD-CfEcRCDEF chimeric polypeptide; prepared as described in Example 1 (switch 1.3).

**Figure 4:** An ecdysone receptor-based gene expression system comprising a first gene  
15 expression cassette encoding a Gal4DBD-MmRXRDEF chimeric polypeptide and a second gene expression cassette encoding a VP16AD-DmEcRCDEF chimeric polypeptide; prepared as described in Example 1 (switch 1.4).

**Figure 5:** An ecdysone receptor-based gene expression system comprising a first gene expression cassette encoding a Gal4DBD-CfUSPDEF chimeric polypeptide and a second gene  
20 expression cassette encoding a VP16AD-CfEcRCDEF chimeric polypeptide; prepared as described in Example 1 (switch 1.5).

**Figure 6:** An ecdysone receptor-based gene expression system comprising a first gene expression cassette encoding a Gal4DBD-CfEcRDEF-VP16AD chimeric polypeptide; prepared as described in Example 1 (switch 1.6).

25 **Figure 7:** An ecdysone receptor-based gene expression system comprising a first gene expression cassette encoding a VP16AD-CfEcRCDEF chimeric polypeptide; prepared as described in Example 1 (switch 1.7).

**Figure 8:** An ecdysone receptor-based gene expression system comprising a first gene expression cassette encoding a VP16AD-DmEcRCDEF chimeric polypeptide and a second  
30 gene expression cassette encoding a MmRXR polypeptide; prepared as described in Example 1 (switch 1.8).

Figure 9: An ecdysone receptor-based gene expression system comprising a first gene expression cassette encoding a VP16AD-CfEcRCDEF chimeric polypeptide and a second gene expression cassette encoding a MmRXR polypeptide; prepared as described in Example 1 (switch 1.9).

- 5 Figure 10: An ecdysone receptor-based gene expression system comprising a gene expression cassette encoding a Gal4DBD-CfEcRCDEF chimeric polypeptide; prepared as described in Example 1 (switch 1.10).

Figure 11: Expression data of GAL4CfEcRA/BCDEF, GAL4CfEcRCDEF, GAL4CfEcR1/2CDEF, GAL4CfEcRDEF, GAL4CfEcREF, GAL4CfEcRDE truncation

- 10 mutants transfected into NIH3T3 cells along with VP16MmRXRDE, pFRLUc and pTKRL plasmid DNAs.

Figure 12: Expression data of GAL4CfEcRA/BCDEF, GAL4CfEcRCDEF, GAL4CfEcR1/2CDEF, GAL4CfEcRDEF, GAL4CfEcREF, GAL4CfEcRDE truncation mutants transfected into 3T3 cells along with VP16MmRXRE, pFRLUc and pTKRL plasmid

- 15 DNAs.

Figure 13: Expression data of VP16MmRXRA/BCDEF, VP16MmRXRCDEF, VP16MmRXRDEF, VP16MmRXREF, VP16MmRXRBam-EF, VP16MmRXRAF2del constructs transfected into NIH3T3 cells along with GAL4CfEcRCDEF, pFRLUc and pTKRL plasmid DNAs.

- 20 Figure 14: Expression data of VP16MmRXRA/BCDEF, VP16MmRXRCDEF, VP16MmRXRDEF, VP16MmRXREF, VP16MmRXRBam-EF, VP16MmRXRAF2del constructs transfected into NIH3T3 cells along with GAL4CfEcRDEF, pFRLUc and pTKRL plasmid DNAs.

Figure 15: Expression data of various truncated CfEcR and MmRXR receptor pairs

- 25 transfected into NIH3T3 cells along with GAL4CfEcRDEF, pFRLUc and pTKRL plasmid DNAs.

#### DETAILED DESCRIPTION OF THE INVENTION

- 30 Applicants have now developed an improved ecdysone receptor-based inducible gene expression system comprising a truncation mutant of an ecdysone receptor or a retinoid X receptor (RXR) polypeptide that affects ligand binding activity or ligand sensitivity. This mutational effect may increase or reduce ligand binding activity or ligand sensitivity and may

be steroid or non-steroid specific. Thus, Applicants' invention provides an improved ecdysone receptor-based inducible gene expression system useful for modulating expression of a gene of interest in a host cell. In a particularly desirable embodiment, Applicants' invention provides an inducible gene expression system that has a reduced level of background gene expression and responds to submicromolar concentrations of non-steroidal ligand. Thus, Applicants' novel inducible gene expression system and its use in methods of modulating gene expression in a host cell overcome the limitations of currently available inducible expression systems and provide the skilled artisan with an effective means to control gene expression.

The present invention provides a novel inducible gene expression system that can be used to modulate gene expression in both prokaryotic and eukaryotic host cells. Applicants' invention is useful for applications such as gene therapy, large scale production of proteins and antibodies, cell-based high throughput screening assays, functional genomics and regulation of traits in transgenic organisms.

## 15 DEFINITIONS

In this disclosure, a number of terms and abbreviations are used. The following definitions are provided and should be helpful in understanding the scope and practice of the present invention.

In a specific embodiment, the term "about" or "approximately" means within 20%, preferably within 10%, more preferably within 5%, and even more preferably within 1% of a given value or range.

The term "substantially free" means that a composition comprising "A" (where "A" is a single protein, DNA molecule, vector, recombinant host cell, etc.) is substantially free of "B" (where "B" comprises one or more contaminating proteins, DNA molecules, vectors, etc.) when at least about 75% by weight of the proteins, DNA, vectors (depending on the category of species to which A and B belong) in the composition is "A". Preferably, "A" comprises at least about 90% by weight of the A+B species in the composition, most preferably at least about 99% by weight. It is also preferred that a composition, which is substantially free of contamination, contain only a single molecular weight species having the activity or characteristic of the species of interest.

The term "isolated" for the purposes of the present invention designates a biological material (nucleic acid or protein) that has been removed from its original environment (the environment in which it is naturally present).

For example, a polynucleotide present in the natural state in a plant or an animal is not isolated. The same polynucleotide separated from the adjacent nucleic acids in which it is naturally present. The term "purified" does not require the material to be present in a form exhibiting absolute purity, exclusive of the presence of other compounds. It is rather a relative  
5 definition.

A polynucleotide is in the "purified" state after purification of the starting material or of the natural material by at least one order of magnitude, preferably 2 or 3 and preferably 4 or 5 orders of magnitude.

A "nucleic acid" is a polymeric compound comprised of covalently linked subunits  
10 called nucleotides. Nucleic acid includes polynucleic acid (RNA) and polydeoxyribonucleic acid (DNA), both of which may be single-stranded or double-stranded. DNA includes but is not limited to cDNA, genomic DNA, plasmids DNA, synthetic DNA, and semi-synthetic DNA. DNA may be linear, circular, or supercoiled.

A "nucleic acid molecule" refers to the phosphate ester polymeric form of  
15 ribonucleosides (adenosine, guanosine, uridine or cytidine; "RNA molecules") or deoxyribonucleosides (deoxyadenosine, deoxyguanosine, deoxythymidine, or deoxycytidine; "DNA molecules"), or any phosphoester analogs thereof, such as phosphorothioates and thioesters, in either single stranded form, or a double-stranded helix. Double stranded DNA-DNA, DNA-RNA and RNA-RNA helices are possible. The term nucleic acid molecule, and in  
20 particular DNA or RNA molecule, refers only to the primary and secondary structure of the molecule, and does not limit it to any particular tertiary forms. Thus, this term includes double-stranded DNA found, *inter alia*, in linear or circular DNA molecules (*e.g.*, restriction fragments), plasmids, and chromosomes. In discussing the structure of particular double-stranded DNA molecules, sequences may be described herein according to the normal  
25 convention of giving only the sequence in the 5' to 3' direction along the non-transcribed strand of DNA (*i.e.*, the strand having a sequence homologous to the mRNA). A "recombinant DNA molecule" is a DNA molecule that has undergone a molecular biological manipulation.

The term "fragment" will be understood to mean a nucleotide sequence of reduced length relative to the reference nucleic acid and comprising, over the common portion, a  
30 nucleotide sequence identical to the reference nucleic acid. Such a nucleic acid fragment according to the invention may be, where appropriate, included in a larger polynucleotide of which it is a constituent. Such fragments comprise, or alternatively consist of, oligonucleotides ranging in length from at least 8, 10, 12, 15, 18, 20 to 25, 30, 40, 50, 70, 80, 100, 200, 500,

1000 or 1500 consecutive nucleotides of a nucleic acid according to the invention.

As used herein, an "isolated nucleic acid fragment" is a polymer of RNA or DNA that is single- or double-stranded, optionally containing synthetic, non-natural or altered nucleotide bases. An isolated nucleic acid fragment in the form of a polymer of DNA may be comprised  
5 of one or more segments of cDNA, genomic DNA or synthetic DNA.

A "gene" refers to an assembly of nucleotides that encode a polypeptide, and includes cDNA and genomic DNA nucleic acids. "Gene" also refers to a nucleic acid fragment that expresses a specific protein or polypeptide, including regulatory sequences preceding (5' non-coding sequences) and following (3' non-coding sequences) the coding sequence. "Native gene"  
10 refers to a gene as found in nature with its own regulatory sequences. "Chimeric gene" refers to any gene that is not a native gene, comprising regulatory and/or coding sequences that are not found together in nature. Accordingly, a chimeric gene may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner different than that  
15 found in nature. A chimeric gene may comprise coding sequences derived from different sources and/or regulatory sequences derived from different sources. "Endogenous gene" refers to a native gene in its natural location in the genome of an organism. A "foreign" gene or "heterologous" gene refers to a gene not normally found in the host organism, but that is introduced into the host organism by gene transfer. Foreign genes can comprise native genes  
20 inserted into a non-native organism, or chimeric genes. A "transgene" is a gene that has been introduced into the genome by a transformation procedure.

"Heterologous" DNA refers to DNA not naturally located in the cell, or in a chromosomal site of the cell. Preferably, the heterologous DNA includes a gene foreign to the cell.

25 The term "genome" includes chromosomal as well as mitochondrial, chloroplast and viral DNA or RNA.

A nucleic acid molecule is "hybridizable" to another nucleic acid molecule, such as a cDNA, genomic DNA, or RNA, when a single stranded form of the nucleic acid molecule can anneal to the other nucleic acid molecule under the appropriate conditions of temperature and  
30 solution ionic strength (see Sambrook *et al.*, 1989 *infra*). Hybridization and washing conditions are well known and exemplified in Sambrook, J., Fritsch, E. F. and Maniatis, T. *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor (1989), particularly Chapter 11 and Table 11.1 therein (entirely

incorporated herein by reference). The conditions of temperature and ionic strength determine the "stringency" of the hybridization.

Stringency conditions can be adjusted to screen for moderately similar fragments, such as homologous sequences from distantly related organisms, to highly similar fragments, such as genes that duplicate functional enzymes from closely related organisms. For preliminary screening for homologous nucleic acids, low stringency hybridization conditions, corresponding to a  $T_m$  of 55°, can be used, *e.g.*, 5x SSC, 0.1% SDS, 0.25% milk, and no formamide; or 30% formamide, 5x SSC, 0.5% SDS). Moderate stringency hybridization conditions correspond to a higher  $T_m$ , *e.g.*, 40% formamide, with 5x or 6x SCC. High stringency hybridization conditions correspond to the highest  $T_m$ , *e.g.*, 50% formamide, 5x or 6x SCC. Hybridization requires that the two nucleic acids contain complementary sequences, although depending on the stringency of the hybridization, mismatches between bases are possible.

The term "complementary" is used to describe the relationship between nucleotide bases that are capable of hybridizing to one another. For example, with respect to DNA, adenosine is complementary to thymine and cytosine is complementary to guanine. Accordingly, the instant invention also includes isolated nucleic acid fragments that are complementary to the complete sequences as disclosed or used herein as well as those substantially similar nucleic acid sequences.

In a specific embodiment, the term "standard hybridization conditions" refers to a  $T_m$  of 55°C, and utilizes conditions as set forth above. In a preferred embodiment, the  $T_m$  is 60°C; in a more preferred embodiment, the  $T_m$  is 65°C.

Post-hybridization washes also determine stringency conditions. One set of preferred conditions uses a series of washes starting with 6X SSC, 0.5% SDS at room temperature for 15 minutes (min), then repeated with 2X SSC, 0.5% SDS at 45°C for 30 minutes, and then repeated twice with 0.2X SSC, 0.5% SDS at 50°C for 30 minutes. A more preferred set of stringent conditions uses higher temperatures in which the washes are identical to those above except for the temperature of the final two 30 min washes in 0.2X SSC, 0.5% SDS was increased to 60°C. Another preferred set of highly stringent conditions uses two final washes in 0.1X SSC, 0.1% SDS at 65°C. Hybridization requires that the two nucleic acids comprise complementary sequences, although depending on the stringency of the hybridization, mismatches between bases are possible.

The appropriate stringency for hybridizing nucleic acids depends on the length of the nucleic acids and the degree of complementation, variables well known in the art. The greater

the degree of similarity or homology between two nucleotide sequences, the greater the value of  $T_m$  for hybrids of nucleic acids having those sequences. The relative stability (corresponding to higher  $T_m$ ) of nucleic acid hybridizations decreases in the following order: RNA:RNA, DNA:RNA, DNA:DNA. For hybrids of greater than 100 nucleotides in length, equations for calculating  $T_m$  have been derived (see Sambrook *et al.*, *supra*, 9.50-9.51). For hybridization with shorter nucleic acids, *i.e.*, oligonucleotides, the position of mismatches becomes more important, and the length of the oligonucleotide determines its specificity (see Sambrook *et al.*, *supra*, 11.7-11.8).

In one embodiment the length for a hybridizable nucleic acid is at least about 10 nucleotides. Preferable a minimum length for a hybridizable nucleic acid is at least about 15 nucleotides; more preferably at least about 20 nucleotides; and most preferably the length is at least 30 nucleotides. Furthermore, the skilled artisan will recognize that the temperature and wash solution salt concentration may be adjusted as necessary according to factors such as length of the probe.

The term "probe" refers to a single-stranded nucleic acid molecule that can base pair with a complementary single stranded target nucleic acid to form a double-stranded molecule.

As used herein, the term "oligonucleotide" refers to a nucleic acid, generally of at least 18 nucleotides, that is hybridizable to a genomic DNA molecule, a cDNA molecule, a plasmid DNA or an mRNA molecule. Oligonucleotides can be labeled, *e.g.*, with  $^{32}\text{P}$ -nucleotides or nucleotides to which a label, such as biotin, has been covalently conjugated. A labeled oligonucleotide can be used as a probe to detect the presence of a nucleic acid.

Oligonucleotides (one or both of which may be labeled) can be used as PCR primers, either for cloning full length or a fragment of a nucleic acid, or to detect the presence of a nucleic acid.

An oligonucleotide can also be used to form a triple helix with a DNA molecule. Generally, oligonucleotides are prepared synthetically, preferably on a nucleic acid synthesizer.

Accordingly, oligonucleotides can be prepared with non-naturally occurring phosphoester analog bonds, such as thioester bonds, etc.

A "primer" is an oligonucleotide that hybridizes to a target nucleic acid sequence to create a double stranded nucleic acid region that can serve as an initiation point for DNA synthesis under suitable conditions. Such primers may be used in a polymerase chain reaction.

"Polymerase chain reaction" is abbreviated PCR and means an *in vitro* method for enzymatically amplifying specific nucleic acid sequences. PCR involves a repetitive series of temperature cycles with each cycle comprising three stages: denaturation of the template

nucleic acid to separate the strands of the target molecule, annealing a single stranded PCR oligonucleotide primer to the template nucleic acid, and extension of the annealed primer(s) by DNA polymerase. PCR provides a means to detect the presence of the target molecule and, under quantitative or semi-quantitative conditions, to determine the relative amount of that target molecule within the starting pool of nucleic acids.

“Reverse transcription-polymerase chain reaction” is abbreviated RT-PCR and means an *in vitro* method for enzymatically producing a target cDNA molecule or molecules from an RNA molecule or molecules, followed by enzymatic amplification of a specific nucleic acid sequence or sequences within the target cDNA molecule or molecules as described above. RT-PCR also provides a means to detect the presence of the target molecule and, under quantitative or semi-quantitative conditions, to determine the relative amount of that target molecule within the starting pool of nucleic acids.

A DNA “coding sequence” is a double-stranded DNA sequence that is transcribed and translated into a polypeptide in a cell *in vitro* or *in vivo* when placed under the control of appropriate regulatory sequences. “Suitable regulatory sequences” refer to nucleotide sequences located upstream (5' non-coding sequences), within, or downstream (3' non-coding sequences) of a coding sequence, and which influence the transcription, RNA processing or stability, or translation of the associated coding sequence. Regulatory sequences may include promoters, translation leader sequences, introns, polyadenylation recognition sequences, RNA processing site, effector binding site and stem-loop structure. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxyl) terminus. A coding sequence can include, but is not limited to, prokaryotic sequences, cDNA from mRNA, genomic DNA sequences, and even synthetic DNA sequences. If the coding sequence is intended for expression in a eukaryotic cell, a polyadenylation signal and transcription termination sequence will usually be located 3' to the coding sequence.

“Open reading frame” is abbreviated ORF and means a length of nucleic acid sequence, either DNA, cDNA or RNA, that comprises a translation start signal or initiation codon, such as an ATG or AUG, and a termination codon and can be potentially translated into a polypeptide sequence.

The term “head-to-head” is used herein to describe the orientation of two polynucleotide sequences in relation to each other. Two polynucleotides are positioned in a head-to-head orientation when the 5' end of the coding strand of one polynucleotide is adjacent



to the 5' end of the coding strand of the other polynucleotide, whereby the direction of transcription of each polynucleotide proceeds away from the 5' end of the other polynucleotide.

The term "head-to-head" may be abbreviated (5')-to-(5') and may also be indicated by the symbols ( $\leftarrow \rightarrow$ ) or ( $3' \leftarrow 5' \rightarrow 3'$ ).

- 5 The term "tail-to-tail" is used herein to describe the orientation of two polynucleotide sequences in relation to each other. Two polynucleotides are positioned in a tail-to-tail orientation when the 3' end of the coding strand of one polynucleotide is adjacent to the 3' end of the coding strand of the other polynucleotide, whereby the direction of transcription of each polynucleotide proceeds toward the other polynucleotide. The term "tail-to-tail" may be
- 10 abbreviated (3')-to-(3') and may also be indicated by the symbols ( $\rightarrow \leftarrow$ ) or ( $5' \rightarrow 3' 3' \leftarrow 5'$ ).

- The term "head-to-tail" is used herein to describe the orientation of two polynucleotide sequences in relation to each other. Two polynucleotides are positioned in a head-to-tail orientation when the 5' end of the coding strand of one polynucleotide is adjacent to the 3' end of the coding strand of the other polynucleotide, whereby the direction of transcription of each
- 15 polynucleotide proceeds in the same direction as that of the other polynucleotide. The term "head-to-tail" may be abbreviated (5')-to-(3') and may also be indicated by the symbols ( $\rightarrow$ ) or ( $5' \rightarrow 3' 5' \rightarrow 3'$ ).

- The term "downstream" refers to a nucleotide sequence that is located 3' to reference nucleotide sequence. In particular, downstream nucleotide sequences generally relate to
- 20 sequences that follow the starting point of transcription. For example, the translation initiation codon of a gene is located downstream of the start site of transcription.

- The term "upstream" refers to a nucleotide sequence that is located 5' to reference nucleotide sequence. In particular, upstream nucleotide sequences generally relate to sequences that are located on the 5' side of a coding sequence or starting point of transcription. For
- 25 example, most promoters are located upstream of the start site of transcription.

The terms "restriction endonuclease" and "restriction enzyme" refer to an enzyme that binds and cuts within a specific nucleotide sequence within double stranded DNA.

- "Homologous recombination" refers to the insertion of a foreign DNA sequence into another DNA molecule, e.g., insertion of a vector in a chromosome. Preferably, the vector
- 30 targets a specific chromosomal site for homologous recombination. For specific homologous recombination, the vector will contain sufficiently long regions of homology to sequences of the chromosome to allow complementary binding and incorporation of the vector into the chromosome. Longer regions of homology, and greater degrees of sequence similarity, may

increase the efficiency of homologous recombination.

Several methods known in the art may be used to propagate a polynucleotide according to the invention. Once a suitable host system and growth conditions are established, recombinant expression vectors can be propagated and prepared in quantity. As described  
5 herein, the expression vectors which can be used include, but are not limited to, the following vectors or their derivatives: human or animal viruses such as vaccinia virus or adenovirus; insect viruses such as baculovirus; yeast vectors; bacteriophage vectors (*e.g.*, lambda), and plasmid and cosmid DNA vectors, to name but a few.

A "vector" is any means for the cloning of and/or transfer of a nucleic acid into a host  
10 cell. A vector may be a replicon to which another DNA segment may be attached so as to bring about the replication of the attached segment. A "replicon" is any genetic element (*e.g.*, plasmid, phage, cosmid, chromosome, virus) that functions as an autonomous unit of DNA replication *in vivo*, *i.e.*, capable of replication under its own control. The term "vector" includes both viral and nonviral means for introducing the nucleic acid into a cell *in vitro*, *ex*  
15 *vivo* or *in vivo*. A large number of vectors known in the art may be used to manipulate nucleic acids, incorporate response elements and promoters into genes, etc. Possible vectors include, for example, plasmids or modified viruses including, for example bacteriophages such as lambda derivatives, or plasmids such as PBR322 or pUC plasmid derivatives, or the Bluescript vector. For example, the insertion of the DNA fragments corresponding to response elements  
20 and promoters into a suitable vector can be accomplished by ligating the appropriate DNA fragments into a chosen vector that has complementary cohesive termini. Alternatively, the ends of the DNA molecules may be enzymatically modified or any site may be produced by ligating nucleotide sequences (linkers) into the DNA termini. Such vectors may be engineered to contain selectable marker genes that provide for the selection of cells that have incorporated  
25 the marker into the cellular genome. Such markers allow identification and/or selection of host cells that incorporate and express the proteins encoded by the marker.

Viral vectors, and particularly retroviral vectors, have been used in a wide variety of gene delivery applications in cells, as well as living animal subjects. Viral vectors that can be used include but are not limited to retrovirus, adeno-associated virus, pox, baculovirus,  
30 vaccinia, herpes simplex, Epstein-Barr, adenovirus, geminivirus, and caulimovirus vectors. Non-viral vectors include plasmids, liposomes, electrically charged lipids (cytofectins), DNA-protein complexes, and biopolymers. In addition to a nucleic acid, a vector may also comprise one or more regulatory regions, and/or selectable markers useful in selecting, measuring, and

monitoring nucleic acid transfer results (transfer to which tissues, duration of expression, etc.).

The term "plasmid" refers to an extra chromosomal element often carrying a gene that is not part of the central metabolism of the cell, and usually in the form of circular double-stranded DNA molecules. Such elements may be autonomously replicating sequences, genome  
5 integrating sequences, phage or nucleotide sequences, linear, circular, or supercoiled, of a single- or double-stranded DNA or RNA, derived from any source, in which a number of nucleotide sequences have been joined or recombined into a unique construction which is capable of introducing a promoter fragment and DNA sequence for a selected gene product along with appropriate 3' untranslated sequence into a cell.

10 A "cloning vector" is a "replicon", which is a unit length of a nucleic acid, preferably DNA, that replicates sequentially and which comprises an origin of replication, such as a plasmid, phage or cosmid, to which another nucleic acid segment may be attached so as to bring about the replication of the attached segment. Cloning vectors may be capable of replication in one cell type and expression in another ("shuttle vector").

15 Vectors may be introduced into the desired host cells by methods known in the art, e.g., transfection, electroporation, microinjection, transduction, cell fusion, DEAE dextran, calcium phosphate precipitation, lipofection (lysosome fusion), use of a gene gun, or a DNA vector transporter (see, e.g., Wu et al., 1992, J. Biol. Chem. 267:963-967; Wu and Wu, 1988, J. Biol. Chem. 263:14621-14624; and Hartmut et al., Canadian Patent Application No.  
20 2,012,311, filed March 15, 1990).

A polynucleotide according to the invention can also be introduced *in vivo* by lipofection. For the past decade, there has been increasing use of liposomes for encapsulation and transfection of nucleic acids *in vitro*. Synthetic cationic lipids designed to limit the difficulties and dangers encountered with liposome mediated transfection can be used to prepare liposomes for *in vivo*  
25 transfection of a gene encoding a marker (Felgner et al., 1987. PNAS 84:7413; Mackey, et al., 1988. Proc. Natl. Acad. Sci. U.S.A. 85:8027-8031; and Ulmer et al., 1993. Science 259:1745-1748). The use of cationic lipids may promote encapsulation of negatively charged nucleic acids, and also promote fusion with negatively charged cell membranes (Felgner and Ringold, 1989. Science 337:387-388). Particularly useful lipid compounds and compositions for transfer of  
30 nucleic acids are described in International Patent Publications WO95/18863 and WO96/17823, and in U.S. Patent No. 5,459,127. The use of lipofection to introduce exogenous genes into the specific organs *in vivo* has certain practical advantages. Molecular targeting of liposomes to specific cells represents one area of benefit. It is clear that directing transfection to particular cell

types would be particularly preferred in a tissue with cellular heterogeneity, such as pancreas, liver, kidney, and the brain. Lipids may be chemically coupled to other molecules for the purpose of targeting (Mackey, et al., 1988, *supra*). Targeted peptides, *e.g.*, hormones or neurotransmitters, and proteins such as antibodies, or non-peptide molecules could be coupled to liposomes

5 chemically.

Other molecules are also useful for facilitating transfection of a nucleic acid *in vivo*, such as a cationic oligopeptide (*e.g.*, WO95/21931), peptides derived from DNA binding proteins (*e.g.*, WO96/25508), or a cationic polymer (*e.g.*, WO95/21931).

It is also possible to introduce a vector *in vivo* as a naked DNA plasmid (see U.S.  
10 Patents 5,693,622, 5,589,466 and 5,580,859). Receptor-mediated DNA delivery approaches can also be used (Curiel et al., 1992. Hum. Gene Ther. 3:147-154; and Wu and Wu, 1987. J. Biol. Chem. 262:4429-4432).

The term "transfection" means the uptake of exogenous or heterologous RNA or DNA by a cell. A cell has been "transfected" by exogenous or heterologous RNA or DNA when  
15 such RNA or DNA has been introduced inside the cell. A cell has been "transformed" by exogenous or heterologous RNA or DNA when the transfected RNA or DNA effects a phenotypic change. The transforming RNA or DNA can be integrated (covalently linked) into chromosomal DNA making up the genome of the cell.

"Transformation" refers to the transfer of a nucleic acid fragment into the genome of a  
20 host organism, resulting in genetically stable inheritance. Host organisms containing the transformed nucleic acid fragments are referred to as "transgenic" or "recombinant" or "transformed" organisms.

The term "genetic region" will refer to a region of a nucleic acid molecule or a nucleotide sequence that comprises a gene encoding a polypeptide.

25 In addition, the recombinant vector comprising a polynucleotide according to the invention may include one or more origins for replication in the cellular hosts in which their amplification or their expression is sought; markers or selectable markers.

The term "selectable marker" means an identifying factor, usually an antibiotic or chemical resistance gene, that is able to be selected for based upon the marker gene's effect,  
30 *i.e.*, resistance to an antibiotic, resistance to a herbicide, colorimetric markers, enzymes, fluorescent markers, and the like, wherein the effect is used to track the inheritance of a nucleic acid of interest and/or to identify a cell or organism that has inherited the nucleic acid of

interest. Examples of selectable marker genes known and used in the art include: genes providing resistance to ampicillin, streptomycin, gentamycin, kanamycin, hygromycin, bialaphos herbicide, sulfonamide, and the like; and genes that are used as phenotypic markers, *i.e.*, anthocyanin regulatory genes, isopentenyl transferase gene, and the like.

- 5           The term "reporter gene" means a nucleic acid encoding an identifying factor that is able to be identified based upon the reporter gene's effect, wherein the effect is used to track the inheritance of a nucleic acid of interest, to identify a cell or organism that has inherited the nucleic acid of interest, and/or to measure gene expression induction or transcription. Examples of reporter genes known and used in the art include: luciferase (Luc), green fluorescent protein
- 10 (GFP), chloramphenicol acetyltransferase (CAT),  $\beta$ -galactosidase (LacZ),  $\beta$ -glucuronidase (Gus), and the like. Selectable marker genes may also be considered reporter genes.

- "Promoter" refers to a DNA sequence capable of controlling the expression of a coding sequence or functional RNA. In general, a coding sequence is located 3' to a promoter sequence. Promoters may be derived in their entirety from a native gene, or be composed of
- 15 different elements derived from different promoters found in nature, or even comprise synthetic DNA segments. It is understood by those skilled in the art that different promoters may direct the expression of a gene in different tissues or cell types, or at different stages of development, or in response to different environmental or physiological conditions. Promoters that cause a gene to be expressed in most cell types at most times are commonly referred to as "constitutive
- 20 promoters". Promoters that cause a gene to be expressed in a specific cell type are commonly referred to as "cell-specific promoters" or "tissue-specific promoters". Promoters that cause a gene to be expressed at a specific stage of development or cell differentiation are commonly referred to as "developmentally-specific promoters" or "cell differentiation-specific promoters". Promoters that are induced and cause a gene to be expressed following exposure or treatment
- 25 of the cell with an agent, biological molecule, chemical, ligand, light, or the like that induces the promoter are commonly referred to as "inducible promoters" or "regulatable promoters". It is further recognized that since in most cases the exact boundaries of regulatory sequences have not been completely defined, DNA fragments of different lengths may have identical promoter activity.

- 30           A "promoter sequence" is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. For purposes of defining the present invention, the promoter sequence is bounded at its 3' terminus by the transcription initiation site and extends upstream (5' direction) to include

the minimum number of bases or elements necessary to initiate transcription at levels detectable above background. Within the promoter sequence will be found a transcription initiation site (conveniently defined for example, by mapping with nuclease S1), as well as protein binding domains (consensus sequences) responsible for the binding of RNA polymerase.

5 A coding sequence is "under the control" of transcriptional and translational control sequences in a cell when RNA polymerase transcribes the coding sequence into mRNA, which is then trans-RNA spliced (if the coding sequence contains introns) and translated into the protein encoded by the coding sequence.

"Transcriptional and translational control sequences" are DNA regulatory sequences, 10 such as promoters, enhancers, terminators, and the like, that provide for the expression of a coding sequence in a host cell. In eukaryotic cells, polyadenylation signals are control sequences.

The term "response element" means one or more cis-acting DNA elements which confer responsiveness on a promoter mediated through interaction with the DNA-binding 15 domains of the first chimeric gene. This DNA element may be either palindromic (perfect or imperfect) in its sequence or composed of sequence motifs or half sites separated by a variable number of nucleotides. The half sites can be similar or identical and arranged as either direct or inverted repeats or as a single half site or multimers of adjacent half sites in tandem. The response element may comprise a minimal promoter isolated from different organisms 20 depending upon the nature of the cell or organism into which the response element will be incorporated. The DNA binding domain of the first hybrid protein binds, in the presence or absence of a ligand, to the DNA sequence of a response element to initiate or suppress transcription of downstream gene(s) under the regulation of this response element. Examples of DNA sequences for response elements of the natural ecdysone receptor include: 25 RRGG/TTCANTGAC/ACY (see Cherbas L., et. al., (1991), *Genes Dev.* 5, 120-131); AGGTCAN<sub>(n)</sub>AGGTCA, where N<sub>(n)</sub> can be one or more spacer nucleotides (see D'Avino PP., et. al., (1995), *Mol. Cell. Endocrinol.* 113, 1-9); and GGGTTGAATGAATTT (see Antoniewski C., et. al., (1994), *Mol. Cell Biol.* 14, 4465-4474).

The term "operably linked" refers to the association of nucleic acid sequences on a 30 single nucleic acid fragment so that the function of one is affected by the other. For example, a promoter is operably linked with a coding sequence when it is capable of affecting the expression of that coding sequence (i.e., that the coding sequence is under the transcriptional control of the promoter). Coding sequences can be operably linked to regulatory sequences in

sense or antisense orientation.

The term "expression", as used herein, refers to the transcription and stable accumulation of sense (mRNA) or antisense RNA derived from a nucleic acid or polynucleotide. Expression may also refer to translation of mRNA into a protein or

5 polypeptide.

The terms "cassette", "expression cassette" and "gene expression cassette" refer to a segment of DNA that can be inserted into a nucleic acid or polynucleotide at specific restriction sites or by homologous recombination. The segment of DNA comprises a polynucleotide that encodes a polypeptide of interest, and the cassette and restriction sites are designed to ensure

10 insertion of the cassette in the proper reading frame for transcription and translation.

"Transformation cassette" refers to a specific vector comprising a polynucleotide that encodes a polypeptide of interest and having elements in addition to the polynucleotide that facilitate transformation of a particular host cell. Cassettes, expression cassettes, gene expression cassettes and transformation cassettes of the invention may also comprise elements that allow

15 for enhanced expression of a polynucleotide encoding a polypeptide of interest in a host cell.

These elements may include, but are not limited to: a promoter, a minimal promoter, an enhancer, a response element, a terminator sequence, a polyadenylation sequence, and the like.

For purposes of this invention, the term "gene switch" refers to the combination of a response element associated with a promoter, and an EcR based system which, in the presence  
20 of one or more ligands, modulates the expression of a gene into which the response element and promoter are incorporated.

The terms "modulate" and "modulates" mean to induce, reduce or inhibit nucleic acid or gene expression, resulting in the respective induction, reduction or inhibition of protein or polypeptide production.

25 The plasmids or vectors according to the invention may further comprise at least one promoter suitable for driving expression of a gene in a host cell. The term "expression vector" means a vector, plasmid or vehicle designed to enable the expression of an inserted nucleic acid sequence following transformation into the host. The cloned gene, i.e., the inserted nucleic acid sequence, is usually placed under the control of control elements such as a promoter, a minimal  
30 promoter, an enhancer, or the like. Initiation control regions or promoters, which are useful to drive expression of a nucleic acid in the desired host cell are numerous and familiar to those skilled in the art. Virtually any promoter capable of driving these genes is suitable for the present invention including but not limited to: viral promoters, plant promoters, bacterial

promoters, animal promoters, mammalian promoters, synthetic promoters, constitutive promoters, tissue specific promoter, developmental specific promoters, inducible promoters, light regulated promoters; *CYC1*, *HIS3*, *GAL1*, *GAL4*, *GAL10*, *ADH1*, *PGK*, *PHO5*, *GAPDH*, *ADC1*, *TRP1*, *URA3*, *LEU2*, *ENO*, *TPI*, alkaline phosphatase promoters (useful for expression in *Saccharomyces*); *AOX1* promoter (useful for expression in *Pichia*); b-lactamase, *lac*, *ara*, *tet*, *trp*, *lP<sub>L</sub>*, *lP<sub>R</sub>*, *T7*, *tac*, and *trc* promoters (useful for expression in *Escherichia coli*); and light regulated-, seed specific-, pollen specific-, ovary specific-, pathogenesis or disease related-, cauliflower mosaic virus 35S, CMV 35S minimal, cassava vein mosaic virus (CsVMV), chlorophyll a/b binding protein, ribulose 1, 5-bisphosphate carboxylase, shoot-specific, root specific, chitinase, stress inducible, rice tungro bacilliform virus, plant super-promoter, potato leucine aminopeptidase, nitrate reductase, mannopine synthase, nopaline synthase, ubiquitin, zein protein, and anthocyanin promoters (useful for expression in plant cells); animal and mammalian promoters known in the art include, but are not limited to, the SV40 early (SV40e) promoter region, the promoter contained in the 3' long terminal repeat (LTR) of Rous sarcoma virus (RSV), the promoters of the E1A or major late promoter (MLP) genes of adenoviruses, the cytomegalovirus early promoter, the herpes simplex virus (HSV) thymidine kinase (TK) promoter, an elongation factor 1 alpha (EF1) promoter, a phosphoglycerate kinase (PGK) promoter, a ubiquitin (Ubc) promoter, an albumin promoter, the regulatory sequences of the mouse metallothionein-L promoter, and transcriptional control regions, the ubiquitous promoters (HPRT, vimentin,  $\alpha$ -actin, tubulin and the like), the promoters of the intermediate filaments (desmin, neurofilaments, keratin, GFAP, and the like), the promoters of therapeutic genes (of the MDR, CFTR or factor VIII type, and the like), and promoters that exhibit tissue specificity and have been utilized in transgenic animals, such as the elastase I gene control region which is active in pancreatic acinar cells; insulin gene control region active in pancreatic beta cells, immunoglobulin gene control region active in lymphoid cells, mouse mammary tumor virus control region active in testicular, breast, lymphoid and mast cells; albumin gene, Apo AI and Apo AII control regions active in liver, alpha-fetoprotein gene control region active in liver, alpha 1-antitrypsin gene control region active in the liver, beta-globin gene control region active in myeloid cells, myelin basic protein gene control region active in oligodendrocyte cells in the brain, myosin light chain-2 gene control region active in skeletal muscle, and gonadotropic releasing hormone gene control region active in the hypothalamus, pyruvate kinase promoter, villin promoter, promoter of the fatty acid binding intestinal protein, promoter of the smooth muscle cell  $\alpha$ -actin, and the like. In a preferred



embodiment of the invention, the promoter is selected from the group consisting of a cauliflower mosaic virus 35S promoter, a cassava vein mosaic virus promoter, and a cauliflower mosaic virus 35S minimal promoter, an elongation factor 1 alpha (EF1) promoter, a phosphoglycerate kinase (PGK) promoter, a ubiquitin (Ubc) promoter, and an albumin promoter. In addition, these expression sequences may be modified by addition of enhancer or regulatory sequences and the like.

Enhancers that may be used in embodiments of the invention include but are not limited to: tobacco mosaic virus enhancer, cauliflower mosaic virus 35S enhancer, tobacco etch virus enhancer, ribulose 1, 5-bisphosphate carboxylase enhancer, rice tungro bacilliform virus enhancer, and other plant and viral gene enhancers, and the like.

Termination control regions, *i.e.*, terminator or polyadenylation sequences, may also be derived from various genes native to the preferred hosts. Optionally, a termination site may be unnecessary, however, it is most preferred if included. In a preferred embodiment of the invention, the termination control region may be comprise or be derived from a synthetic sequence, synthetic polyadenylation signal, an SV40 late polyadenylation signal, an SV40 polyadenylation signal, a bovine growth hormone (BGH) polyadenylation signal, nopaline synthase (nos), cauliflower mosaic virus (CaMV), octopine synthase (ocs), Agrobacterium, viral, and plant terminator sequences, or the like.

The terms "3' non-coding sequences" or "3' untranslated region (UTR)" refer to DNA sequences located downstream (3') of a coding sequence and may comprise polyadenylation [poly(A)] recognition sequences and other sequences encoding regulatory signals capable of affecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by affecting the addition of polyadenylic acid tracts to the 3' end of the mRNA precursor.


"Regulatory region" means a nucleic acid sequence which regulates the expression of a second nucleic acid sequence. A regulatory region may include sequences which are naturally responsible for expressing a particular nucleic acid (a homologous region) or may include sequences of a different origin that are responsible for expressing different proteins or even synthetic proteins (a heterologous region). In particular, the sequences can be sequences of prokaryotic, eukaryotic, or viral genes or derived sequences that stimulate or repress transcription of a gene in a specific or non-specific manner and in an inducible or non-inducible manner. Regulatory regions include origins of replication, RNA splice sites, promoters, enhancers, transcriptional termination sequences, and signal sequences which direct the

polypeptide into the secretory pathways of the target cell.

A regulatory region from a "heterologous source" is a regulatory region that is not naturally associated with the expressed nucleic acid. Included among the heterologous regulatory regions are regulatory regions from a different species, regulatory regions from a different gene, hybrid regulatory sequences, and regulatory sequences which do not occur in nature, but which are designed by one having ordinary skill in the art.

"RNA transcript" refers to the product resulting from RNA polymerase-catalyzed transcription of a DNA sequence. When the RNA transcript is a perfect complementary copy of the DNA sequence, it is referred to as the primary transcript or it may be a RNA sequence derived from post-transcriptional processing of the primary transcript and is referred to as the mature RNA. "Messenger RNA (mRNA)" refers to the RNA that is without introns and that can be translated into protein by the cell. "cDNA" refers to a double-stranded DNA that is complementary to and derived from mRNA. "Sense" RNA refers to RNA transcript that includes the mRNA and so can be translated into protein by the cell. "Antisense RNA" refers to a RNA transcript that is complementary to all or part of a target primary transcript or mRNA and that blocks the expression of a target gene. The complementarity of an antisense RNA may be with any part of the specific gene transcript, i.e., at the 5' non-coding sequence, 3' non-coding sequence, or the coding sequence. "Functional RNA" refers to antisense RNA, ribozyme RNA, or other RNA that is not translated yet has an effect on cellular processes.

A "polypeptide" is a polymeric compound comprised of covalently linked amino acid residues. Amino acids have the following general structure:



25

Amino acids are classified into seven groups on the basis of the side chain R: (1) aliphatic side chains, (2) side chains containing a hydroxylic (OH) group, (3) side chains containing sulfur atoms, (4) side chains containing an acidic or amide group, (5) side chains containing a basic group, (6) side chains containing an aromatic ring, and (7) proline, an imino acid in which the side chain is fused to the amino group. A polypeptide of the invention preferably comprises at least about 14 amino acids.

30

A "protein" is a polypeptide that performs a structural or functional role in a living

cell.

An "isolated polypeptide" or "isolated protein" is a polypeptide or protein that is substantially free of those compounds that are normally associated therewith in its natural state (e.g., other proteins or polypeptides, nucleic acids, carbohydrates, lipids). "Isolated" is not meant to exclude artificial or synthetic mixtures with other compounds, or the presence of impurities which do not interfere with biological activity, and which may be present, for example, due to incomplete purification, addition of stabilizers, or compounding into a pharmaceutically acceptable preparation.

"Fragment" of a polypeptide according to the invention will be understood to mean a polypeptide whose amino acid sequence is shorter than that of the reference polypeptide and which comprises, over the entire portion with these reference polypeptides, an identical amino acid sequence. Such fragments may, where appropriate, be included in a larger polypeptide of which they are a part. Such fragments of a polypeptide according to the invention may have a length of 10, 15, 20, 30 to 40, 50, 100, 200 or 300 amino acids.

A "variant" of a polypeptide or protein is any analogue, fragment, derivative, or mutant which is derived from a polypeptide or protein and which retains at least one biological property of the polypeptide or protein. Different variants of the polypeptide or protein may exist in nature. These variants may be allelic variations characterized by differences in the nucleotide sequences of the structural gene coding for the protein, or may involve differential splicing or post-translational modification. The skilled artisan can produce variants having single or multiple amino acid substitutions, deletions, additions, or replacements. These variants may include, *inter alia*: (a) variants in which one or more amino acid residues are substituted with conservative or non-conservative amino acids, (b) variants in which one or more amino acids are added to the polypeptide or protein, (c) variants in which one or more of the amino acids includes a substituent group, and (d) variants in which the polypeptide or protein is fused with another polypeptide such as serum albumin. The techniques for obtaining these variants, including genetic (suppressions, deletions, mutations, etc.), chemical, and enzymatic techniques, are known to persons having ordinary skill in the art. A variant polypeptide preferably comprises at least about 14 amino acids.

A "heterologous protein" refers to a protein not naturally produced in the cell.

A "mature protein" refers to a post-translationally processed polypeptide; i.e., one from which any pre- or propeptides present in the primary translation product have been removed. "Precursor" protein refers to the primary product of translation of mRNA; i.e., with

pre- and propeptides still present. Pre- and propeptides may be but are not limited to intracellular localization signals.

The term "signal peptide" refers to an amino terminal polypeptide preceding the secreted mature protein. The signal peptide is cleaved from and is therefore not present in the  
5 mature protein. Signal peptides have the function of directing and translocating secreted proteins across cell membranes. Signal peptide is also referred to as signal protein.

A "signal sequence" is included at the beginning of the coding sequence of a protein to be expressed on the surface of a cell. This sequence encodes a signal peptide, N-terminal to the mature polypeptide, that directs the host cell to translocate the polypeptide. The term  
10 "translocation signal sequence" is used herein to refer to this sort of signal sequence. Translocation signal sequences can be found associated with a variety of proteins native to eukaryotes and prokaryotes, and are often functional in both types of organisms.

The term "homology" refers to the percent of identity between two polynucleotide or two polypeptide moieties. The correspondence between the sequence from one moiety to  
15 another can be determined by techniques known to the art. For example, homology can be determined by a direct comparison of the sequence information between two polypeptide molecules by aligning the sequence information and using readily available computer programs. Alternatively, homology can be determined by hybridization of polynucleotides under conditions that form stable duplexes between homologous regions, followed by digestion  
20 with single-stranded-specific nuclease(s) and size determination of the digested fragments.

As used herein, the term "homologous" in all its grammatical forms and spelling variations refers to the relationship between proteins that possess a "common evolutionary origin," including proteins from superfamilies (*e.g.*, the immunoglobulin superfamily) and homologous proteins from different species (*e.g.*, myosin light chain, etc.) (Reeck et al., 1987,  
25 Cell 50:667.). Such proteins (and their encoding genes) have sequence homology, as reflected by their high degree of sequence similarity.

Accordingly, the term "sequence similarity" in all its grammatical forms refers to the degree of identity or correspondence between nucleic acid or amino acid sequences of proteins that may or may not share a common evolutionary origin (*see* Reeck et al., 1987, Cell 50:667).  
30 As used herein, the term "homologous" in all its grammatical forms and spelling variations refers to the relationship between proteins that possess a "common evolutionary origin," including proteins from superfamilies and homologous proteins from different species (Reeck *et al.*, *supra*). Such proteins (and their encoding genes) have sequence homology, as reflected by

their high degree of sequence similarity. However, in common usage and in the instant application, the term "homologous," when modified with an adverb such as "highly," may refer to sequence similarity and not a common evolutionary origin.

In a specific embodiment, two DNA sequences are "substantially homologous" or  
5 "substantially similar" when at least about 50% (preferably at least about 75%, and most preferably at least about 90 or 95%) of the nucleotides match over the defined length of the DNA sequences. Sequences that are substantially homologous can be identified by comparing the sequences using standard software available in sequence data banks, or in a Southern hybridization experiment under, for example, stringent conditions as defined for that particular  
10 system. Defining appropriate hybridization conditions is within the skill of the art. See, e.g., Sambrook *et al.*, 1989, *supra*.

As used herein, "substantially similar" refers to nucleic acid fragments wherein changes in one or more nucleotide bases results in substitution of one or more amino acids, but do not affect the functional properties of the protein encoded by the DNA sequence.  
15 "Substantially similar" also refers to nucleic acid fragments wherein changes in one or more nucleotide bases does not affect the ability of the nucleic acid fragment to mediate alteration of gene expression by antisense or co-suppression technology. "Substantially similar" also refers to modifications of the nucleic acid fragments of the instant invention such as deletion or insertion of one or more nucleotide bases that do not substantially affect the functional  
20 properties of the resulting transcript. It is therefore understood that the invention encompasses more than the specific exemplary sequences. Each of the proposed modifications is well within the routine skill in the art, as is determination of retention of biological activity of the encoded products.

Moreover, the skilled artisan recognizes that substantially similar sequences  
25 encompassed by this invention are also defined by their ability to hybridize, under stringent conditions (0.1X SSC, 0.1% SDS, 65°C and washed with 2X SSC, 0.1% SDS followed by 0.1X SSC, 0.1% SDS), with the sequences exemplified herein. Substantially similar nucleic acid fragments of the instant invention are those nucleic acid fragments whose DNA sequences are at least 70% identical to the DNA sequence of the nucleic acid fragments reported herein.  
30 Preferred substantially nucleic acid fragments of the instant invention are those nucleic acid fragments whose DNA sequences are at least 80% identical to the DNA sequence of the nucleic acid fragments reported herein. More preferred nucleic acid fragments are at least 90% identical to the DNA sequence of the nucleic acid fragments reported herein. Even more

preferred are nucleic acid fragments that are at least 95% identical to the DNA sequence of the nucleic acid fragments reported herein.

Two amino acid sequences are "substantially homologous" or "substantially similar" when greater than about 40% of the amino acids are identical, or greater than 60% are similar  
5 (functionally identical). Preferably, the similar or homologous sequences are identified by alignment using, for example, the GCG (Genetics Computer Group, Program Manual for the GCG Package, Version 7, Madison, Wisconsin) pileup program.

The term "corresponding to" is used herein to refer to similar or homologous sequences, whether the exact position is identical or different from the molecule to which the  
10 similarity or homology is measured. A nucleic acid or amino acid sequence alignment may include spaces. Thus, the term "corresponding to" refers to the sequence similarity, and not the numbering of the amino acid residues or nucleotide bases.

A "substantial portion" of an amino acid or nucleotide sequence comprises enough of the amino acid sequence of a polypeptide or the nucleotide sequence of a gene to putatively  
15 identify that polypeptide or gene, either by manual evaluation of the sequence by one skilled in the art, or by computer-automated sequence comparison and identification using algorithms such as BLAST (Basic Local Alignment Search Tool; Altschul, S. F., et al., (1993) *J. Mol. Biol.* 215:403-410; see also [www.ncbi.nlm.nih.gov/BLAST/](http://www.ncbi.nlm.nih.gov/BLAST/)). In general, a sequence of ten or more contiguous amino acids or thirty or more nucleotides is necessary in order to putatively  
20 identify a polypeptide or nucleic acid sequence as homologous to a known protein or gene. Moreover, with respect to nucleotide sequences, gene specific oligonucleotide probes comprising 20-30 contiguous nucleotides may be used in sequence-dependent methods of gene identification (e.g., Southern hybridization) and isolation (e.g., *in situ* hybridization of bacterial colonies or bacteriophage plaques). In addition, short oligonucleotides of 12-15 bases may be  
25 used as amplification primers in PCR in order to obtain a particular nucleic acid fragment comprising the primers. Accordingly, a "substantial portion" of a nucleotide sequence comprises enough of the sequence to specifically identify and/or isolate a nucleic acid fragment comprising the sequence.

The term "percent identity", as known in the art, is a relationship between two or more  
30 polypeptide sequences or two or more polynucleotide sequences, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. "Identity" and "similarity" can be readily calculated by

- known methods, including but not limited to those described in: *Computational Molecular Biology* (Lesk, A. M., ed.) Oxford University Press, New York (1988); *Biocomputing: Informatics and Genome Projects* (Smith, D. W., ed.) Academic Press, New York (1993); *Computer Analysis of Sequence Data, Part I* (Griffin, A. M., and Griffin, H. G., eds.) Humana Press, New Jersey (1994); *Sequence Analysis in Molecular Biology* (von Heinje, G., ed.) Academic Press (1987); and *Sequence Analysis Primer* (Gribskov, M. and Devereux, J., eds.) Stockton Press, New York (1991). Preferred methods to determine identity are designed to give the best match between the sequences tested. Methods to determine identity and similarity are codified in publicly available computer programs. Sequence alignments and percent identity calculations may be performed using the Megalign program of the LASERGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). Multiple alignment of the sequences may be performed using the Clustal method of alignment (Higgins and Sharp (1989) *CABIOS*. 5:151-153) with the default parameters (GAP PENALTY=10, GAP LENGTH PENALTY=10). Default parameters for pairwise alignments using the Clustal method may be selected: KTUPLE 1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5.

- The term "sequence analysis software" refers to any computer algorithm or software program that is useful for the analysis of nucleotide or amino acid sequences. "Sequence analysis software" may be commercially available or independently developed. Typical sequence analysis software will include but is not limited to the GCG suite of programs (Wisconsin Package Version 9.0, Genetics Computer Group (GCG), Madison, WI), BLASTP, BLASTN, BLASTX (Altschul et al., *J. Mol. Biol.* 215:403-410 (1990), and DNASTAR (DNASTAR, Inc. 1228 S. Park St. Madison, WI 53715 USA). Within the context of this application it will be understood that where sequence analysis software is used for analysis, that the results of the analysis will be based on the "default values" of the program referenced, unless otherwise specified. As used herein "default values" will mean any set of values or parameters which originally load with the software when first initialized.

- "Synthetic genes" can be assembled from oligonucleotide building blocks that are chemically synthesized using procedures known to those skilled in the art. These building blocks are ligated and annealed to form gene segments that are then enzymatically assembled to construct the entire gene. "Chemically synthesized", as related to a sequence of DNA, means that the component nucleotides were assembled *in vitro*. Manual chemical synthesis of DNA may be accomplished using well established procedures, or automated chemical synthesis can

be performed using one of a number of commercially available machines. Accordingly, the genes can be tailored for optimal gene expression based on optimization of nucleotide sequence to reflect the codon bias of the host cell. The skilled artisan appreciates the likelihood of successful gene expression if codon usage is biased towards those codons favored by the host.

- 5 Determination of preferred codons can be based on a survey of genes derived from the host cell where sequence information is available.

#### GENE EXPRESSION MODULATION SYSTEM OF THE INVENTION

- Applicants have now shown that separating the transactivation and DNA binding domains by placing them on two different proteins results in greatly reduced background activity in the absence of a ligand and significantly increased activity over background in the presence of a ligand. Applicants' improved gene expression system comprises two chimeric gene expression; the first encoding a DNA binding domain fused to a nuclear receptor polypeptide and the second encoding a transactivation domain fused to a nuclear receptor polypeptide. The interaction of the first protein with the second protein effectively tethers the DNA binding domain to the transactivation domain. Since the DNA binding and transactivation domains reside on two different molecules, the background activity in the absence of ligand is greatly reduced.

- In general, the inducible gene expression modulation system of the invention comprises
- 20 a) a first chimeric gene that is capable of being expressed in a host cell comprising a polynucleotide sequence that encodes a first hybrid polypeptide comprising i) a DNA-binding domain that recognizes a response element associated with a gene whose expression is to be modulated; and ii) a ligand binding domain comprising the ligand binding domain from a nuclear receptor; and b) a second chimeric gene that is capable of being expressed in the host
- 25 cell comprising a polynucleotide sequence that encodes a second hybrid polypeptide comprising: i) a transactivation domain; and ii) a ligand binding domain comprising the ligand binding domain from a nuclear receptor other than ultraspiracle (USP); wherein the transactivation domain are from other than EcR, RXR, or USP; and wherein the ligand binding domains from the first hybrid polypeptide and the second hybrid polypeptide are different and
- 30 dimerize.

This two-hybrid system exploits the ability of a pair of interacting proteins to bring the transcription activation domain into a more favorable position relative to the DNA binding domain such that when the DNA binding domain binds to the DNA binding site on the gene,



the transactivation domain more effectively activates the promoter (see, for example, U.S. Patent No. 5,283,173). This two-hybrid system is a significantly improved inducible gene expression modulation system compared to the two systems disclosed in International Patent Applications PCT/US97/05330 and PCT/US98/14215.

5       The ecdysone receptor-based gene expression modulation system of the invention may be either heterodimeric and homodimeric. A functional EcR complex generally refers to a heterodimeric protein complex consisting of two members of the steroid receptor family, an ecdysone receptor protein obtained from various insects, and an ultraspiracle (USP) protein or the vertebrate homolog of USP, retinoid X receptor protein (see Yao, et al. (1993) Nature 366, 10 476-479; Yao, et al., (1992) Cell 71, 63-72). However, the complex may also be a homodimer as detailed below. The functional ecdysteroid receptor complex may also include additional protein(s) such as immunophilins. Additional members of the steroid receptor family of proteins, known as transcriptional factors (such as DHR38 or *betaFTZ-I*), may also be ligand dependent or independent partners for EcR, USP, and/or RXR. Additionally, other cofactors 15 may be required such as proteins generally known as coactivators (also termed adapters or mediators). These proteins do not bind sequence-specifically to DNA and are not involved in basal transcription. They may exert their effect on transcription activation through various mechanisms, including stimulation of DNA-binding of activators, by affecting chromatin structure, or by mediating activator-initiation complex interactions. Examples of such 20 coactivators include RIP140, TIF1, RAP46/Bag-1, ARA70, SRC-1/NCoA-1, TIF2/GRIP/NCoA-2, ACTR/AIB1/RAC3/pCIP as well as the promiscuous coactivator C response element B binding protein, CBP/p300 (for review see Glass et al, Curr. Opin. Cell Biol. 9:222-232, 1997). Also, protein cofactors generally known as corepressors (also known as repressors, silencers, or silencing mediators) may be required to effectively inhibit 25 transcriptional activation in the absence of ligand. These corepressors may interact with the unliganded ecdysone receptor to silence the activity at the response element. Current evidence suggests that binding of ligand changes the conformation of the receptor, which results in release of the corepressor and recruitment of the above described coactivators, thereby abolishing their silencing activity. Examples of corepressors include N-CoR and SMRT (for 30 review, see Horwitz et al. Mol Endocrinol. 10: 1167-1177, 1996). These cofactors may either be endogenous within the cell or organism, or may be added exogenously as transgenes to be expressed in either a regulated or unregulated fashion. Homodimer complexes of the ecdysone receptor protein, USP, or RXR may also be functional under some circumstances.

The ecdysone receptor complex typically includes proteins which are members of the nuclear receptor superfamily wherein all members are characterized by the presence of an amino-terminal transactivation domain, a DNA binding domain ("DBD"), and a ligand binding domain ("LBD") separated from the DBD by a hinge region. As used herein, the term "DNA  
5 binding domain" comprises a minimal peptide sequence of a DNA binding protein, up to the entire length of a DNA binding protein, so long as the DNA binding domain functions to associate with a particular response element. Members of the nuclear receptor superfamily are also characterized by the presence of four or five domains: A/B, C, D, E, and in some members F (see Evans, *Science* 240:889-895 (1988)). The "A/B" domain corresponds to the  
10 transactivation domain, "C" corresponds to the DNA binding domain, "D" corresponds to the hinge region, and "E" corresponds to the ligand binding domain. Some members of the family may also have another transactivation domain on the carboxy-terminal side of the LBD corresponding to "F".

The DBD is characterized by the presence of two cysteine zinc fingers between which  
15 are two amino acid motifs, the P-box and the D-box, which confer specificity for ecdysone response elements. These domains may be either native, modified, or chimeras of different domains of heterologous receptor proteins. This EcR receptor, like a subset of the steroid receptor family, also possesses less well defined regions responsible for heterodimerization properties. Because the domains of EcR, USP, and RXR are modular in nature, the LBD,  
20 DBD, and transactivation domains may be interchanged.

Gene switch systems are known that incorporate components from the ecdysone receptor complex. However, in these known systems, whenever EcR is used it is associated with native or modified DNA binding domains and transactivation domains on the same molecule. USP or RXR are typically used as silent partners. We have now shown that when  
25 DNA binding domains and transactivation domains are on the same molecule the background activity in the absence of ligand is high and that such activity is dramatically reduced when DNA binding domains and transactivation domains are on different molecules, that is, on each of two partners of a heterodimeric or homodimeric complex. This two-hybrid system also provides improved sensitivity to non-steroidal ligands for example, diacylhydrazines, when  
30 compared to steroidal ligands for example, ponasterone A ("PonA") or munisterone A ("MurA"). That is, when compared to steroids, the non-steroidal ligands provide higher activity at a lower concentration. In addition, since transactivation based on EcR gene switches is often cell-line dependent, it is easier to tailor switching system to obtain maximum

transactivation capability for each application. Furthermore, this two-hybrid system avoids some side effects due to overexpression of RXR that often occur when unmodified RXR is used as a switching partner. In this two-hybrid system, native DNA binding and transactivation domains of EcR or RXR are eliminated. As a result, these chimeric molecules  
5 have less chance of interacting with other steroid hormone receptors present in the cell resulting in reduced side effects.

Specifically, Applicants' invention relates to a gene expression modulation system comprising: a) a first gene expression cassette that is capable of being expressed in a host cell, wherein the first gene expression cassette comprises a polynucleotide that encodes a first  
10 polypeptide comprising i) a DNA-binding domain that recognizes a response element associated with a gene whose expression is to be modulated; and ii) a ligand binding domain comprising a ligand binding domain from a nuclear receptor; and b) a second gene expression cassette that is capable of being expressed in the host cell, wherein the second gene expression cassette comprises a polynucleotide sequence that encodes a second polypeptide comprising i) a  
15 transactivation domain; and ii) a ligand binding domain comprising a ligand binding domain from a nuclear receptor other than ultraspiracle (USP); wherein the DNA binding domain and the transactivation domain are from other than EcR, RXR, or USP; wherein the ligand binding domains from the first polypeptide and the second polypeptide are different and dimerize.

The present invention also relates to a gene expression modulation system according to  
20 the present invention further comprising c) a third gene expression cassette comprising: i) the response element to which the DNA-binding domain of the first polypeptide binds; ii) a promoter that is activated by the transactivation domain of the second polypeptide; and iii) the gene whose expression is to be modulated.

In a specific embodiment, the gene whose expression is to be modulated is a  
25 homologous gene with respect to the host cell. In another specific embodiment, the gene whose expression is to be modulated is a heterologous gene with respect to the host cell.

In a specific embodiment, the ligand binding domain of the first polypeptide comprises an ecdysone receptor ligand binding domain.

In another specific embodiment, the ligand binding domain of the first polypeptide  
30 comprises a retinoid X receptor ligand binding domain.

In a specific embodiment, the ligand binding domain of the second polypeptide comprises an ecdysone receptor ligand binding domain.

In another specific embodiment, the ligand binding domain of the second polypeptide

comprises a retinoid X receptor ligand binding domain.

In a preferred embodiment, the ligand binding domain of the first polypeptide comprises an ecdysone receptor ligand binding domain, and the ligand binding domain of the second polypeptide comprises a retinoid X receptor ligand binding domain.

- 5 In another preferred embodiment, the ligand binding domain of the first polypeptide is from a retinoid X receptor polypeptide, and the ligand binding domain of the second polypeptide is from an ecdysone receptor polypeptide.

- Preferably, the ligand binding domain is an EcR or RXR related steroid/thyroid hormone nuclear receptor family member ligand binding domain, or analogs, combinations, or  
10 modifications thereof. More preferably, the LBD is from EcR or RXR. Even more preferably, the LBD is from a truncated EcR or RXR. A truncation mutation may be made by any method used in the art, including but not limited to restriction endonuclease digestion/deletion, PCR-mediated/oligonucleotide-directed deletion, chemical mutagenesis, UV strand breakage, and the like.

- 15 Preferably, the EcR is an insect EcR selected from the group consisting of a Lepidopteran EcR, a Dipteran EcR, an Arthropod EcR, a Homopteran EcR and a Hemipteran EcR. More preferably, the EcR for use is a spruce budworm *Choristoneura fumiferana* EcR ("CfEcR"), a *Tenebrio molitor* EcR ("TmEcR"), a *Manduca sexta* EcR ("MsEcR"), a *Heliothies virescens* EcR ("HvEcR"), a silk moth *Bombyx mori* EcR ("BmEcR"), a fruit fly  
20 *Drosophila melanogaster* EcR ("DmEcR"), a mosquito *Aedes aegypti* EcR ("AaEcR"), a blowfly *Lucilia capitata* EcR ("LcEcR"), a Mediterranean fruit fly *Ceratitidis capitata* EcR ("CcEcR"), a locust *Locusta migratoria* EcR ("LmEcR"), an aphid *Myzus persicae* EcR ("MpEcR"), a fiddler crab *Uca pugilator* EcR ("UpEcR"), or an ixodid tick *Amblyomma americanum* EcR ("AmaEcR"). Even more preferably, the LBD is from spruce budworm  
25 (*Choristoneura fumiferana*) EcR ("CfEcR") or fruit fly *Drosophila melanogaster* EcR ("DmEcR").

- Preferably, the LBD is from a truncated insect EcR. The insect EcR polypeptide truncation comprises a deletion of at least 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55,  
60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155,  
30 160, 165, 170, 175, 180, 185, 190, 195, 200, 205, 210, 215, 220, 225, 230, 235, 240, 245, 250, 255, 260, or 265 amino acids. More preferably, the insect EcR polypeptide truncation comprises a deletion of at least a partial polypeptide domain. Even more preferably, the insect EcR polypeptide truncation comprises a deletion of at least an entire polypeptide domain. In a

specific embodiment, the insect EcR polypeptide truncation comprises a deletion of at least an A/B-domain deletion, a C-domain deletion, a D-domain deletion, an E-domain deletion, an F-domain deletion, an A/B/C-domains deletion, an A/B/1/2-C-domains deletion, an A/B/C/D-domains deletion, an A/B/C/D/F-domains deletion, an A/B/F-domains, and an A/B/C/F-

5 domains deletion. A combination of several complete and/or partial domain deletions may also be performed.

In a preferred embodiment, the ecdysone receptor ligand binding domain is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ

10 SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.

In another preferred embodiment, the ecdysone receptor ligand binding domain comprises a polypeptide sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, and SEQ ID NO: 20.

15 Preferably, the RXR polypeptide is a mouse *Mus musculus* RXR ("MmRXR") or a human *Homo sapiens* RXR ("HsRXR"). The RXR polypeptide may be an RXR $\alpha$ , RXR $\beta$ , or RXR $\gamma$  isoform.

Preferably, the LBD is from a truncated RXR. The RXR polypeptide truncation comprises a deletion of at least 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70,

20 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 205, 210, 215, 220, 225, 230, 235, 240, 245, 250, 255, 260, or 265 amino acids. More preferably, the RXR polypeptide truncation comprises a deletion of at least a partial polypeptide domain. Even more preferably, the RXR polypeptide truncation comprises a deletion of at least an entire polypeptide domain. In a specific

25 embodiment, the RXR polypeptide truncation comprises a deletion of at least an A/B-domain deletion, a C-domain deletion, a D-domain deletion, an E-domain deletion, an F-domain deletion, an A/B/C-domains deletion, an A/B/1/2-C-domains deletion, an A/B/C/D-domains deletion, an A/B/C/D/F-domains deletion, an A/B/F-domains, and an A/B/C/F-domains deletion. A combination of several complete and/or partial domain deletions may also be

30 performed.

In a preferred embodiment, the retinoid X receptor ligand binding domain is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ

ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30.

In another preferred embodiment, the retinoid X receptor ligand binding domain comprises a polypeptide sequence selected from the group consisting of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 38, SEQ ID NO: 39, and SEQ ID NO: 40.

For purposes of this invention EcR and RXR also include synthetic and chimeric EcR and RXR and their homologs.

The DNA binding domain can be any DNA binding domain with a known response element, including synthetic and chimeric DNA binding domains, or analogs, combinations, or modifications thereof. Preferably, the DBD is a GAL4 DBD, a LexA DBD, a transcription factor DBD, a steroid/thyroid hormone nuclear receptor superfamily member DBD, a bacterial LacZ DBD, or a yeast put DBD. More preferably, the DBD is a GAL4 DBD [SEQ ID NO: 41 (polynucleotide) or SEQ ID NO: 42 (polypeptide)] or a LexA DBD [(SEQ ID NO: 43 (polynucleotide) or SEQ ID NO: 44 (polypeptide)].

The transactivation domain (abbreviated "AD" or "TA") may be any steroid/thyroid hormone nuclear receptor AD, synthetic or chimeric AD, polyglutamine AD, basic or acidic amino acid AD, a VP16 AD, a GAL4 AD, an NF- $\kappa$ B AD, a BP64 AD, or an analog, combination, or modification thereof. Preferably, the AD is a synthetic or chimeric AD, or is obtained from a VP16, GAL4, or NF- $\kappa$ B. Most preferably, the AD is a VP16 AD [SEQ ID NO: 45 (polynucleotide) or SEQ ID NO: 46 (polypeptide)].

The response element ("RE") may be any response element with a known DNA binding domain, or an analog, combination, or modification thereof. Preferably, the RE is an RE from GAL4 ("GAL4RE"), LexA, a steroid/thyroid hormone nuclear receptor RE, or a synthetic RE that recognizes a synthetic DNA binding domain. More preferably, the RE is a GAL4RE comprising a polynucleotide sequence of SEQ ID NO: 47 or a LexA 8X operon comprising a polynucleotide sequence of SEQ ID NO: 48. Preferably, the first hybrid protein is substantially free of a transactivation domain and the second hybrid protein is substantially free of a DNA binding domain. For purposes of this invention, "substantially free" means that the protein in question does not contain a sufficient sequence of the domain in question to provide activation or binding activity.

The ligands for use in the present invention as described below, when combined with the ligand binding domain of an EcR, USP, RXR, or another polypeptide which in turn are bound to the response element linked to a gene, provide the means for external temporal

regulation of expression of the gene. The binding mechanism or the order in which the various components of this invention bind to each other, that is, ligand to receptor, first polypeptide to response element, second polypeptide to promoter, etc., is not critical. Binding of the ligand to the ligand binding domains of an EcR, USP, RXR, or another protein, enables expression or  
5 suppression of the gene. This mechanism does not exclude the potential for ligand binding to EcR, USP, or RXR, and the resulting formation of active homodimer complexes (e.g. EcR+EcR or USP+USP). Preferably, one or more of the receptor domains can be varied producing a chimeric gene switch. Typically, one or more of the three domains, DBD, LBD, and transactivation domain, may be chosen from a source different than the source of the other  
10 domains so that the chimeric genes and the resulting hybrid proteins are optimized in the chosen host cell or organism for transactivating activity, complementary binding of the ligand, and recognition of a specific response element. In addition, the response element itself can be modified or substituted with response elements for other DNA binding protein domains such as the GAL-4 protein from yeast (see Sadowski, et al. (1988) *Nature*, 335:563-564) or LexA  
15 protein from *E. coli* (see Brent and Ptashne (1985), *Cell*, 43:729-736), or synthetic response elements specific for targeted interactions with proteins designed, modified, and selected for such specific interactions (see, for example, Kim, et al. (1997), *Proc. Natl. Acad. Sci., USA*, 94:3616-3620) to accommodate chimeric receptors. Another advantage of chimeric systems is that they allow choice of a promoter used to drive the gene expression according to a desired  
20 end result. Such double control can be particularly important in areas of gene therapy, especially when cytotoxic proteins are produced, because both the timing of expression as well as the cells wherein expression occurs can be controlled. When genes, operatively linked to a suitable promoter, are introduced into the cells of the subject, expression of the exogenous genes is controlled by the presence of the system of this invention. Promoters may be  
25 constitutively or inducibly regulated or may be tissue-specific (that is, expressed only in a particular type of cells) or specific to certain developmental stages of the organism.

#### GENE EXPRESSION CASSETTES OF THE INVENTION

The novel ecdysone receptor-based inducible gene expression system of the invention  
30 comprises a novel gene expression cassette that is capable of being expressed in a host cell, wherein the gene expression cassette comprises a polynucleotide encoding a hybrid polypeptide. Thus, Applicants' invention also provides novel gene expression cassettes for use in the gene expression system of the invention.

Specifically, the present invention provides a gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide. The hybrid polypeptide comprises either 1) a DNA-binding domain that recognizes a response element and a ligand binding domain of a nuclear receptor or 2) a transactivation domain and a ligand binding domain of a nuclear receptor, wherein the transactivation domain is from a nuclear receptor other than an EcR, an RXR, or a USP.

In a specific embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a DNA-binding domain that recognizes a response element and an ecdysone receptor ligand binding domain, wherein the DNA binding domain is from a nuclear receptor other than an ecdysone receptor.

In another specific embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a DNA-binding domain that recognizes a response element and a retinoid X receptor ligand binding domain, wherein the DNA binding domain is from a nuclear receptor other than a retinoid X receptor.

The DNA binding domain can be any DNA binding domain with a known response element, including synthetic and chimeric DNA binding domains, or analogs, combinations, or modifications thereof. Preferably, the DBD is a GAL4 DBD, a LexA DBD, a transcription factor DBD, a steroid/thyroid hormone nuclear receptor superfamily member DBD, a bacterial LacZ DBD, or a yeast put DBD. More preferably, the DBD is a GAL4 DBD [SEQ ID NO: 41 (polynucleotide) or SEQ ID NO: 42 (polypeptide)] or a LexA DBD [(SEQ ID NO: 43 (polynucleotide) or SEQ ID NO: 44 (polypeptide)].

In another specific embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a transactivation domain and an ecdysone receptor ligand binding domain, wherein the transactivation domain is from a nuclear receptor other than an ecdysone receptor.

In another specific embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a transactivation domain and a retinoid X receptor ligand binding domain, wherein the transactivation domain is from a nuclear receptor other than a retinoid X receptor.

The transactivation domain (abbreviated "AD" or "TA") may be any steroid/thyroid hormone nuclear receptor AD, synthetic or chimeric AD, polyglutamine AD, basic or acidic amino acid AD, a VP16 AD, a GAL4 AD, an NF- $\kappa$ B AD, a BP64 AD, or an analog, combination, or modification thereof. Preferably, the AD is a synthetic or chimeric AD, or is



obtained from a VP16, GAL4, or NF-kB. Most preferably, the AD is a VP16 AD [SEQ ID NO: 45 (polynucleotide) or SEQ ID NO: 46 (polypeptide)].

Preferably, the ligand binding domain is an EcR or RXR related steroid/thyroid hormone nuclear receptor family member ligand binding domain, or analogs, combinations, or 5 modifications thereof. More preferably, the LBD is from EcR or RXR. Even more preferably, the LBD is from a truncated EcR or RXR.

Preferably, the EcR is an insect EcR selected from the group consisting of a Lepidopteran EcR, a Dipteran EcR, an Arthropod EcR, a Homopteran EcR and a Hemipteran EcR. More preferably, the EcR for use is a spruce budworm *Choristoneura fumiferana* EcR 10 ("CfEcR"), a *Tenebrio molitor* EcR ("TmEcR"), a *Manduca sexta* EcR ("MsEcR"), a *Heliothis virescens* EcR ("HvEcR"), a silk moth *Bombyx mori* EcR ("BmEcR"), a fruit fly *Drosophila melanogaster* EcR ("DmEcR"), a mosquito *Aedes aegypti* EcR ("AaEcR"), a blowfly *Lucilia capitata* EcR ("LcEcR"), a Mediterranean fruit fly *Ceratitis capitata* EcR ("CcEcR"), a locust *Locusta migratoria* EcR ("LmEcR"), an aphid *Myzus persicae* EcR 15 ("MpEcR"), a fiddler crab *Uca pugilator* EcR ("UpEcR"), or an ixodid tick *Amblyomma americanum* EcR ("AmaEcR"). Even more preferably, the LBD is from spruce budworm (*Choristoneura fumiferana*) EcR ("CfEcR") or fruit fly *Drosophila melanogaster* EcR ("DmEcR").

Preferably, the LBD is from a truncated insect EcR. The insect EcR polypeptide 20 truncation comprises a deletion of at least 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 205, 210, 215, 220, 225, 230, 235, 240, 245, 250, 255, 260, or 265 amino acids. More preferably, the insect EcR polypeptide truncation comprises a deletion of at least a partial polypeptide domain. Even more preferably, the insect 25 EcR polypeptide truncation comprises a deletion of at least an entire polypeptide domain. In a specific embodiment, the insect EcR polypeptide truncation comprises a deletion of at least an A/B-domain deletion, a C-domain deletion, a D-domain deletion, an E-domain deletion, an F-domain deletion, an A/B/C-domains deletion, an A/B/1/2-C-domains deletion, an A/B/C/D-domains deletion, an A/B/C/D/F-domains deletion, an A/B/F-domains, and an A/B/C/F- 30 domains deletion. A combination of several complete and/or partial domain deletions may also be performed.

In a preferred embodiment, the ecdysone receptor ligand binding domain is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ

ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.

In another preferred embodiment, the ecdysone receptor ligand binding domain comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, and SEQ ID NO: 20.

Preferably, the RXR polypeptide is a mouse *Mus musculus* RXR ("MmRXR") or a human *Homo sapiens* RXR ("HsRXR"). The RXR polypeptide may be an RXR $\alpha$ , RXR $\beta$ , or RXR $\gamma$  isoform.

10 Preferably, the LBD is from a truncated RXR. The RXR polypeptide truncation comprises a deletion of at least 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 205, 210, 215, 220, 225, 230, 235, 240, 245, 250, 255, 260, or 265 amino acids. More preferably, the RXR polypeptide truncation comprises a  
15 deletion of at least a partial polypeptide domain. Even more preferably, the RXR polypeptide truncation comprises a deletion of at least an entire polypeptide domain. In a specific embodiment, the RXR polypeptide truncation comprises a deletion of at least an A/B-domain deletion, a C-domain deletion, a D-domain deletion, an E-domain deletion, an F-domain deletion, an A/B/C-domains deletion, an A/B/1/2-C-domains deletion, an A/B/C/D-domains  
20 deletion, an A/B/C/D/F-domains deletion, an A/B/F-domains, and an A/B/C/F-domains deletion. A combination of several complete and/or partial domain deletions may also be performed.

In a preferred embodiment, the retinoid X receptor ligand binding domain is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of  
25 SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30.

In another preferred embodiment, the retinoid X receptor ligand binding domain comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID  
30 NO: 37, SEQ ID NO: 38, SEQ ID NO: 39, and SEQ ID NO: 40.

In a preferred embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a DNA-binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of a GAL4 DBD (SEQ ID NO: 41) or a LexA

DBD (SEQ ID NO: 43) and an ecdysone receptor ligand binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.

5 In another preferred embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a DNA-binding domain comprising a polypeptide sequence selected from the group consisting of a GAL4 DBD (SEQ ID NO: 42) or a LexA DBD (SEQ ID NO: 44) and an ecdysone receptor ligand binding domain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ  
10 ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, and SEQ ID NO: 20.

In another preferred embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a DNA-binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of a GAL4 DBD (SEQ ID NO: 41) or  
15 a LexA DBD (SEQ ID NO: 43) and a retinoid X receptor ligand binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30.

In another preferred embodiment, the gene expression cassette encodes a hybrid  
20 polypeptide comprising a DNA-binding domain comprising a polypeptide sequence selected from the group consisting of a GAL4 DBD (SEQ ID NO: 42) or a LexA DBD (SEQ ID NO: 44) and a retinoid X receptor ligand binding domain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 38, SEQ ID  
25 NO: 39, and SEQ ID NO: 40.

In another preferred embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a transactivation domain encoded by a polynucleotide comprising a nucleic acid sequence of SEQ ID NO: 45 and an ecdysone receptor ligand binding domain  
30 encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.

In another preferred embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a transactivation domain comprising a polypeptide sequence of SEQ

ID NO: 46 and an ecdysone receptor ligand binding domain comprising a polypeptide sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, and SEQ ID NO: 20.

5 In another preferred embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a transactivation domain encoded by a polynucleotide comprising a nucleic acid sequence of SEQ ID NO: 45 and a retinoid X receptor ligand binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID  
10 NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30.

In another preferred embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a transactivation domain comprising a polypeptide sequence of SEQ ID NO: 46 and a retinoid X receptor ligand binding domain comprising an amino acid sequence  
15 selected from the group consisting of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 38, SEQ ID NO: 39, and SEQ ID NO: 40.

For purposes of this invention EcR and RXR also include synthetic and chimeric EcR and RXR and their homologs.

20

#### POLYNUCLEOTIDES OF THE INVENTION

The novel ecdysone receptor-based inducible gene expression system of the invention comprises a gene expression cassette comprising a polynucleotide that encodes a truncated EcR or RXR polypeptide comprising a truncation mutation and is useful in methods of  
25 modulating the expression of a gene within a host cell.

Thus, the present invention also relates to a polynucleotide that encodes an EcR or RXR polypeptide comprising a truncation mutation. Specifically, the present invention relates to an isolated polynucleotide encoding an EcR or RXR polypeptide comprising a truncation mutation that affects ligand binding activity or ligand sensitivity.

30 Preferably, the truncation mutation results in a polynucleotide that encodes a truncated EcR polypeptide or a truncated RXR polypeptide comprising a deletion of at least 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 205, 210,

215, 220, 225, 230, 235, 240, 245, 250, 255, 260, or 265 amino acids. More preferably, the EcR or RXR polypeptide truncation comprises a deletion of at least a partial polypeptide domain. Even more preferably, the EcR or RXR polypeptide truncation comprises a deletion of at least an entire polypeptide domain. In a specific embodiment, the EcR or RXR

5 polypeptide truncation comprises a deletion of at least an A/B-domain deletion, a C-domain deletion, a D-domain deletion, an E-domain deletion, an F-domain deletion, an A/B/C-domains deletion, an A/B/1/2-C-domains deletion, an A/B/C/D-domains deletion, an A/B/C/D/F-domains deletion, an A/B/F-domains, and an A/B/C/F-domains deletion. A combination of several complete and/or partial domain deletions may also be performed.

10 In a specific embodiment, the EcR polynucleotide according to the invention comprises a polynucleotide sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10. In a specific embodiment, the polynucleotide according to the invention encodes a ecdysone receptor polypeptide comprising an amino acid  
15 sequence selected from the group consisting of SEQ ID NO: 11 (CfEcR-CDEF), SEQ ID NO: 12 (CfEcR-1/2CDEF, which comprises the last 33 carboxy-terminal amino acids of C domain), SEQ ID NO: 13 (CfEcR-DEF), SEQ ID NO: 14 (CfEcR-EF), SEQ ID NO: 15 (CfEcR-DE), SEQ ID NO: 16 (DmEcR-CDEF), SEQ ID NO: 17 (DmEcR-1/2CDEF), SEQ ID NO: 18 (DmEcR-DEF), SEQ ID NO: 19 (DmEcR-EF), and SEQ ID NO: 20 (DmEcR-  
20 DE).

In another specific embodiment, the RXR polynucleotide according to the invention comprises a polynucleotide sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30. In another specific  
25 embodiment, the polynucleotide according to the invention encodes a truncated RXR polypeptide comprising an amino acid sequence consisting of SEQ ID NO: 31 (MmRXR-CDEF), SEQ ID NO: 32 (MmRXR-DEF), SEQ ID NO: 33 (MmRXR-EF), SEQ ID NO: 34 (MmRXR-truncatedEF), SEQ ID NO: 35 (MmRXR-E), SEQ ID NO: 36 (HsRXR-CDEF), SEQ ID NO: 37 (HsRXR-DEF), SEQ ID NO: 38 (HsRXR-EF), SEQ ID NO: 39 (HsRXR-truncated EF), and SEQ ID NO: 40 (HsRXR-E).  
30

In particular, the present invention relates to an isolated polynucleotide encoding an EcR or RXR polypeptide comprising a truncation mutation, wherein the mutation reduces ligand binding activity or ligand sensitivity of the EcR or RXR polypeptide. In a specific

embodiment, the present invention relates to an isolated polynucleotide encoding an EcR or RXR polypeptide comprising a truncation mutation that reduces steroid binding activity or steroid sensitivity of the EcR or RXR polypeptide. In a preferred embodiment, the present invention relates to an isolated polynucleotide encoding an EcR polypeptide comprising a truncation mutation that reduces steroid binding activity or steroid sensitivity of the EcR polypeptide, wherein the polynucleotide comprises a nucleic acid sequence of SEQ ID NO: 3 (CfEcR-DEF), SEQ ID NO: 4 (CfEcR-EF), SEQ ID NO: 8 (DmEcR-DEF), or SEQ ID NO: 9 (DmEcR-EF). In another specific embodiment, the present invention relates to an isolated polynucleotide encoding an EcR or RXR polypeptide comprising a truncation mutation that reduces non-steroid binding activity or non-steroid sensitivity of the EcR or RXR polypeptide. In a preferred embodiment, the present invention relates to an isolated polynucleotide encoding an EcR polypeptide comprising a truncation mutation that reduces non-steroid binding activity or non-steroid sensitivity of the EcR polypeptide, wherein the polynucleotide comprises a nucleic acid sequence of SEQ ID NO: 4 (CfEcR-EF) or SEQ ID NO: 9 (DmEcR-EF).

The present invention also relates to an isolated polynucleotide encoding an EcR or RXR polypeptide comprising a truncation mutation, wherein the mutation enhances ligand binding activity or ligand sensitivity of the EcR or RXR polypeptide. In a specific embodiment, the present invention relates to an isolated polynucleotide encoding an EcR or RXR polypeptide comprising a truncation mutation that enhances steroid binding activity or steroid sensitivity of the EcR or RXR polypeptide. In another specific embodiment, the present invention relates to an isolated polynucleotide encoding an EcR or RXR polypeptide comprising a truncation mutation that enhances non-steroid binding activity or non-steroid sensitivity of the EcR or RXR polypeptide. In a preferred embodiment, the present invention relates to an isolated polynucleotide encoding an EcR polypeptide comprising a truncation mutation that enhances non-steroid binding activity or non-steroid sensitivity of the EcR polypeptide, wherein the polynucleotide comprises a nucleic acid sequence of SEQ ID NO: 3 (CfEcR-DEF) or SEQ ID NO: 8 (DmEcR-DEF).

The present invention also relates to an isolated polynucleotide encoding a retinoid X receptor polypeptide comprising a truncation mutation that increases ligand sensitivity of a heterodimer comprising the mutated retinoid X receptor polypeptide and a dimerization partner. Preferably, the isolated polynucleotide encoding a retinoid X receptor polypeptide comprising a truncation mutation that increases ligand sensitivity of a heterodimer comprises a polynucleotide sequence selected from the group consisting of SEQ ID NO: 23 (MmRXR-EF),

SEQ ID NO: 24 (MmRXR-truncatedEF), SEQ ID NO: 28 (HsRXR-EF), or SEQ ID NO: 29 (HsRXR-truncated EF). In a specific embodiment, the dimerization partner is an ecdysone receptor polypeptide. Preferably, the dimerization partner is a truncated EcR polypeptide. More preferably, the dimerization partner is an EcR polypeptide in which domains A/B/C have  
 5 been deleted. Even more preferably, the dimerization partner is an EcR polypeptide comprising an amino acid sequence of SEQ ID NO: 13 (CfEcR-DEF) or SEQ ID NO: 18 (DmEcR-DEF).

### POLYPEPTIDES OF THE INVENTION

The novel ecdysone receptor-based inducible gene expression system of the invention  
 10 comprises a polynucleotide that encodes a truncated EcR or RXR polypeptide and is useful in methods of modulating the expression of a gene within a host cell. Thus, the present invention also relates to an isolated truncated EcR or RXR polypeptide encoded by a polynucleotide or a gene expression cassette according to the invention. Specifically, the present invention relates to an isolated truncated EcR or RXR polypeptide comprising a truncation mutation that affects  
 15 ligand binding activity or ligand sensitivity encoded by a polynucleotide according to the invention.

The present invention also relates to an isolated truncated EcR or RXR polypeptide comprising a truncation mutation. Specifically, the present invention relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that affects ligand binding activity or  
 20 ligand sensitivity.

Preferably, the truncation mutation results in a truncated EcR polypeptide or a truncated RXR polypeptide comprising a deletion of at least 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 205, 210, 215, 220, 225,  
 25 230, 235, 240, 245, 250, 255, 260, or 265 amino acids. More preferably, the EcR or RXR polypeptide truncation comprises a deletion of at least a partial polypeptide domain. Even more preferably, the EcR or RXR polypeptide truncation comprises a deletion of at least an entire polypeptide domain. In a specific embodiment, the EcR or RXR polypeptide truncation comprises a deletion of at least an A/B-domain deletion, a C-domain deletion, a D-domain  
 30 deletion, an E-domain deletion, an F-domain deletion, an A/B/C-domains deletion, an A/B/1/2-C-domains deletion, an A/B/C/D-domains deletion, an A/B/C/D/F-domains deletion, an A/B/F-domains, and an A/B/C/F-domains deletion. A combination of several complete and/or partial domain deletions may also be performed.

In a preferred embodiment, the isolated truncated ecdysone receptor polypeptide is encoded by a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO: 1 (CfEcR-CDEF), SEQ ID NO: 2 (CfEcR-1/2CDEF), SEQ ID NO: 3 (CfEcR-DEF), SEQ ID NO: 4 (CfEcR-EF), SEQ ID NO: 5 (CfEcR-DE), SEQ ID NO: 6 (DmEcR-CDEF), SEQ ID NO: 7 (DmEcR-1/2CDEF), SEQ ID NO: 8 (DmEcR-DEF), SEQ ID NO: 9 (DmEcR-EF), and SEQ ID NO: 10 (DmEcR-DE). In another preferred embodiment, the isolated truncated ecdysone receptor polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 11 (CfEcR-CDEF), SEQ ID NO: 12 (CfEcR-1/2CDEF), SEQ ID NO: 13 (CfEcR-DEF), SEQ ID NO: 14 (CfEcR-EF), SEQ ID NO: 15 (CfEcR-DE), SEQ ID NO: 16 (DmEcR-CDEF), SEQ ID NO: 17 (DmEcR-1/2CDEF), SEQ ID NO: 18 (DmEcR-DEF), SEQ ID NO: 19 (DmEcR-EF), and SEQ ID NO: 20 (DmEcR-DE).

In a preferred embodiment, the isolated truncated RXR polypeptide is encoded by a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO: 21 (MmRXR-CDEF), SEQ ID NO: 22 (MmRXR-DEF), SEQ ID NO: 23 (MmRXR-EF), SEQ ID NO: 24 (MmRXR-truncatedEF), SEQ ID NO: 25 (MmRXR-E), SEQ ID NO: 26 (HsRXR-CDEF), SEQ ID NO: 27 (HsRXR-DEF), SEQ ID NO: 28 (HsRXR-EF), SEQ ID NO: 29 (HsRXR-truncatedEF) and SEQ ID NO: 30 (HsRXR-E). In another preferred embodiment, the isolated truncated RXR polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 31 (MmRXR-CDEF), SEQ ID NO: 32 (MmRXR-DEF), SEQ ID NO: 33 (MmRXR-EF), SEQ ID NO: 34 (MmRXR-truncatedEF), SEQ ID NO: 35 (MmRXR-E), SEQ ID NO: 36 (HsRXR-CDEF), SEQ ID NO: 37 (HsRXR-DEF), SEQ ID NO: 38 (HsRXR-EF), SEQ ID NO: 39 (HsRXR-truncatedEF), and SEQ ID NO: 40 (HsRXR-E).

The present invention relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that reduces ligand binding activity or ligand sensitivity of the EcR or RXR polypeptide, wherein the polypeptide is encoded by a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO: 1 (CfEcR-CDEF), SEQ ID NO: 2 (CfEcR-1/2CDEF), SEQ ID NO: 3 (CfEcR-DEF), SEQ ID NO: 4 (CfEcR-EF), SEQ ID NO: 5 (CfEcR-DE), SEQ ID NO: 6 (DmEcR-CDEF), SEQ ID NO: 7 (DmEcR-1/2CDEF), SEQ ID NO: 8 (DmEcR-DEF), SEQ ID NO: 9 (DmEcR-EF), SEQ ID NO: 10 (DmEcR-DE), SEQ ID NO: 21 (MmRXR-CDEF), SEQ ID NO: 22 (MmRXR-DEF), SEQ ID NO: 23 (MmRXR-EF), SEQ ID NO: 24 (MmRXR-truncatedEF), SEQ ID NO: 25 (MmRXR-



E), SEQ ID NO: 26 (HsRXR-CDEF), SEQ ID NO: 27 (HsRXR-DEF), SEQ ID NO: 28 (HsRXR-EF), SEQ ID NO: 29 (HsRXR-truncatedEF), and SEQ ID NO: 30 (HsRXR-E).

Thus, the present invention relates to an isolated truncated EcR or RXR polypeptide comprising a truncation mutation that reduces ligand binding activity or ligand sensitivity of  
5 the EcR or RXR polypeptide, wherein the polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 11 (CfEcR-CDEF), SEQ ID NO: 12 (CfEcR-1/2CDEF), SEQ ID NO: 13 (CfEcR-DEF), SEQ ID NO: 14 (CfEcR-EF), SEQ ID NO: 15 (CfEcR-DE), SEQ ID NO: 16 (DmEcR-CDEF), SEQ ID NO: 17 (DmEcR-1/2CDEF), SEQ ID NO: 18 (DmEcR-DEF), SEQ ID NO: 19 (DmEcR-EF), SEQ ID NO: 20 (DmEcR-  
10 DE), SEQ ID NO: 31 (MmRXR-CDEF), SEQ ID NO: 32 (MmRXR-DEF), SEQ ID NO: 33 (MmRXR-EF), SEQ ID NO: 34 (MmRXR-truncatedEF), SEQ ID NO: 35 (MmRXR-E), SEQ ID NO: 36 (HsRXR-CDEF), SEQ ID NO: 37 (HsRXR-DEF), SEQ ID NO: 38 (HsRXR-EF), SEQ ID NO: 39 (HsRXR-truncatedEF), and SEQ ID NO: 40 (HsRXR-E).

In a specific embodiment, the present invention relates to an isolated EcR or RXR  
15 polypeptide comprising a truncation mutation that reduces steroid binding activity or steroid sensitivity of the EcR or RXR polypeptide. In a preferred embodiment, the present invention relates to an isolated EcR polypeptide comprising a truncation mutation that reduces steroid binding activity or steroid sensitivity of the EcR polypeptide, wherein the EcR polypeptide is encoded by a polynucleotide comprising a nucleic acid sequence of SEQ ID NO: 3 (CfEcR-  
20 DEF), SEQ ID NO: 4 (CfEcR-EF), SEQ ID NO: 8 (DmEcR-DEF), or SEQ ID NO: 9 (DmEcR-EF). Accordingly, the present invention also relates to an isolated truncated EcR or RXR polypeptide comprising a truncation mutation that reduces steroid binding activity or steroid sensitivity of the EcR or RXR polypeptide. In a preferred embodiment, the present invention relates to an isolated EcR polypeptide comprising a truncation mutation that reduces  
25 steroid binding activity or steroid sensitivity of the EcR polypeptide, wherein the EcR polypeptide comprises an amino acid sequence of SEQ ID NO: 13 (CfEcR-DEF), SEQ ID NO: 14 (CfEcR-EF), SEQ ID NO: 18 (DmEcR-DEF), or SEQ ID NO: 19 (DmEcR-EF).

In another specific embodiment, the present invention relates to an isolated EcR or  
RXR polypeptide comprising a truncation mutation that reduces non-steroid binding activity or  
30 non-steroid sensitivity of the EcR or RXR polypeptide. In a preferred embodiment, the present invention relates to an isolated EcR polypeptide comprising a truncation mutation that reduces non-steroid binding activity or non-steroid sensitivity of the EcR polypeptide, wherein the EcR polypeptide is encoded by a polynucleotide comprising a nucleic acid sequence of SEQ ID NO:

4 (CfEcR-EF) or SEQ ID NO: 9 (DmEcR-EF). Accordingly, the present invention also relates to an isolated truncated EcR or RXR polypeptide comprising a truncation mutation that reduces non-steroid binding activity or steroid sensitivity of the EcR or RXR polypeptide. In a preferred embodiment, the present invention relates to an isolated EcR polypeptide comprising  
 5 a truncation mutation that reduces non-steroid binding activity or non-steroid sensitivity of the EcR polypeptide, wherein the EcR polypeptide comprises an amino acid sequence of SEQ ID NO: 14 (CfEcR-EF) or SEQ ID NO: 19 (DmEcR-EF).

In particular, the present invention relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that enhances ligand binding activity or ligand sensitivity of  
 10 the EcR or RXR polypeptide, wherein the polypeptide is encoded by a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO: 1 (CfEcR-CDEF), SEQ ID NO: 2 (CfEcR-1/2CDEF), SEQ ID NO: 3 (CfEcR-DEF), SEQ ID NO: 4 (CfEcR-EF), SEQ ID NO: 5 (CfEcR-DE), SEQ ID NO: 6 (DmEcR-CDEF), SEQ ID NO: 7 (DmEcR-1/2CDEF), SEQ ID NO: 8 (DmEcR-DEF), SEQ ID NO: 9 (DmEcR-EF),  
 15 SEQ ID NO: 10 (DmEcR-DE), SEQ ID NO: 21 (MmRXR-CDEF), SEQ ID NO: 22 (MmRXR-DEF), SEQ ID NO: 23 (MmRXR-EF), SEQ ID NO: 24 (MmRXR-truncatedEF), SEQ ID NO: 25 (MmRXR-E), SEQ ID NO: 26 (HsRXR-CDEF), SEQ ID NO: 27 (HsRXR-DEF), SEQ ID NO: 28 (HsRXR-EF), SEQ ID NO: 29 (HsRXR-truncated EF), and SEQ ID NO: 30 (HsRXR-E).

20 The present invention relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that enhances ligand binding activity or ligand sensitivity of the EcR or RXR polypeptide, wherein the polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 11 (CfEcR-CDEF), SEQ ID NO: 12 (CfEcR-1/2CDEF), SEQ ID NO: 13 (CfEcR-DEF), SEQ ID NO: 14 (CfEcR-EF), SEQ ID NO: 15 (CfEcR-DE),  
 25 SEQ ID NO: 16 (DmEcR-CDEF), SEQ ID NO: 17 (DmEcR-1/2CDEF), SEQ ID NO: 18 (DmEcR-DEF), SEQ ID NO: 19 (DmEcR-EF), SEQ ID NO: 20 (DmEcR-DE), SEQ ID NO: 31 (MmRXR-CDEF), SEQ ID NO: 32 (MmRXR-DEF), SEQ ID NO: 33 (MmRXR-EF),  
 30 SEQ ID NO: 34 (MmRXR-truncatedEF), SEQ ID NO: 35 (MmRXR-E), SEQ ID NO: 36 (HsRXR-CDEF), SEQ ID NO: 37 (HsRXR-DEF), SEQ ID NO: 39 (HsRXR-EF), SEQ ID NO: 39 (HsRXR-truncatedEF), and SEQ ID NO: 40 (HsRXR-E).

The present invention relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that enhances ligand binding activity or ligand sensitivity of the EcR or RXR polypeptide. In a specific embodiment, the present invention relates to an isolated EcR or

RXR polypeptide comprising a truncation mutation that enhances steroid binding activity or steroid sensitivity of the EcR or RXR polypeptide. Accordingly, the present invention also relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that enhances steroid binding activity or steroid sensitivity of the EcR or RXR polypeptide.

- 5 In another specific embodiment, the present invention relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that enhances non-steroid binding activity or non-steroid sensitivity of the EcR or RXR polypeptide. In a preferred embodiment, the present invention relates to an isolated EcR polypeptide comprising a truncation mutation that enhances non-steroid binding activity or non-steroid sensitivity of the EcR polypeptide, wherein
- 10 the EcR polypeptide is encoded by a polynucleotide comprising a nucleic acid sequence of SEQ ID NO: 3 (CfEcR-DEF) or SEQ ID NO: 8 (DmEcR-DEF). Accordingly, the present invention also relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that enhances non-steroid binding activity or steroid sensitivity of the EcR or RXR polypeptide. In a preferred embodiment, the present invention relates to an isolated EcR polypeptide comprising
- 15 a truncation mutation that enhances non-steroid binding activity or non-steroid sensitivity of the EcR polypeptide, wherein the EcR polynucleotide comprises an amino acid sequence of SEQ ID NO: 13 (CfEcR-DEF) or SEQ ID NO: 18 (DmEcR-DEF).

- The present invention also relates to an isolated retinoid X receptor polypeptide comprising a truncation mutation that increases ligand sensitivity of a heterodimer comprising
- 20 the mutated retinoid X receptor polypeptide and a dimerization partner. Preferably, the isolated retinoid X receptor polypeptide comprising a truncation mutation that increases ligand sensitivity of a heterodimer is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 23 (MmRXR-EF), SEQ ID NO: 24 (MmRXR-truncatedEF), SEQ ID NO: 28 (HsRXR-EF), or SEQ ID NO: 29 (HsRXR-
- 25 truncatedEF). More preferably, the isolated polynucleotide encoding a retinoid X receptor polypeptide comprising a truncation mutation that increases ligand sensitivity of a heterodimer comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 33 (MmRXR-EF), SEQ ID NO: 34 (MmRXR-truncatedEF), SEQ ID NO: 38 (HsRXR-EF), or SEQ ID NO: 39 (HsRXR-truncatedEF).

- 30 In a specific embodiment, the dimerization partner is an ecdysone recaptor polypeptide. Preferably, the dimerization partner is a truncated EcR polypeptide. More preferably, the dimerization partner is an EcR polypeptide in which domains A/B/C have been deleted. Even more preferably, the dimerization partner is an EcR polypeptide comprising an amino acid

sequence of SEQ ID NO: 13 (CfEcR-DEF) or SEQ ID NO: 18 (DmEcR-DEF).

#### METHOD OF MODULATING GENE EXPRESSION OF THE INVENTION


Applicants' invention also relates to methods of modulating gene expression in a host  
5 cell using a gene expression modulation system according to the invention. Specifically,  
Applicants' invention provides a method of modulating the expression of a gene in a host cell  
comprising the steps of: a) introducing into the host cell a gene expression modulation system  
according to the invention; and b) introducing into the host cell a ligand that independently  
combines with the ligand binding domains of the first polypeptide and the second polypeptide  
10 of the gene expression modulation system; wherein the gene to be expressed is a component of  
a gene expression cassette comprising: i) a response element comprising a domain to which the  
DNA binding domain of the first polypeptide binds; ii) a promoter that is activated by the  
transactivation domain of the second polypeptide; and iii) a gene whose expression is to be  
modulated, whereby a complex is formed comprising the ligand, the first polypeptide of the  
15 gene expression modulation system and the second polypeptide of the gene expression  
modulation system, and whereby the complex modulates expression of the gene in the host cell.

Genes of interest for expression in a host cell using Applicants' methods may be  
endogenous genes or heterologous genes. Nucleic acid or amino acid sequence information for  
a desired gene or protein can be located in one of many public access databases, for example,  
20 GENBANK, EMBL, Swiss-Prot, and PIR, or in many biology related journal publications.  
Thus, those skilled in the art have access to nucleic acid sequence information for virtually all  
known genes. Such information can then be used to construct the desired constructs for the  
insertion of the gene of interest within the gene expression cassettes used in Applicants'  
methods described herein.

25 Examples of genes of interest for expression in a host cell using Applicants' methods  
include, but are not limited to: antigens produced in plants as vaccines, enzymes like alpha-  
amylase, phytase, glucanase, and xylanase, genes for resistance against insects, nematodes, fungi,  
bacteria, viruses, and abiotic stresses, nutraceuticals, pharmaceuticals, vitamins, genes for  
modifying amino acid content, herbicide resistance, cold, drought, and heat tolerance, industrial  
30 products, oils, protein, carbohydrates, antioxidants, male sterile plants, flowers, fuels, other  
output traits, genes encoding therapeutically desirable polypeptides or products, such as genes  
that can provide, modulate, alleviate, correct and/or restore polypeptides important in treating a  
condition, a disease, a disorder, a dysfunction, a genetic defect, and the like.

Acceptable ligands are any that modulate expression of the gene when binding of the DNA binding domain of the two hybrid system to the response element in the presence of the ligand results in activation or suppression of expression of the genes. Preferred ligands include ponasterone, muristerone A, N,N'-diacylhydrazines such as those disclosed in U. S. Patents 5 No. 6,013,836; 5,117,057; 5,530,028; and 5,378,726; dibenzoylalkyl cyanohydrazines such as those disclosed in European Application No. 461,809; N-alkyl-N,N'-diaroylhydrazines such as those disclosed in U. S. Patent No. 5,225,443; N-acyl-N-alkylcarbonylhydrazines such as those disclosed in European Application No. 234,994; N-aryl-N-alkyl-N'-aroylhydrazines such as those described in U. S. Patent No. 4,985,461; each of which is incorporated herein by reference and other similar materials including 3,5-di-tert-butyl-4-hydroxy-N-isobutylbenzamide, 8-O-acetylharpagide, and the like.

Preferably, the ligand for use in Applicants' method of modulating expression of gene is a compound of the formula:



15 wherein:

E is a (C<sub>4</sub>-C<sub>6</sub>)alkyl containing a tertiary carbon or a cyano(C<sub>3</sub>-C<sub>5</sub>)alkyl containing a tertiary carbon;

R<sup>1</sup> is H, Me, Et, i-Pr, F, formyl, CF<sub>3</sub>, CHF<sub>2</sub>, CHCl<sub>2</sub>, CH<sub>2</sub>F, CH<sub>2</sub>Cl, CH<sub>2</sub>OH, CH<sub>2</sub>OMe, CH<sub>2</sub>CN, CN, C<sup>o</sup>CH, 1-propynyl, 2-propynyl, vinyl, OH, OMe, OEt, cyclopropyl, CF<sub>2</sub>CF<sub>3</sub>, CH=CHCN, allyl, azido, SCN, or SCHF<sub>2</sub>;

R<sup>2</sup> is H, Me, Et, n-Pr, i-Pr, formyl, CF<sub>3</sub>, CHF<sub>2</sub>, CHCl<sub>2</sub>, CH<sub>2</sub>F, CH<sub>2</sub>Cl, CH<sub>2</sub>OH, CH<sub>2</sub>OMe, CH<sub>2</sub>CN, CN, C<sup>o</sup>CH, 1-propynyl, 2-propynyl, vinyl, Ac, F, Cl, OH, OMe, OEt, O-n-Pr, OAc, NMe<sub>2</sub>, NEt<sub>2</sub>, SMe, SEt, SOCF<sub>3</sub>, OCF<sub>2</sub>CF<sub>2</sub>H, COEt, cyclopropyl, CF<sub>2</sub>CF<sub>3</sub>, CH=CHCN, allyl, azido, OCF<sub>3</sub>, OCHF<sub>2</sub>, O-i-Pr, SCN, SCHF<sub>2</sub>, SOMe, NH-CN, or joined with R<sup>3</sup> and the phenyl carbons to which R<sup>2</sup> and R<sup>3</sup> are attached to form an ethylenedioxy, a dihydrofuryl ring with the oxygen adjacent to a phenyl carbon, or a dihydropyryl ring with the oxygen adjacent to a phenyl carbon;

R<sup>3</sup> is H, Et, or joined with R<sup>2</sup> and the phenyl carbons to which R<sup>2</sup> and R<sup>3</sup> are attached to

form an ethylenedioxy, a dihydrofuryl ring with the oxygen adjacent to a phenyl carbon, or a dihydropyryl ring with the oxygen adjacent to a phenyl carbon; R<sup>4</sup>, R<sup>5</sup>, and R<sup>6</sup> are independently H, Me, Et, F, Cl, Br, formyl, CF<sub>3</sub>, CHF<sub>2</sub>, CHCl<sub>2</sub>, CH<sub>2</sub>F, CH<sub>2</sub>Cl, CH<sub>2</sub>OH, CN, C<sup>o</sup>CH, 1-propynyl, 2-propynyl, vinyl, OMe, OEt, SMe, or SEt.

- 5 Applicants' invention provides for modulation of gene expression in prokaryotic and eukaryotic host cells. Thus, the present invention also relates to a method for modulating gene expression in a host cell selected from the group consisting of a bacterial cell, a fungal cell, a yeast cell, a plant cell, an animal cell, and a mammalian cell. Preferably, the host cell is a yeast cell, a plant cell, a murine cell, or a human cell.
- 10 Expression in transgenic host cells may be useful for the expression of various polypeptides of interest including but not limited to therapeutic polypeptides, pathway intermediates; for the modulation of pathways already existing in the host for the synthesis of new products heretofore not possible using the host; cell based assays; and the like. Additionally the gene products may be useful for conferring higher growth yields of the host
- 15 or for enabling alternative growth mode to be utilized.

#### HOST CELLS AND NON-HUMAN ORGANISMS OF THE INVENTION

- As described above, the gene expression modulation system of the present invention may be used to modulate gene expression in a host cell. Expression in transgenic host cells
- 20 may be useful for the expression of various genes of interest. Thus, Applicants' invention also provides an isolated host cell comprising a gene expression system according to the invention. The present invention also provides an isolated host cell comprising a gene expression cassette according to the invention. Applicants' invention also provides an isolated host cell comprising a polynucleotide or polypeptide according to the invention. The isolated
- 25 host cell may be either a prokaryotic or a eukaryotic host cell.

- Preferably, the host cell is selected from the group consisting of a bacterial cell, a fungal cell, a yeast cell, a plant cell, an animal cell, and a mammalian cell. Examples of preferred host cells include, but are not limited to, fungal or yeast species such as *Aspergillus*, *Trichoderma*, *Saccharomyces*, *Pichia*, *Candida*, *Hansenula*, or bacterial species such as
- 30 those in the genera *Synechocystis*, *Synechococcus*, *Salmonella*, *Bacillus*, *Acinetobacter*, *Rhodococcus*, *Sireptomycetes*, *Escherichia*, *Pseudomonas*, *Methylobacter*, *Alcaligenes*, *Synechocystis*, *Anabaena*, *Thiobacillus*, *Methanobacterium* and *Klebsiella*, plant, animal, and mammalian host cells. More preferably, the host cell is a yeast cell, a plant

cell, a murine cell, or a human cell.

In a specific embodiment, the host cell is a yeast cell selected from the group consisting of a *Saccharomyces*, a *Pichia*, and a *Candida* host cell.

In another specific embodiment, the host cell is a plant cell selected from the group consisting of an apple, *Arabidopsis*, bajra, banana, barley, bean, beet, blackgram, chickpea, chili, cucumber, eggplant, favabean, maize, melon, millet, mungbean, oat, okra, *Panicum*, papaya, peanut, pea, pepper, pigeonpea, pineapple, *Phaseolus*, potato, pumpkin, rice, sorghum, soybean, squash, sugarcane, sugarbeet, sunflower, sweet potato, tea, tomato, tobacco, watermelon, and wheat host cell.

10 In another specific embodiment, the host cell is a murine cell.

In another specific embodiment, the host cell is a human cell.

Host cell transformation is well known in the art and may be achieved by a variety of methods including but not limited to electroporation, viral infection, plasmid/vector transfection, non-viral vector mediated transfection, *Agrobacterium*-mediated transformation, particle bombardment, and the like. Expression of desired gene products involves culturing the transformed host cells under suitable conditions and inducing expression of the transformed gene. Culture conditions and gene expression protocols in prokaryotic and eukaryotic cells are well known in the art (see General Methods section of Examples). Cells may be harvested and the gene products isolated according to protocols specific for the gene product.

20 In addition, a host cell may be chosen which modulates the expression of the inserted polynucleotide, or modifies and processes the polypeptide product in the specific fashion desired. Different host cells have characteristic and specific mechanisms for the translational and post-translational processing and modification (*e.g.*, glycosylation, cleavage [*e.g.*, of signal sequence]) of proteins. Appropriate cell lines or host systems can be chosen to ensure the desired modification and processing of the foreign protein expressed. For example, expression in a bacterial system can be used to produce a non-glycosylated core protein product.

However, a polypeptide expressed in bacteria may not be properly folded. Expression in yeast can produce a glycosylated product. Expression in eukaryotic cells can increase the likelihood of "native" glycosylation and folding of a heterologous protein. Moreover, expression in mammalian cells can provide a tool for reconstituting, or constituting, the polypeptide's activity. Furthermore, different vector/host expression systems may affect processing reactions, such as proteolytic cleavages, to a different extent.

30

Applicants' invention also relates to a non-human organism comprising an isolated host cell according to the invention. Preferably, the non-human organism is selected from the group consisting of a bacterium, a fungus, a yeast, a plant, an animal, and a mammal. More preferably, the non-human organism is a yeast, a plant, a mouse, a rat, a rabbit, a cat, a dog, a bovine, a goat, a pig, a horse, a sheep, a monkey, or a chimpanzee.

In a specific embodiment, the non-human organism is a yeast selected from the group consisting of *Saccharomyces*, *Pichia*, and *Candida*.

In another specific embodiment, the non-human organism is a plant selected from the group consisting of an apple, *Arabidopsis*, bajra, banana, barley, beans, beet, blackgram, chickpea, chili, cucumber, eggplant, favabean, maize, melon, millet, mungbean, oat, okra, *Panicum*, papaya, peanut, pea, pepper, pigeonpea, pineapple, *Phaseolus*, potato, pumpkin, rice, sorghum, soybean, squash, sugarcane, sugarbeet, sunflower, sweet potato, tea, tomato, tobacco, watermelon, and wheat.

In another specific embodiment, the non-human organism is a *Mus musculus* mouse.

#### MEASURING GENE EXPRESSION/TRANSCRIPTION

One useful measurement of Applicants' methods of the invention is that of the transcriptional state of the cell including the identities and abundances of RNA, preferably mRNA species. Such measurements are conveniently conducted by measuring cDNA abundances by any of several existing gene expression technologies.

Nucleic acid array technology is a useful technique for determining differential mRNA expression. Such technology includes, for example, oligonucleotide chips and DNA microarrays. These techniques rely on DNA fragments or oligonucleotides which correspond to different genes or cDNAs which are immobilized on a solid support and hybridized to probes prepared from total mRNA pools extracted from cells, tissues, or whole organisms and converted to cDNA. Oligonucleotide chips are arrays of oligonucleotides synthesized on a substrate using photolithographic techniques. Chips have been produced which can analyze for up to 1700 genes. DNA microarrays are arrays of DNA samples, typically PCR products, that are robotically printed onto a microscope slide. Each gene is analyzed by a full or partial-length target DNA sequence. Microarrays with up to 10,000 genes are now routinely prepared commercially. The primary difference between these two techniques is that oligonucleotide chips typically utilize 25-mer oligonucleotides which allow fractionation of short DNA molecules whereas the larger DNA targets of microarrays, approximately 1000 base pairs, may



provide more sensitivity in fractionating complex DNA mixtures.

Another useful measurement of Applicants' methods of the invention is that of determining the translation state of the cell by measuring the abundances of the constituent protein species present in the cell using processes well known in the art.

5       Where identification of genes associated with various physiological functions is desired, an assay may be employed in which changes in such functions as cell growth, apoptosis, senescence, differentiation, adhesion, binding to a specific molecules, binding to another cell, cellular organization, organogenesis, intracellular transport, transport facilitation, energy conversion, metabolism, myogenesis, neurogenesis, and/or hematopoiesis is measured.

10       In addition, selectable marker or reporter gene expression may be used to measure gene expression modulation using Applicants' invention.

Other methods to detect the products of gene expression are well known in the art and include Southern blots (DNA detection), dot or slot blots (DNA, RNA), Northern blots (RNA), and RT-PCR (RNA) analyses. Although less preferred, labeled proteins can be used to detect  
15 a particular nucleic acid sequence to which it hybridizes.

In some cases it is necessary to amplify the amount of a nucleic acid sequence. This may be carried out using one or more of a number of suitable methods including, for example, polymerase chain reaction ("PCR"), ligase chain reaction ("LCR"), strand displacement amplification ("SDA"), transcription-based amplification, and the like. PCR is carried out in  
20 accordance with known techniques in which, for example, a nucleic acid sample is treated in the presence of a heat stable DNA polymerase, under hybridizing conditions, with one oligonucleotide primer for each strand of the specific sequence to be detected. An extension product of each primer that is synthesized is complementary to each of the two nucleic acid strands, with the primers sufficiently complementary to each strand of the specific sequence to  
25 hybridize therewith. The extension product synthesized from each primer can also serve as a template for further synthesis of extension products using the same primers. Following a sufficient number of rounds of synthesis of extension products, the sample may be analyzed as described above to assess whether the sequence or sequences to be detected are present.

30       The present invention may be better understood by reference to the following non-limiting Examples, which are provided as exemplary of the invention.

## EXAMPLES

### GENERAL METHODS

- Standard recombinant DNA and molecular cloning techniques used herein are well known in the art and are described by Sambrook, J., Fritsch, E. F. and Maniatis, T. *Molecular Cloning: A Laboratory Manual*; Cold Spring Harbor Laboratory Press: Cold Spring Harbor, (1989) (Maniatis) and by T. J. Silhavy, M. L. Bennis, and L. W. Enquist, *Experiments with Gene Fusions*, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y. (1984) and by Ausubel, F. M. et al., *Current Protocols in Molecular Biology*, Greene Publishing Assoc. and Wiley-Interscience (1987).
- 10 Methods for plant tissue culture, transformation, plant molecular biology, and plant, general molecular biology may be found in *Plant Tissue Culture Concepts and Laboratory Exercises* edited by RN Trigiano and DJ Gray, 2<sup>nd</sup> edition, 2000, CRC press, New York; *Agrobacterium Protocols* edited by KMA Gartland and MR Davey, 1995, Humana Press, Totowa, New Jersey; *Methods in Plant Molecular Biology*, P. Maliga et al., 1995, Cold Spring Harbor Lab Press, New York; and *Molecular Cloning*, J. Sambrook et al., 1989, Cold Spring Harbor Lab Press, New York.

- Materials and methods suitable for the maintenance and growth of bacterial cultures are well known in the art. Techniques suitable for use in the following examples may be found as set out in *Manual of Methods for General Bacteriology* (Phillipp Gerhardt, R. G. E. Murray, Ralph N. Costilow, Eugene W. Nester, Willis A. Wood, Noel R. Krieg and G. Briggs Phillips, eds), American Society for Microbiology, Washington, DC. (1994)) or by Thomas D. Brock in *Biotechnology: A Textbook of Industrial Microbiology*, Second Edition, Sinauer Associates, Inc., Sunderland, MA (1989). All reagents, restriction enzymes and materials used for the growth and maintenance of host cells were obtained from Aldrich Chemicals (Milwaukee, WI), DIFCO Laboratories (Detroit, MI), GIBCO/BRL (Gaithersburg, MD), or Sigma Chemical Company (St. Louis, MO) unless otherwise specified.

- Manipulations of genetic sequences may be accomplished using the suite of programs available from the Genetics Computer Group Inc. (Wisconsin Package Version 9.0, Genetics Computer Group (GCG), Madison, WI). Where the GCG program "Pileup" is used the gap creation default value of 12, and the gap extension default value of 4 may be used. Where the GCG "Gap" or "Bestfit" programs are used the default gap creation penalty of 50 and the default gap extension penalty of 3 may be used. In any case where GCG program parameters are not prompted for, in these or any other GCG program, default values may be used.

The meaning of abbreviations is as follows: "h" means hour(s), "min" means minute(s), "sec" means second(s), "d" means day(s), "μl" means microliter(s), "ml" means milliliter(s), "L" means liter(s), "μM" means micromolar, "mM" means millimolar, "μg" means microgram(s), "mg" means milligram(s), "A" means adenine or adenosine, "T" means thymine or thymidine, "G" means guanine or guanosine, "C" means cytidine or cytosine, "x g" means times gravity, "nt" means nucleotide(s), "aa" means amino acid(s), "bp" means base pair(s), "kb" means kilobase(s), "k" means kilo, "μ" means micro, and "°C" means degrees Celsius.

10

**EXAMPLE 1**

Applicants' improved EcR-based inducible gene modulation system was developed for use in various applications including gene therapy, expression of proteins of interest in host cells, production of transgenic organisms, and cell-based assays. This Example describes the construction and evaluation of several gene expression cassettes for use in the EcR-based inducible gene expression system of the invention.

In various cellular backgrounds, including mammalian cells, insect ecdysone receptor (EcR) heterodimerizes with retinoid X receptor (RXR) and, upon binding of ligand, transactivates genes under the control of ecdysone response elements. Applicants constructed several EcR-based gene expression cassettes based on the spruce budworm *Choristoneura fumiferana* EcR ("CfEcR"; full length polynucleotide and amino acid sequences are set forth in SEQ ID NO: 49 and SEQ ID NO: 50, respectively), *C. fumiferana* ultraspiracle ("CfUSP"; full length polynucleotide and amino acid sequences are set forth in SEQ ID NO: 51 and SEQ ID NO: 52, respectively), and mouse *Mus musculus* RXRα (MmRXRα; full length polynucleotide and amino acid sequences are set forth in SEQ ID NO: 53 and SEQ ID NO: 54, respectively). The prepared receptor constructs comprise a ligand binding domain of EcR and of RXR or of USP; a DNA binding domain of GAL4 or of EcR; and an activation domain of VP16. The reporter constructs include a reporter gene, luciferase or LacZ, operably linked to a synthetic promoter construct that comprises either GAL4 or EcR/USP binding sites (response elements). Various combinations of these receptor and reporter constructs were cotransfected into CHO, NIH3T3, CV1 and 293 cells. Gene induction potential (magnitude of induction) and ligand specificity and sensitivity were examined using four different ligands: two steroidal ligands (ponasterone A and muristerone A) and two non-steroidal ligands (N-(2-ethyl-3-

methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine and N-(3,4-(1,2-ethylenedioxy)-2-methylbenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine) in a dose-dependent induction of reporter gene expression in the transfected cells. Reporter gene expression activities were assayed at 24hr or 48hr after ligand addition.

5

**Gene Expression Cassettes:** Ecdysone receptor-based, chemically inducible gene expression cassettes (switches) were constructed as followed, using standard cloning methods available in the art. The following is brief description of preparation and composition of each switch.

- 1.1 - GAL4EcR/VP16RXR: The D, E, and F domains from spruce budworm *Choristoneura fumiferana* EcR ("CfEcRDEF"; SEQ ID NO: 3) were fused to GAL4 DNA binding domain ("DNABD"; SEQ ID NO: 41) and placed under the control of an SV40e promoter (SEQ ID NO: 55). The DEF domains from mouse (*Mus musculus*) RXR ("MmRXRDEF"; SEQ ID NO: 22) were fused to the activation domain from VP16 ("VP16AD"; SEQ ID NO: 45) and placed under the control of an SV40e promoter (SEQ ID NO: 55). Five consensus GAL4 binding sites ("5XGAL4RE"; comprising 5, GAL4RE comprising SEQ ID NO: 47) were fused to a synthetic E1b minimal promoter (SEQ ID NO: 56) and placed upstream of the luciferase gene (SEQ ID NO: 57).

- 1.2 - GAL4EcR/VP16USP: This construct was prepared in the same way as in switch 1.1 above except MmRXRDEF was replaced with the D, E and F domains from spruce budworm USP ("CfUSPDEF"; SEQ ID NO: 58). The constructs used in this example are similar to those disclosed in U. S. Patent No. 5,880,333 except that *Choristoneura fumiferana* USP rather than *Drosophila melanogaster* USP was utilized.

- 1.3 - GAL4RXR/VP16CfEcR: MmRXRDEF (SEQ ID NO: 22) was fused to a GAL4DNABD (SEQ ID NO: 41) and CfEcRCDEF (SEQ ID NO: 1) was fused to a VP16AD (SEQ ID NO: 45).

- 1.4 - GAL4RXR/VP16DmEcR: This construct was prepared in the same way as switch 1.3 except CfEcRCDEF was replaced with DmEcRCDEF (SEQ ID NO: 6).

- 1.5 - GAL4USP/VP16CfEcR: This construct was prepared in the same way as switch 1.3 except MmRXRDEF was replaced with CfUSPDEF (SEQ ID NO: 58).

- 1.6 - GAL4RXRCfEcRV16: This construct was prepared so that both the GAL4 DNABD and the VP16AD were placed on the same molecule. GAL4DNABD (SEQ ID NO: 41) and VP16AD (SEQ ID NO: 45) were fused to CfEcRDEF (SEQ ID NO: 3) at N- and C-termini respectively. The fusion was placed under the control of an SV40e promoter (SEQ ID NO:

55).

1.7 - VP16CfEcR: This construct was prepared such that CfEcRCDEF (SEQ ID NO: 1) was fused to VP16AD (SEQ ID NO: 45) and placed under the control of an SV40e promoter (SEQ ID NO: 55). Six ecdysone response elements ("EcRE"; SEQ ID NO: 59) from the hsp27 gene were placed upstream of the promoter and a luciferase gene (SEQ ID NO: 57). This switch most probably uses endogenous RXR.

1.8 - DmVgRXR: This system was purchased from Invitrogen Corp., Carlsbad, California. It comprises a *Drosophila melanogaster* EcR ("DmEcR") with a modified DNABD fused to VP16AD and placed under the control of a CMV promoter (SEQ ID NO: 60). Full length MmRXR (SEQ ID NO: 53) was placed under the control of the RSV promoter (SEQ ID NO: 61). The reporter, pIND(SP1)LacZ, contains five copies of a modified ecdysone response element ("EcRE", E/GRE), three copies of an SP1 enhancer, and a minimal heat shock promoter, all of which were placed upstream to the LacZ reporter gene.

1.9 - CfVgRXR: This example was prepared in the same way as switch 1.8 except DmEcR was replaced with a truncated CfEcR comprising a partial A/B domain and the complete CDEF domains [SEQ ID NO: 62 (polynucleotide) and SEQ ID NO: 63 (polypeptide)].

1.10 - CfVgRXRdel: This example was prepared in the same way as switch 1.9 except MmRXR (SEQ ID NO: 53) was deleted.

**20 Cell lines:** Four cell lines: CHO, Chinese hamster *Cricetulus griseus* ovarian cell line; NIH3T3 (3T3) mouse *Mus musculus* cell line; 293 human *Homo sapiens* kidney cell line, and CV1 African green monkey kidney cell line were used in these experiments. Cells were maintained in their respective media and were subcultured when they reached 60% confluency. Standard methods for culture and maintenance of the cells were followed.

25

**Transfections:** Several commercially available lipofactors as well as electroporation methods were evaluated and the best conditions for transfection of each cell line were developed. CHO,

NIH3T3, 293 and CV1 cells were grown to 60% confluency. DNAs corresponding to the various switch constructs outlined in Examples 1.1 through 1.10 were transfected into CHO

30 cells, NIH3T3 cells, 293 cells, or CV1 cells as follows.

CHO cells: Cells were harvested when they reach 60-80% confluency and plated in 6- or 12- or 24- well plates at 250,000, 100,000, or 50,000 cells in 2.5, 1.0, or 0.5 ml of growth medium containing 10% Fetal bovine serum respectively. The next day, the cells were rinsed with

growth medium and transfected for four hours. LipofectAMINE™ 2000 (Life Technologies Inc.) was found to be the best transfection reagent for these cells. For 12- well plates, 4 µl of LipofectAMINE™ 2000 was mixed with 100 µl of growth medium. 1.0 µg of reporter construct and 0.25 µg of receptor construct(s) were added to the transfection mix. A second  
5 reporter construct was added [pTKRL (Promega), 0.1 µg/transfection mix] and comprised a *Renilla* luciferase gene (SEQ ID NO: 64) operably linked and placed under the control of a thymidine kinase (TK) constitutive promoter and was used for normalization. The contents of the transfection mix were mixed in a vortex mixer and let stand at room temperature for 30 min. At the end of incubation, the transfection mix was added to the cells maintained in 400 µl  
10 growth medium. The cells were maintained at 37°C and 5% CO<sub>2</sub> for four hours. At the end of incubation, 500 µl of growth medium containing 20% FBS and either DMSO (control) or a DMSO solution of appropriate ligands were added and the cells were maintained at 37 °C and 5% CO<sub>2</sub> for 24-48 hr. The cells were harvested and reporter activity was assayed. The same procedure was followed for 6 and 24 well plates as well except all the reagents were doubled  
15 for 6 well plates and reduced to half for 24-well plates.

NIH3T3 Cells: Superfect™ (Qiagen Inc.) was found to be the best transfection reagent for 3T3 cells. The same procedures described for CHO cells were followed for 3T3 cells as well with two modifications. The cells were plated when they reached 50% confluency. 125,000 or 50,000 or 25,000 cells were plated per well of 6- or 12- or 24-well plates respectively. The  
20 GA14EcR/VP16RXR and reporter vector DNAs were transfected into NIH3T3 cells, the transfected cells were grown in medium containing PonA, MurA, N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-t-butylhydrazine, or N-(3,4-(1,2-ethylenedioxy)-2-methylbenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine for 48 hr. The ligand treatments were performed as described in the CHO cell section above.

25 293 Cells: LipofectAMINE™ 2000 (Life Technologies) was found to be the best lipofactor for 293 cells. The same procedures described for CHO were followed for 293 cells except that the cells were plated in biocoated plates to avoid clumping. The ligand treatments were performed  
as described in the CHO cell section above.

CV1 Cells: LipofectAMINE™ plus (Life Technologies) was found to be the best lipofactor  
30 for CV1 cells. The same procedures described for NIH3T3 cells were followed for CV1 cells

**Ligands:** Ponasterone A and Muristerone A were purchased from Sigma Chemical Company. The two non-steroids N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-t-

butylhydrazine, or N-(3,4-(1,2-ethylenedioxy)-2-methylbenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine are synthetic stable ecdysteroids synthesized at Rohm and Haas Company. All ligands were dissolved in DMSO and the final concentration of DMSO was maintained at 0.1% in both controls and treatments.

5

- Reporter Assays:** Cells were harvested 24-48 hr after adding ligands. 125, 250, or 500 µl of passive lysis buffer (part of Dual-luciferase<sup>TM</sup> reporter assay system from Promega Corporation) were added to each well of 24- or 12- or 24-well plates respectively. The plates were placed on a rotary shaker for 15 min. Twenty µl of lysate was assayed. Luciferase activity was measured using Dual-luciferase<sup>TM</sup> reporter assay system from Promega Corporation following the manufacturer's instructions. β-Galactosidase was measured using Galacto-Star<sup>TM</sup> assay kit from TROPIX following the manufacturer's instructions. All luciferase and β-galactosidase activities were normalized using *Renilla* luciferase as a standard. Fold activities were calculated by dividing normalized relative light units ("RLU") in ligand treated cells with normalized RLU in DMSO treated cells (untreated control).

10

The results of these experiments are provided in the following tables.

Table 1

Transactivation of reporter genes through various switches in CHO cells

20

Composition of Switch	Mean Fold Activation with 50µM N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-t-butylhydrazine
1.1 GAL4EcR + VP16RXR pGAL4RELuc	267
1.2 GAL4EcR + VP16USP pGAL4RELuc	2
1.3 GAL4RXR + VP16CfEcR pGAL4RELuc	85
1.4 GAL4RXR + VP16DmEcR pGAL4RELuc	312
1.5 GAL4USP + VP16CfEcR pGAL4RELuc	2
1.6 GAL4CfEcRVP16 pGAL4RELuc	9
1.7 VP16CfEcR pEcRELuc	36
1.8 DmVgRXR + MmRXR pIND(SP1)LacZ	14

1.9 CfVgRXR + MmRXR pIND(SP1)LacZ	27
1.10 CfVgRXR pIND(SP1)LacZ	29

Table 2

Transactivation of reporter genes through various switches in 3T3 cells

Composition of Switch	Mean Fold Activation Through N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-t-butylhydrazine
1.1 GAL4EcR + VP16RXR pGAL4RELuc	1118
1.2 GAL4EcR + VP16USP pGAL4RELuc	2
1.3 GAL4RXR + VP16CfEcR pGAL4RELuc	47
1.4 GAL4RXR + VP16DmEcR pGAL4RELuc	269
1.5 GAL4USP + VP16CfEcR pGAL4RELuc	3
1.6 GAL4CfEcRVP16 pGAL4RELuc	7
1.7 VP16CfEcR pEcRELuc	1
1.8 DmVgRXR + MmRXR pIND(SP1)LacZ	21
1.9 CfVgRXR + MmRXR pIND(SP1)LacZ	19
1.10 CfVgRXR pIND(SP1)LacZ	2

5

Table 3

Transactivation of reporter genes through various switches in 293 cells

Composition of Switch	Mean Fold Activation Through N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-t-butylhydrazine
1.1 GAL4EcR + VP16RXR pGAL4RELuc	125
1.2 GAL4EcR + VP16USP pGAL4RELuc	2
1.3 GAL4RXR + VP16CfEcR pGAL4RELuc	17
1.4 GAL4RXR + VP16DmEcR pGAL4RELuc	3
1.5 GAL4USP + VP16CfEcR pGAL4RELuc	2
1.6 GAL4CfEcRVP16 pGAL4RELuc	3



1.7 VP16CfEcR pEcRELuc	2
1.8 DmVgRXR + MmRXR pIND(SP1)LacZ	21
1.9 CfVgRXR + MmRXR pIND(SP1)LacZ	12
1.10 CfVgRXR pIND(SP1)LacZ	3

**Table 4**  
Transactivation of reporter genes through various switches in CV1 cells

Composition of Switch	Mean Fold Activation Through N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-t-butylhydrazine
1.1 GAL4EcR + VP16RXR pGAL4RELuc	279
1.2 GAL4EcR + VP16USP pGAL4RELuc	2
1.3 GAL4RXR + VP16CfEcR pGAL4RELuc	25
1.4 GAL4RXR + VP16DmEcR pGAL4RELuc	80
1.5 GAL4USP + VP16CfEcR pGAL4RELuc	3
1.6 GAL4CfEcRVP16 pGAL4RELuc	6
1.7 VP16CfEcR pEcRELuc	1
1.8 DmVgRXR + MmRXR pIND(SP1)LacZ	12
1.9 CfVgRXR + MmRXR pIND(SP1)LacZ	7
1.10 CfVgRXR pIND(SP1)LacZ	1

5

**Table 5**  
Transactivation of reporter gene GAL4CfEcRDEF/VP16MmRXRDEF (switch 1.1)  
through steroids and non-steroids in 3T3 cells.

Ligand	Mean Fold Induction at 1.0 $\mu$ M Concentration
1. Ponasterone A	1.0
2. Muriesterone A	1.0
3. N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine	116
4. N'-(3,4-(1,2-ethylenedioxy)-2-methylbenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine	601

10

**Table 6**  
**Transactivation of reporter gene GAL4MmRXRDEF/VP16CfEcRCDEF (switch 1.3)**  
**through steroids and non-steroids in 3T3 cells.**

Ligand	Mean Fold Induction at 1.0 $\mu$ M Concentration
1. Ponasterone A	1.0
2. Muristerone A	1.0
3. N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine	71
4. N'-(3,4-(1,2-ethylenedioxy)-2-methylbenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine	54

5

Applicants' results demonstrate that the non-steroidal ecdysone agonists, N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine and N'-(3,4-(1,2-ethylenedioxy)-2-methylbenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine, were more potent activators of CfEcR as compared to *Drosophila melanogaster* EcR (DmEcR). (see  
 10 Tables 1-4). Also, in the mammalian cell lines tested, MmRXR performed better than CfUSP as a heterodimeric partner for CfEcR. (see Tables 1-4). Additionally, Applicants' inducible gene expression modulation system performed better when exogenous MmRXR was used than when the system relied only on endogenous RXR levels (see Tables 1-4).

Applicants' results also show that in a CfEcR-based inducible gene expression system,  
 15 the non-steroidal ecdysone agonists induced reporter gene expression at a lower concentration (i.e., increased ligand sensitivity) as compared to the steroid ligands, ponasterone A and muristerone A (see Tables 5 and 6).

Out of 10 EcR based gene switches tested, the GAL4EcR/VP16RXR switch (Switch 1.1) performed better than any other switch in all four cell lines examined and was more  
 20 sensitive to non-steroids than steroids. The results also demonstrate that placing the activation domain (AD) and DNA binding domain (DNABD) on each of the two partners reduced background when compared to placing both AD and DNABD together on one of the two partners. Therefore, a switch format where the AD and DNABD are separated between two partners, works well for EcR-based gene switch applications.

25 In addition, the MmRXR/EcR-based switches performed better than CfUSP/EcR-based switches, which have a higher background activity than the MmRXR/EcR switches in the absence of ligand.

Finally, the GAL4EcR/VP16RXR switch (Switch 1.1) was more sensitive to non-steroid ligands than to the steroid ligands (see Tables 5 and 6). In particular, steroid ligands

initiated transactivation at concentrations of 50  $\mu$ M, whereas the non-steroid ligands initiated transactivation at less than 1  $\mu$ M (submicromolar) concentration.

## EXAMPLE 2

5

This Example describes Applicants' further analysis of truncated EcR and RXR polypeptides in the improved EcR-based inducible gene expression system of the invention. To identify the best combination and length of two receptors that give a switch with a) maximum induction in the presence of ligand; b) minimum background in the absence of ligand; c) highly sensitive to ligand concentration; and d) minimum cross-talk among ligands and receptors, Applicants made and analyzed several truncation mutations of the CfEcR and MmRXR receptor polypeptides in NIH3T3 cells.

Briefly, polynucleotides encoding EcR or RXR receptors were truncated at the junctions of A/B, C, D, E and F domains and fused to either a GAL4 DNA binding domain encoding polynucleotide (SEQ ID NO: 41) for CfEcR, or a VP16 activation domain encoding polynucleotide (SEQ ID NO: 45) for MmRXR as described in Example 1. The resulting receptor truncation/fusion polypeptides were assayed in NIH3T3 cells. Plasmid pFRLUC (Stratagene) encoding a luciferase polypeptide was used as a reporter gene construct and pTKRL (Promega) encoding a *Renilla* luciferase polypeptide under the control of the constitutive TK promoter was used to normalize the transfections as described above. The analysis was performed in triplicates and mean luciferase counts were determined as described above.

### Gene Expression Cassettes Encoding Truncated Ecdysone Receptor Polypeptides

Gene expression cassettes comprising polynucleotides encoding either full length or truncated CfEcR polypeptides fused to a GAL4 DNA binding domain (SEQ ID NO: 41): GAL4CfEcRA/BCDEF (full length CfEcRA/BCDEF; SEQ ID NO: 49), GAL4CfEcRCDEF (CfEcRCDEF; SEQ ID NO: 1), GAL4CfEcR1/2CDEF (CfEcR1/2CDEF; SEQ ID NO: 2), GAL4CfEcRDEF (CfEcRDEF; SEQ ID NO: 3), GAL4CfEcREF (CfEcREF; SEQ ID NO: 4), and GAL4CfEcRDE (CfEcRDE; SEQ ID NO: 5) were transfected into NIH3T3 cells along with VP16MmRXRDEF (constructed as in Example 1.1; Figure 11) or VP16MmRXREF [constructed as in Example 1.1 except that MmRXRDEF was replaced with MmRXREF (SEQ ID NO: 23); Figure 12], and pFRLUC and pTKRL plasmid DNAs. The transfected cells were grown in the presence 0, 1, 5 or 25  $\mu$ M of N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-

dimethylbenzoyl)-N'-tert-butylhydrazine or PonA for 48 hr. The cells were harvested, lysed and luciferase reporter activity was measured in the cell lysates. Total fly luciferase relative light units are presented. The number on the top of each bar is the maximum fold induction for that treatment.

- 5 Applicants' results show that the EF domain of MmRXR is sufficient and performs better than DEF domains of this receptor (see Figures 11 and 12). Applicants have also shown that, in general, EcR/RXR receptor combinations are insensitive to PonA (see Figures 11 and 12). As shown in the Figures 11 and 12, the GAL4CfEcRCDEF hybrid polypeptide (SEQ ID NO: 7) performed better than any other CfEcR hybrid polypeptide.

10 Gene Expression Cassettes Encoding Truncated Retinoid X Receptor Polypeptides

- Gene expression cassettes comprising polynucleotides encoding either full length or truncated MmRXR polypeptides fused to a VP16 transactivation domain (SEQ ID NO: 45): VP16MmRXRA/BCDEF (full length MmRXRA/BCDEF; SEQ ID NO: 53), VP16MmRXRCDEF (MmRXRCDEF; SEQ ID NO: 21), VP16MmRXRDEF (MmRXRDEF; SEQ ID NO: 22), VP16MmRXREF (MmRXREF; SEQ ID NO: 23), VP16MmRXRBam-EF ("MmRXRBam-EF" or "MmRXR-truncatedEF"; SEQ ID NO: 24), and VP16MmRXRAF2del ("MmRXRAF2del" or "MmRXR-E"; SEQ ID NO: 25) constructs were transfected into NIH3T3 cells along with GAL4CfEcRCDEF (constructed as in Example 1.1; Figure 13) or GAL4CfEcRDEF [constructed as in Example 1.1 except CfEcRCDEF was replaced with CfEcRDEF (SEQ ID NO: 3); Figure 14], pFRLuc and pTKRL plasmid DNAs as described above. The transfected cells were grown in the presence 0, 1, 5 and 25 uM of N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine or PonA for 48 hr. The cells were harvested and lysed and reporter activity was measured in the cell lysate. Total fly luciferase relative light units are presented. The number on top of each bar is the maximum fold induction in that treatment.

- Of all the truncations of MmRXR tested, Applicants' results show that the MmRXREF receptor was the best partner for CfEcR (Figures 13 and 14). CfEcRCDEF showed better induction than CfEcRDEF using MmRXREF. Deleting AF2 (abbreviated "EF-AF2del") or helices 1-3 of the E domain (abbreviated "EF-Bamdel") resulted in an RXR receptor that reduced gene induction and ligand sensitivity when partnered with either CfEcRCDEF (Figure 13) or CfEcRDEF (Figure 14) in NIH3T3 cells. In general, the CfEcR/RXR-based switch was much more sensitive to the non-steroid N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine than to the steroid PonA.

**EXAMPLE 3**

This Example describes Applicants' further analysis of gene expression cassettes encoding truncated EcR or RXR receptor polypeptides that affect either ligand binding activity or ligand sensitivity, or both. Briefly, six different combinations of chimeric receptor pairs, constructed as described in Examples 1 and 2, were further analyzed in a single experiment in NIH3T3 cells. These six receptor pair combinations and their corresponding sample numbers are depicted in Table 7.

10

**Table 7****CfEcR + MmRXR Truncation Receptor Combinations in NIH3T3 Cells**

<b>Figure 15 X-Axis Sample No.</b>	<b>EcR Polypeptide Construct</b>	<b>RXR Polypeptide Construct</b>
Samples 1 and 2	GAL4CfEcRCDEF	VP16RXRA/BCDEF (Full length)
Samples 3 and 4	GAL4CfEcRCDEF	VP16RXRDEF
Samples 5 and 6	GAL4CfEcRCDEF	VP16RXREF
Samples 7 and 8	GAL4CfEcRDEF	VP16RXRA/BCDEF (Full length)
Samples 9 and 10	GAL4CfEcRDEF	VP16RXRDEF
Samples 11 and 12	GAL4CfEcRDEF	VP16RXREF

The above receptor construct pairs, along with the reporter plasmid pFRLuc were transfected into NIH3T3 cells as described above. The six CfEcR truncation receptor combinations were duplicated into two groups and treated with either steroid (odd numbers on x-axis of Figure 15) or non-steroid (even numbers on x-axis of Figure 15). In particular, the cells were grown in media containing 0, 1, 5 or 25 uM PonA (steroid) or N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine (non-steroid) ligand. The reporter gene activity was measured and total RLU are shown. The number on top of each bar is the maximum fold induction for that treatment and is the mean of three replicates.

As shown in Figure 15, the CfEcRCDEF/MmRXREF receptor combinations were the best switch pairs both in terms of total RLU and fold induction (compare columns 1-6 to columns 7-12). This confirms Applicants' earlier findings as described in Example 2 (Figures 11-14). The same gene expression cassettes encoding the truncated EcR and RXR polypeptides were also assayed in a human lung carcinoma cell line A549 (ATCC) and similar results were observed (data not shown).

**WE CLAIM:**

1. A gene expression modulation system comprising:
  - a) a first gene expression cassette that is capable of being expressed in a host cell comprising a polynucleotide sequence that encodes a first polypeptide comprising:
    - 5 i) a DNA-binding domain that recognizes a response element associated with a gene whose expression is to be modulated;
    - i) a ligand binding domain comprising a ligand binding domain from a nuclear receptor;
  - b) a second gene expression cassette that is capable of being expressed in
    - 10 the host cell comprising a polynucleotide sequence that encodes a second polypeptide comprising:
      - i) a transactivation domain; and
      - ii) a ligand binding domain comprising a ligand binding domain from a nuclear receptor other than ultraspiracle (USP);
- 1.5 wherein the transactivation domain is from a nuclear receptor other than an ecdysone receptor, a retinoid X receptor, or an ultraspiracle receptor; and wherein the ligand binding domains from the first polypeptide and the second polypeptide are different and dimerize.
2. The gene expression modulation system according to claim 1, further comprising a third gene expression cassette comprising:
  - 20 i) a response element to which the DNA-binding domain of the first polypeptide binds;
  - ii) a promoter that is activated by the transactivation domain of the second polypeptide; and
  - iii) the gene whose expression is to be modulated.
- 2.5 3. The gene expression modulation system according to claim 1, wherein the ligand binding domain of the first polypeptide is an ecdysone receptor polypeptide.
4. The gene expression modulation system according to claim 1, wherein the ligand binding domain of the second polypeptide is a retinoid X receptor polypeptide.
5. A gene expression modulation system comprising:
  - 30 a) a first gene expression cassette that is capable of being expressed in a host cell comprising a polynucleotide sequence that encodes a first polypeptide comprising:
    - i) a DNA-binding domain that recognizes a response element associated with a gene whose expression is to be modulated; and

- ii) a ligand binding domain comprising a ligand binding domain from an ecdysone receptor; and
- b) a second gene expression cassette that is capable of being expressed in the host cell comprising a polynucleotide sequence that encodes a second polypeptide comprising:
- 5 i) a transactivation domain; and
- ii) a ligand binding domain comprising a ligand binding domain from a retinoid X receptor;

wherein the ligand binding domains from the first polypeptide and the second polypeptide are different and dimerize.

10

6. The gene expression modulation system according to claim 5, further comprising a third gene expression cassette comprising:

- i) a response element to which the DNA-binding domain of the first polypeptide binds;
- 15 ii) a promoter that is activated by the transactivation domain of the second polypeptide; and
- iii) the gene whose expression is to be modulated.

7. The gene expression modulation system according to claim 5, wherein the ligand binding domain of the first polypeptide is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.

20

8. The gene expression modulation system according to claim 5, wherein the ligand binding domain of the first polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, and SEQ ID NO: 20.

25

9. The gene expression modulation system according to claim 5, wherein the ligand binding domain of the second polypeptide is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30.

30

10. The gene expression modulation system according to claim 5, wherein the

ligand binding domain of the second polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 38, SEQ ID NO: 39, and SEQ ID NO: 40.

5 11. A gene expression modulation system comprising:

a) a first gene expression cassette that is capable of being expressed in a host cell comprising a polynucleotide sequence that encodes a first polypeptide comprising:

- 10 i) a DNA-binding domain that recognizes a response element associated with a gene whose expression is to be modulated; and  
ii) a ligand binding domain comprising a ligand binding domain from a retinoid X receptor; and

b) a second gene expression cassette that is capable of being expressed in the host cell comprising a polynucleotide sequence that encodes a second polypeptide comprising:

- 15 i) a transactivation domain; and  
ii) a ligand binding domain comprising a ligand binding domain from an ecdysone receptor;

wherein the ligand binding domains from the first polypeptide and the second polypeptide are different and dimerize.

20 12. The gene expression modulation system according to claim 11, further comprising a third gene expression cassette comprising:

- i) a response element to which the DNA-binding domain of the first polypeptide binds;  
ii) a promoter that is activated by the transactivation domain of the second polypeptide; and  
25 iii) the gene whose expression is to be modulated.

13. The gene expression modulation system according to claim 11, wherein the ligand binding domain of the first polypeptide is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30.

14. The gene expression modulation system according to claim 11, wherein the ligand binding domain of the first polypeptide comprises an amino acid sequence selected from



the group consisting of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 38, SEQ ID NO: 39, and SEQ ID NO: 40.

15. The gene expression modulation system according to claim 11, wherein the  
5 ligand binding domain of the second polypeptide is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.

16. The gene expression modulation system according to claim 11, wherein the  
10 ligand binding domain of the second polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, and SEQ ID NO: 20.

17. A gene expression cassette comprising a polynucleotide encoding a hybrid  
15 polypeptide comprising a DNA-binding domain and an ecdysone receptor ligand binding domain, wherein the DNA binding domain is from a nuclear receptor other than an ecdysone receptor.

18. The gene expression cassette according to claim 18, wherein the DNA-binding domain is a GAL4 DNA-binding domain or a LexA DNA-binding domain.

- 20 19. A gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide comprising a DNA-binding domain and a retinoid X receptor ligand binding domain, wherein the DNA binding domain is from a nuclear receptor other than a retinoid X receptor.

20. The gene expression cassette according to claim 19, wherein the DNA-binding  
25 domain is a GAL4 DNA-binding domain or a LexA DNA-binding domain.

21. A gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide comprising a transactivation domain and an ecdysone receptor ligand binding domain, wherein the transactivation domain is from a nuclear receptor other than an ecdysone receptor.

- 30 22. The gene expression cassette according to claim 21, wherein the transactivation domain is a VP16 transactivation domain.

23. A gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide comprising a transactivation domain and a retinoid X receptor ligand binding

domain, wherein the transactivation domain is from a nuclear receptor other than a retinoid X receptor.

24. The gene expression cassette according to claim 22, wherein the transactivation domain is a VP16 transactivation domain.

5 25. A gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide comprising a DNA-binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of a GAL4 DBD (SEQ ID NO: 41) or a LexA DBD (SEQ ID NO: 43) and an ecdysone receptor ligand binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ  
10 ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.

26. A gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide comprising a DNA-binding domain comprising an amino acid sequence selected from the group consisting of a GAL4 DBD (SEQ ID NO: 42) or a LexA DBD (SEQ ID NO:  
15 44) and an ecdysone receptor ligand binding domain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, and SEQ ID NO: 20.

27. A gene expression cassette comprising a polynucleotide encoding a hybrid  
20 polypeptide comprising a DNA-binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of a GAL4 DBD (SEQ ID NO: 41) or a LexA DBD (SEQ ID NO: 43) and a retinoid X receptor ligand binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID  
25 NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30.

28. A gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide comprising a DNA-binding domain comprising an amino acid sequence selected from the group consisting of a GAL4 DBD (SEQ ID NO: 42) or a LexA DBD (SEQ ID NO:  
44) and a retinoid X receptor ligand binding domain comprising an amino acid sequence  
30 selected from the group consisting of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 38, SEQ ID NO: 39, and SEQ ID NO: 40.

29. A gene expression cassette comprising a polynucleotide encoding a hybrid

polypeptide comprising a transactivation domain encoded by a polynucleotide comprising a nucleic acid sequence of SEQ ID NO: 45 and an ecdysone receptor ligand binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, 5 SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.

30. A gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide comprising a transactivation domain comprising an amino acid sequence of SEQ ID NO: 46 and an ecdysone receptor ligand binding domain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 10 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, and SEQ ID NO: 20.

31. A gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide comprising a transactivation domain encoded by a polynucleotide comprising a nucleic acid sequence of SEQ ID NO: 45 and a retinoid X receptor ligand binding domain 15 encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30.

32. A gene expression cassette comprising a polynucleotide encoding a hybrid 20 polypeptide comprising a transactivation domain comprising an amino acid sequence of SEQ ID NO: 46 and a retinoid X receptor ligand binding domain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 38, SEQ ID NO: 39, and SEQ ID NO: 40.

25 33. An isolated polynucleotide encoding an ecdysone receptor polypeptide or a retinoid X receptor polypeptide comprising a truncation mutation, wherein the truncation mutation reduces ligand binding activity of the ecdysone receptor polypeptide or the retinoid X receptor polypeptide.

34. An isolated polynucleotide encoding an ecdysone receptor polypeptide or a 30 retinoid X receptor polypeptide comprising a truncation mutation, wherein the truncation mutation reduces steroid binding activity of the ecdysone receptor polypeptide or the retinoid X receptor polypeptide.

35. An isolated polynucleotide encoding an ecdysone receptor polypeptide or a

retinoid X receptor polypeptide comprising a truncation mutation, wherein the truncation mutation reduces non-steroid binding activity of the ecdysone receptor polypeptide or the retinoid X receptor polypeptide.

36. An isolated polynucleotide encoding an ecdysone receptor polypeptide or a  
5 retinoid X receptor polypeptide comprising a truncation mutation, wherein the truncation mutation enhances ligand binding activity of the ecdysone receptor polypeptide or the retinoid X receptor polypeptide.

37. An isolated polynucleotide encoding an ecdysone receptor polypeptide or a  
retinoid X receptor polypeptide comprising a truncation mutation, wherein the truncation  
10 mutation enhances steroid binding activity of the ecdysone receptor polypeptide or the retinoid X receptor polypeptide.

38. An isolated polynucleotide encoding an ecdysone receptor polypeptide or a  
retinoid X receptor polypeptide comprising a truncation mutation, wherein the truncation  
mutation enhances non-steroid binding activity of the ecdysone receptor polypeptide or the  
15 retinoid X receptor polypeptide.

39. An isolated polynucleotide encoding a retinoid X receptor polypeptide  
comprising a truncation mutation, wherein the truncation mutation increases ligand sensitivity  
of the retinoid X receptor polypeptide.

40. An isolated polynucleotide encoding a retinoid X receptor polypeptide  
20 comprising a truncation mutation, wherein the truncation mutation increases ligand sensitivity  
of a heterodimer, wherein the heterodimer comprises said retinoid X receptor polypeptide and a  
dimerization partner.

41. The isolated polynucleotide according to claim 40, wherein the dimerization  
partner is an ecdysone receptor polypeptide.

25 42. An isolated polynucleotide encoding a truncated ecdysone receptor  
polypeptide, wherein the polynucleotide comprises a nucleic acid sequence selected from the  
group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID  
NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.

43. An isolated polypeptide encoded by the isolated polynucleotide according to  
30 claim 42.

44. An isolated truncated ecdysone receptor polypeptide comprising an amino acid  
sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO:  
13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18,

SEQ ID NO: 19, and SEQ ID NO: 20.

45. An isolated polynucleotide encoding a truncated retinoid X receptor polypeptide, wherein the polynucleotide comprises a nucleic acid sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30.

46. An isolated polypeptide encoded by the isolated polynucleotide according to claim 45.

47. An isolated truncated retinoid X receptor polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 38, SEQ ID NO: 39, and SEQ ID NO: 40.

48. A method of modulating the expression of a gene in a host cell comprising the gene to be modulated comprising the steps of:

a) introducing into the host cell the gene expression modulation system according to claim 1; and

b) introducing into the host cell a ligand that independently combines with the ligand binding domains of the first polypeptide and the second polypeptide;

wherein the gene to be expressed is a component of a chimeric gene comprising:


i) a response element to which the DNA binding domain from the first polypeptide binds;

ii) a promoter that is activated by the transactivation domain of the second polypeptide; and

iii) a gene whose expression is to be modulated,

whereby a complex is formed comprising the ligand, the first polypeptide, and the second polypeptide, and whereby the complex modulates expression of the gene in the host cell.

49. The method according to claim 48, wherein the ligand is a compound of the formula:



wherein:

E is a (C<sub>4</sub>-C<sub>6</sub>)alkyl containing a tertiary carbon or a cyano(C<sub>3</sub>-C<sub>5</sub>)alkyl containing a tertiary carbon;

5 R<sup>1</sup> is H, Me, Et, i-Pr, F, formyl, CF<sub>3</sub>, CHF<sub>2</sub>, CHCl<sub>2</sub>, CH<sub>2</sub>F, CH<sub>2</sub>Cl, CH<sub>2</sub>OH, CH<sub>2</sub>OMe, CH<sub>2</sub>CN, CN, C<sup>o</sup>CH, 1-propynyl, 2-propynyl, vinyl, OH, OMe, OEt, cyclopropyl, CF<sub>2</sub>CF<sub>3</sub>, CH=CHCN, allyl, azido, SCN, or SCHF<sub>2</sub>;

R<sup>2</sup> is H, Me, Et, n-Pr, i-Pr, formyl, CF<sub>3</sub>, CHF<sub>2</sub>, CHCl<sub>2</sub>, CH<sub>2</sub>F, CH<sub>2</sub>Cl, CH<sub>2</sub>OH, CH<sub>2</sub>OMe, CH<sub>2</sub>CN, CN, C<sup>o</sup>CH, 1-propynyl, 2-propynyl, vinyl, Ac, F, Cl, OH, OMe, OEt, O-n-Pr, OAc, NMe<sub>2</sub>, NEt<sub>2</sub>, SMe, SEt, SOCF<sub>3</sub>, OCF<sub>2</sub>CF<sub>2</sub>H, COEt, cyclopropyl, CF<sub>2</sub>CF<sub>3</sub>, CH=CHCN, allyl, azido, OCF<sub>3</sub>, OCHF<sub>2</sub>, O-i-Pr, SCN, SCHF<sub>2</sub>, SMe, NH-CN, or  
10 joined with R<sup>3</sup> and the phenyl carbons to which R<sup>2</sup> and R<sup>3</sup> are attached to form an ethylenedioxy, a dihydrofuryl ring with the oxygen adjacent to a phenyl carbon, or a dihydropyryl ring with the oxygen adjacent to a phenyl carbon;

15 R<sup>3</sup> is H, Et, or joined with R<sup>2</sup> and the phenyl carbons to which R<sup>2</sup> and R<sup>3</sup> are attached to form an ethylenedioxy, a dihydrofuryl ring with the oxygen adjacent to a phenyl carbon, or a dihydropyryl ring with the oxygen adjacent to a phenyl carbon;

R<sup>4</sup>, R<sup>5</sup>, and R<sup>6</sup> are independently H, Me, Et, F, Cl, Br, formyl, CF<sub>3</sub>, CHF<sub>2</sub>, CHCl<sub>2</sub>, CH<sub>2</sub>F, CH<sub>2</sub>Cl, CH<sub>2</sub>OH, CN, C<sup>o</sup>CH, 1-propynyl, 2-propynyl, vinyl, OMe, OEt, SMe, or SEt.

20 50. A method of modulating the expression of a gene in a host cell comprising the gene to be modulated comprising the steps of:

a) introducing into the host cell the gene expression modulation system of claim 5; and

25 b) introducing into the host cell a ligand that independently combines with the ligand binding domains of the first polypeptide and the second polypeptide;

wherein the gene to be expressed is a component of a chimeric gene comprising:


i) a response element to which the DNA binding domain from the first polypeptide binds;

ii) a promoter that is activated by the transactivation domain of the second polypeptide; and

iii) a gene whose expression is to be modulated,

whereby a complex is formed comprising the ligand, the first polypeptide, and the second polypeptide, and whereby the complex modulates expression of the gene in the host cell.

51. The method according to claim 50, wherein the ligand is a compound of the formula:



wherein:

10 E is a (C<sub>4</sub>-C<sub>6</sub>)alkyl containing a tertiary carbon or a cyano(C<sub>3</sub>-C<sub>5</sub>)alkyl containing a tertiary carbon;

R<sup>1</sup> is H, Me, Et, i-Pr, F, formyl, CF<sub>3</sub>, CHF<sub>2</sub>, CHCl<sub>2</sub>, CH<sub>2</sub>F, CH<sub>2</sub>Cl, CH<sub>2</sub>OH, CH<sub>2</sub>OMe, CH<sub>2</sub>CN, CN, C<sup>o</sup>CH, 1-propynyl, 2-propynyl, vinyl, OH, OMe, OEt, cyclopropyl, CF<sub>2</sub>CF<sub>3</sub>, CH=CHCN, allyl, azido, SCN, or SCHF<sub>2</sub>;

15 R<sup>2</sup> is H, Me, Et, n-Pr, i-Pr, formyl, CF<sub>3</sub>, CHF<sub>2</sub>, CHCl<sub>2</sub>, CH<sub>2</sub>F, CH<sub>2</sub>Cl, CH<sub>2</sub>OH, CH<sub>2</sub>OMe, CH<sub>2</sub>CN, CN, C<sup>o</sup>CH, 1-propynyl, 2-propynyl, vinyl, Ac, F, Cl, OH, OMe, OEt, O-n-Pr, OAc, NMe<sub>2</sub>, NEt<sub>2</sub>, SMe, SEt, SOCF<sub>3</sub>, OCF<sub>2</sub>CF<sub>2</sub>H, COEt, cyclopropyl, CF<sub>2</sub>CF<sub>3</sub>, CH=CHCN, allyl, azido, OCF<sub>3</sub>, OCHF<sub>2</sub>, O-i-Pr, SCN, SCHF<sub>2</sub>, SMe, NH-CN, or joined with R<sup>3</sup> and the phenyl carbons to which R<sup>2</sup> and R<sup>3</sup> are attached to form an ethylenedioxy, a dihydrofuryl ring with the oxygen adjacent to a phenyl carbon, or a dihydropyryl ring with the oxygen adjacent to a phenyl carbon;


20 R<sup>3</sup> is H, Et, or joined with R<sup>2</sup> and the phenyl carbons to which R<sup>2</sup> and R<sup>3</sup> are attached to form an ethylenedioxy, a dihydrofuryl ring with the oxygen adjacent to a phenyl carbon, or a dihydropyryl ring with the oxygen adjacent to a phenyl carbon;

25 R<sup>4</sup>, R<sup>5</sup>, and R<sup>6</sup> are independently H, Me, Et, F, Cl, Br, formyl, CF<sub>3</sub>, CHF<sub>2</sub>, CHCl<sub>2</sub>, CH<sub>2</sub>F, CH<sub>2</sub>Cl, CH<sub>2</sub>OH, CN, C<sup>o</sup>CH, 1-propynyl, 2-propynyl, vinyl, OMe, OEt, SMe, or SEt.

52. A method of modulating the expression of a gene in a host cell comprising the gene to be modulated comprising the steps of:

- a) introducing into the host cell the gene expression modulation system of claim 11; and
- b) introducing into the host cell a ligand that independently combines with the ligand binding domains of the first polypeptide and the second polypeptide;
- 5 wherein the gene to be expressed is a component of a chimeric gene comprising:
- i) a response element to which the DNA binding domain from the first polypeptide binds;
  - ii) a promoter that is activated by the transactivation domain of the second polypeptide; and
  - 10 iii) a gene whose expression is to be modulated,
- whereby a complex is formed comprising the ligand, the first polypeptide, and the second polypeptide, and whereby the complex modulates expression of the gene in the host cell.

53. The method according to claim 52, wherein the ligand is a compound of the formula:



wherein:

- E is a (C<sub>4</sub>-C<sub>6</sub>)alkyl containing a tertiary carbon or a cyano(C<sub>3</sub>-C<sub>5</sub>)alkyl containing a tertiary carbon;
- R<sup>1</sup> is H, Me, Et, i-Pr, F, formyl, CF<sub>3</sub>, CHF<sub>2</sub>, CHCl<sub>2</sub>, CH<sub>2</sub>F, CH<sub>2</sub>Cl, CH<sub>2</sub>OH, CH<sub>2</sub>OMe, CH<sub>2</sub>CN, CN, C<sup>o</sup>CH, 1-propynyl, 2-propynyl, vinyl, OH, OMe, OEt, cyclopropyl, CF<sub>2</sub>CF<sub>3</sub>, CH=CHCN, allyl, azido, SCN, or SCHF<sub>2</sub>;
- 20 R<sup>2</sup> is H, Me, Et, n-Pr, i-Pr, formyl, CF<sub>3</sub>, CHF<sub>2</sub>, CHCl<sub>2</sub>, CH<sub>2</sub>F, CH<sub>2</sub>Cl, CH<sub>2</sub>OH, CH<sub>2</sub>OMe, CH<sub>2</sub>CN, CN, C<sup>o</sup>CH, 1-propynyl, 2-propynyl, vinyl, Ac, F, Cl, OH, OMe, OEt, O-n-Pr, OAc, NMe<sub>2</sub>, NEt<sub>2</sub>, SMe, SEt, SOCF<sub>3</sub>, OCF<sub>2</sub>CF<sub>2</sub>H, COEt, cyclopropyl, CF<sub>2</sub>CF<sub>3</sub>, CH=CHCN, allyl, azido, OCF<sub>3</sub>, OCHF<sub>2</sub>, O-i-Pr, SCN, SCHF<sub>2</sub>, SOMe, NH-CN, or
- 25 joined with R<sup>3</sup> and the phenyl carbons to which R<sup>2</sup> and R<sup>3</sup> are attached to form an ethylenedioxy, a dihydrofuryl ring with the oxygen adjacent to a phenyl carbon, or a dihydropyryl ring with the oxygen adjacent to a phenyl carbon;



R<sup>3</sup> is H, Et, or joined with R<sup>2</sup> and the phenyl carbons to which R<sup>2</sup> and R<sup>3</sup> are attached to form an ethylenedioxy, a dihydrofuryl ring with the oxygen adjacent to a phenyl carbon, or a dihydropyryl ring with the oxygen adjacent to a phenyl carbon;  
R<sup>4</sup>, R<sup>5</sup>, and R<sup>6</sup> are independently H, Me, Et, F, Cl, Br, formyl, CF<sub>3</sub>, CHF<sub>2</sub>, CHCl<sub>2</sub>,  
5 CH<sub>2</sub>F, CH<sub>2</sub>Cl, CH<sub>2</sub>OH, CN, C<sup>o</sup>CH, 1-propynyl, 2-propynyl, vinyl, OMe, OEt, SMe, or SEt.

54. An isolated host cell into which the gene expression modulation system according to claim 1 has been introduced.

55. The isolated host cell according to claim 54, wherein the host cell is selected  
10 from the group consisting of a bacterial cell, a fungal cell, a yeast cell, a plant cell, an animal cell, and a mammalian cell.

56. The isolated host cell according to claim 55, wherein the host cell is a plant cell, a murine cell, or a human cell.

57. An isolated host cell into which the gene expression modulation system  
15 according to claim 5 has been introduced.

58. The isolated host cell according to claim 57, wherein the host cell is selected from the group consisting of a bacterial cell, a fungal cell, a yeast cell, a plant cell, an animal cell, and a mammalian cell.

59. The isolated host cell according to claim 58, wherein the host cell is a plant  
20 cell, a murine cell, or a human cell.

60. An isolated host cell into which the gene expression modulation system according to claim 11 has been introduced.

61. The isolated host cell according to claim 60, wherein the host cell is selected from the group consisting of a bacterial cell, a fungal cell, a yeast cell, a plant cell, an animal  
25 cell, and a mammalian cell.

62. The isolated host cell according to claim 61, wherein the host cell is a plant cell, a murine cell, or a human cell.

63. A non-human organism comprising a host cell into which the gene expression modulation system according to claim 1 has been introduced.

64. The non-human organism according to claim 63, wherein the non-human  
30 organism is selected from the group consisting of a bacterium, a fungus, a yeast, a plant, an animal, and a mammal.

65. The non-human organism according to claim 64, wherein the non-human

organism is selected from the group consisting of a plant, a mouse, a rat, a rabbit, a cat, a dog, a bovine, a goat, a pig, a horse, a sheep, a monkey, and a chimpanzee.

66. A non-human organism comprising a host cell into which the gene expression modulation system according to claim 5 has been introduced.

5 67. The non-human organism according to claim 66, wherein the non-human organism is selected from the group consisting of a bacterium, a fungus, a yeast, a plant, an animal, and a mammal.

68. The non-human organism according to claim 67, wherein the non-human organism is selected from the group consisting of a plant, a mouse, a rat, a rabbit, a cat, a dog,  
10 a bovine, a goat, a pig, a horse, a sheep, a monkey, and a chimpanzee.

69. A non-human organism comprising a host cell into which the gene expression modulation system according to claim 11 has been introduced.

70. The non-human organism according to claim 69, wherein the non-human organism is selected from the group consisting of a bacterium, a fungus, a yeast, a plant, an  
15 animal, and a mammal.

71. The non-human organism according to claim 70, wherein the non-human organism is selected from the group consisting of a plant, a mouse, a rat, a rabbit, a cat, a dog, a bovine, a goat, a pig, a horse, a sheep, a monkey, and a chimpanzee.

GAL4CfEcR GAL4DNABD CHeRDEF  
VP16RXR VP16AD MmRXRDEF  
pGAL4RELuc GAL4RE TATA

Figure 1

GAL4CfEcR GAL4DNABD CHeRDEF  
VP16USP VP16AD CUSPDEF  
pGAL4RELuc GAL4RE TATA

Figure 2

GAL4RXR GAL4DNAB MmRXRDEF  
VP16CfEcR VP16AD CHeRCDEF  
pGAL4RELuc GAL4RE TATA

Figure 3

GAL4RXR GAL4DNAB MmRXRDEF  
VP16DmEcR VP16AD DmEcRCDEF  
pGAL4RELu GAL4RE TATA Luciferase

Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10

Analysis of CfEcR Truncations with MmRXRE in 3T3 Cells




Figure 11

Analysis of CfEcR Truncations with MmRXRE in 3T3 Cells




Figure 12

## Analysis of MmRXR Truncations with C1EcRCDEF in 3T3 Cells




Figure 13




Figure 14




Figure 15

RH0020.ST25  
SEQUENCE LISTING

<110> Rohm and Haas Company  
Palli, Subba Reddy  
Kapitskaya, Marianna Zinovjevna  
Cress, Dean Ervin

<120> Novel Ecdysone Receptor-Based Inducible Gene Expression System

<130> RH0020

<150> 60/191,355  
<151> 2000-03-22

<150> 60/269,799  
<151> 2001-02-20

<160> 64

<170> PatentIn version 3.0

<210> 1  
<211> 1288  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> misc feature  
<223> Novel Sequence

<400> 1  
aaggggccctg cgccccgtca gcaagaggaa ctgtgtcttg tatgcgggga cagagcctcc 60  
ggataccact acaatgcgct cacgtgtgaa ggggtgtaaag ggttcttcag acggagtgtt 120  
accaaaaatg cggtttatat ttgtaaattc ggtcacgctt gcgaaatgga catgtacatg 180  
cgacggaaat gccaggagtg ccgcctgaag aagtgccttag ctgtaggcat gaggcctgag 240  
tgcgtagtac ccgagactca gtgcgccatg aagcggaaag agaagaaagc acagaaggag 300  
aaggacaaac tgctgtcag cagcagcagc gtggacgacc acatgccgcc cattatgcag 360  
tgtgaacctc cacctcctga agcagcaagg attcacgaag tgggcccaag gtttctctcc 420  
gacaagctgt tggagacaaa ccggcagaaa aacatcccc agttgacagc caaccagcag 480  
ttccttatcg ccagggtcat ctgggtaccag gacgggtacg agcagccttc tgatgaagat 540  
ttgaagagga ttacgcagac gtggcagcaa gcggacgatg aaaacgaaga gtctgacact 600  
cccttcgcgc agatcacaga gatgactatc ctcacggtec aacttatcgt ggagttcgcg 660  
aagggattgc cagggttcgc caagatctcg cagcctgac aaattacgct gcttaaggct 720  
tgctcaagtg aggtaatgat gctccgagtc gcgcgacgat acgatgcggc ctcagacagt 780  
gttctgttcg cgaacaacca agcgtacact cgcgacaact accgcaaggc tggcatggcc 840  
tacgtcatcg aggatctact gcacttctgc cgggtcatgt actctatggc gttggacaac 900  
atccattacg cgctgctcac ggctgtcgtc atcttttctg accggccagg gttggagcag 960  
ccgcaactgg ttgaagaaat ccagcggtag tacctgaata cgctccgcat ctatatcctg 1020



## RH0020.ST25

aaccagctga gcgggtcggc gcgttcgtcc gtcataatac gcaagatcct ctcaatcctc 1080  
tctgagctac gcacgctcgg catgcaaaac tccaacatgt gcattctcct caagctcaag 1140  
aacagaaagc tgccgccttt cctcgaggag atctgggatg tggcggacat gtcgcacacc 1200  
caaccgccgc ctatcctcga gtccccacag aatctctagc ccttcgcgcg acgcatcgcc 1260  
gatgccgcgt ccggccgcgc tgctctga 1288

<210> 2  
<211> 1110  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> misc\_feature  
<223> Novel Sequence

<400> 2  
gcggtttata tttgtaaatt cggtcacgct tgcgaaatgg acatgtacat gcgacggaaa 60  
tgccaggagt gccgcctgaa gaagtgttta gctgtaggca tgaggcctga gtgcgtagta 120  
cccagactc agtgcgccat gaagcggaaa gagaagaaag cacagaagga gaaggacaaa 180  
ctgcctgtca gcacgacgac ggtggacgac cacatgccgc ccattatgca gtgtgaacct 240  
ccacctcctg aagcagcaag gattcacgaa gtggtcccaa ggtttctctc cgacaagctg 300  
ttggagacaa accggcagaa aaacatcccc cagttgacag ccaaccagca gttccttata 360  
gccaggctca tctggtacca ggacgggtac gacgagcctt ctgatgaaga tttgaagagg 420  
attacgcaga cgtggcagca agcggacgat gaaaacgaag agtctgacac tcccttccgc 480  
cagatcacag agatgactat cctcacggtc caacttatcg tggagttcgc gaagggattg 540  
ccagggttcg ccaagatctc gcagcctgat caaattacgc tgcttaaggc ttgctcaagt 600  
gaggtaatga tgctccgagt cgcgcgacga tacgatgcgg cctcagacag tgttctgttc 660  
gcgaacaacc aagcgtagac tcgcgacaac taccgcaagg ctggcatggc ctacgtcatc 720  
gaggatctac tgcaattctg ccggtgcatg tactctatgg cgttggacaa catccattac 780  
gcgctgctca cggctgtcgt catcttttct gaccggccag ggttgagca gccgcaactg 840  
gtggaagaaa tccagcggta ctacctgaat acgctccgca tctatctcct gaaccagctg 900  
agcgggtcgg cgcgttcgtc cgtcatatac ggcaagatcc tctcaatcct ctctgagcta 960  
cgcacgctcg gcattcaaaa ctccaacatg tgcatctccc tcaagctcaa gaacagaaag 1020  
ctgcgcctt tctcgagga gatctgggat gtggcggaca tgcgcacac ccaaccgccg 1080  
cctatcctcg agtccccac gaattcttag 1110

<210> 3  
<211> 1054  
<212> DNA

WO 01/70816

PCT/US01/09050

RH0020.ST25

<213> Artificial Sequence

<220>

<221> misc\_feature

<223> Novel Sequence

<400> 3

```
cctgagtgcg tagtaccgga gactcagtgc gccatgaagc ggaaagagaa gaaagcacag    60
aaggagaagg acaaactgcc tgtcagcacg acgacgggtg acgaccacat gccgcccatt    120
atgcagtgtg aacctccacc tctgaagca gcaaggattc acgaagtggc cccaaggttt    180
ctctccgaca agctgttggg gacaaaccgg cagaaaaaca tccccagtt gacagccaac    240
cagcagttcc ttatgccag gctcatctgg taccaggacg ggtacgagca gccttctgat    300
gaagatttga agaggattac gcagacgtgg cagcaagcgg acgatgaaaa cgaagagtct    360
gacactccct tccgccagat cacagagatg actatcctca cgggtccaact tatcgtggag    420
ttcgcaagg gattgccagg gttcgccaag atctcgagc ctgatcaaat tacgctgctt    480
aaggcttgct caagtgggt aatgatgctc cgagtcgagc gacgatacga tgcggcctca    540
gacagtgttc tgttcgcgaa caaccaagcg tactctcg acaactaccg caaggctggc    600
atggcctacg tcatcgagga tctactgcac ttctgccggt gcatgtactc tatggcgttg    660
gacaacatcc attacgcgct gctcacggct gtcgtcatct tttctgaccg gccagggttg    720
gagcagcccg aactggtgga agaatccag cgggtactacc tgaatacgt cgcgcatctat    780
atcctgaacc agctgagcgg gtcggcgctg tcgtccgtca tatacggcaa gatcctctca    840
atcctctctg agctacgcac gctcgcatg caaaactcca acatgtgcat ctccctcaag    900
ctcaagaaca gaaagctgcc gcctttctc gaggagatct gggatgtggc ggacatgtcg    960
cacaccaac cgcgcctat cctcgagtcc cccacgaatc tctagcccct gcgcgcacgc   1020
atcgccgatg ccgcgtccgg ccgcgctgct ctga                               1054
```

<210> 4

<211> 735

<212> DNA

<213> Artificial Sequence

<220>

<221> misc\_feature

<223> Novel Sequence

<400> 4

```
taccaggacg ggtacgagca gccttctgat gaagatttga agaggattac gcagacgtgg    60
cagcaagcgg acgatgaaaa cgaagagtct gacactccct tccgccagat cacagagatg   120
actatcctca cgggtccaact tatcgtggag ttcgcaagg gattgccagg gttcgccaag   180
atctcgagc ctgatcaaat tacgctgctt aaggcttgct caagtgggt aatgatgctc   240
cgagtcgagc gacgatacga tgcggcctca gacagtgttc tgttcgcgaa caaccaagcg   300
```

RH0020.ST25

tacactcgcg acaactaccg caaggctggc atggcctacg tcatcgagga tctactgcac 360  
ttctgccggt gcatgtactc tatggcggtg gacaacatcc attacgcgct gctcacggct 420  
gtcgtcatct tttctgaccg gccagggttg gagcagccgc aactggtgga agaaatccag 480  
cgggtactacc tgaatacgct ccgcattctat atcctgaacc agctgagcgg gtcggcgcg 540  
tcgtccgtca tatacggcaa gatcctctca atcctctctg agctacgcac gtcggcgatg 600  
caaaactcca acatgtgcat ctccctcaag ctcaagaaca gaaagctgcc gcctttcttc 660  
gaggagatct gggatgtggc ggacatgtcg cacaccaac cgccgcctat cctcgagtcc 720  
cccacgaatc tctag 735

<210> 5  
<211> 960  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> misc\_feature  
<223> Novel Sequence

<400> 5  
cctgagtgcg tagtaccgga gactcagtgc gccatgaagc ggaaagagaa gaaagcacag 60  
aaggagaagg acaaaactgcc tgtcagcacg acgacgggtg acgaccacat gccgccatt 120  
atgcagtgtg aacctccacc tcctgaagca gcaaggattc acgaagtggc cccaaggttt 180  
ctctccgaca agctgttgga gacaaaccgg cagaaaaaca tccccagtt gacagccaac 240  
cagcagttcc ttatcgccag gctcatctgg taccaggacg ggtacgagca gccttctgat 300  
gaagatttga agaggattac gcagacgtgg cagcaagcgg acgatgaaaa cgaagagtct 360  
gacactccct tccgccagat cacagagatg actatcctca cggccaact tatcgtggag 420  
ttcgcaagg gattgccagg gttcgccaag atctcgagc ctgatcaaat tacgtgctt 480  
aaggcttgct caagtgggt aatgatgtc cgagtcgagc gacgatacga tgcggcctca 540  
gacagtgttc tgttcgcgaa caaccaagcg tacactcgcg acaactaccg caaggctggc 600  
atggcctacg tcatcgagga tctactgcac ttctgccggt gcatgtactc tatggcggtg 660  
gacaacatcc attacgcgct gctcacggct gtcgtcatct tttctgaccg gccagggttg 720  
gagcagccgc aactggtgga agaaatccag cgggtactacc tgaatacgct ccgcattctat 780  
atcctgaacc agctgagcgg gtcggcgcggt tcgtccgtca tatacggcaa gatcctctca 840  
atcctctctg agctacgcac gtcggcgatg caaaactcca acatgtgcat ctccctcaag 900  
ctcaagaaca gaaagctgcc gcctttcttc gaggagatct gggatgtggc ggacatgtcg 960

<210> 6  
<211> 1878  
<212> DNA

RH0020.ST25

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Novel Sequence

&lt;400&gt; 6

ggacctgccc caccgggtgca agaggagctg tgcctgggtt gcggcgacag ggcctccggc 60  
taccactaca acgccctcac ctgtgagggc tgcaaggggt tctttcgacg cagcgttacg 120  
aagagcgccg tctactgctg caagtccggg cgcgcctcgc aaatggacat gtacatgagg 180  
cgaaagtgtc aggagtgccc cctgaaaaag tgcctggccg tgggtatgcy gccggaatgc 240  
gtcgtcccg agaaccaatg tgcgatgaag cggcgcgaaa agaaggccca gaaggagaag 300  
gacaaaaatga ccacttcgcc gagctctcag catggcggca atggcagctt ggcctctggt 360  
ggcgccaag actttgttaa gaaggagatt cttgacctta tgacatgcga gccgccccag 420  
catgccacta ttccgctact acctgatgaa atattggcca agtgtcaagc gcgcaatata 480  
ccttccttaa cgtacaatca gttggccgtt atatacaagt taatttggtta ccaggatggc 540  
tatgagcagc catctgaaga ggatctcagg cgtataatga gtcaaccgca tgagaacgag 600  
agccaaacgg acgtcagctt tcggcatata accgagataa ccatactcac ggtccagttg 660  
attgttgagt ttgctaaagg tctaccagcy tttacaaaga taccocagga ggaccagatc 720  
acgttactaa aggcctgctc gtcggaggtg atgatgctgc gtatggcacg acgctatgac 780  
cacagctcgg actcaatatt ctccgcgaat aatagatcat atacgcggga ttcttataaa 840  
atggccggaa tggctgataa cattgaagac ctgctgcatt tctgccgcca aatgttctcg 900  
atgaaggtgg acaacgtcga atacgcgctt ctactgcca ttgtgatctt ctccgaccgg 960  
ccgggcctcg agaaggccca actagtcgaa gcgatccaga gctactacat cgacacgcta 1020  
cgcatattata tactcaaccg ccactgcggc gactcaatga gcctcgtctt ctacgcaaag 1080  
ctgctctcga tcctcaccga gctgcgtacg ctgggcaacc agaacgccga gatgtgtttc 1140  
tactaaagc tcaaaaaccg caaactgccc aagttcctcg aggagatctg ggacgttcat 1200  
gccatcccg catcgggtcca gtcgcacctt cagattaccc aggaggagaa cgagcgtctc 1260  
gagcgggctg agcgtatgcy ggcacggtt gggggcgcca ttaccgccg cattgattgc 1320  
gactctgctt ccacttcggc ggcggcagcc gcggcccagc atcagcctca gcctcagccc 1380  
cagccccaac cctcctccct gaccagaac gattcccagc accagacaca gccgcagcta 1440  
caacctcagc taccacctca gctgcaaggt caactgcaac cccagctcca accacagctt 1500  
cagacgcaac tccagccaca gattcaacca cagccacagc tcttcccggt ctccgctccc 1560  
gtgccgcct ccgtaaccg acctgggtcc ttgtccggc tcagtacgag cagcgaatac 1620  
atgggcggaa gtgcggccat aggacccatc acgccggcaa ccaccagcag tatcacggct 1680  
gccgttaccg ctagctccac cacatcagcy gtaccgatgg gcaacggagt tggagtcggt 1740

WO 01/70816

PCT/US01/09050

RH0020.ST25

gttggggtgg gcggcaacgt cagcatgtat gcgaacgccc agacggcgat ggccttgatg 1800  
 ggtgtagccc tgcattcgca ccaagagcag cttatcgggg gagtggcggg taagtcggag 1860  
 cactcgacga ctgcatag 1878

<210> 7  
 <211> 1752  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> misc feature  
 <223> Novel Sequence

<400> 7  
 gccgtctact gctgcaagtt cgggcgcgcc tgcgaaatgg acatgtacat gaggcgaaag 60  
 tgtcaggagt gccgcctgaa aaagtgcctg gccgtgggta tgcggccgga atgcgtcgtc 120  
 ccggagaacc aatgtgcgat gaagcggcgc gaaaagaagg ccagaagga gaaggacaaa 180  
 atgaccactt cgcgagctc tcagcatgyc ggcaatggca gcttggcctc tgggtggcggc 240  
 caagactttg ttaagaagga gattcttgac cttatgacat gcgagccgcc ccagcatgcc 300  
 actattccgc tactacctga tgaaatattg gccaaagtgc aagcgcgcaa tataccttcc 360  
 ttaacgtaca atcagttggc cgttatatac aagttaattt ggtaccagga tggctatgag 420  
 cagccatctg aagaggatct caggcgata atgagtcaac ccgatgagaa cgagagccaa 480  
 acggacgtca gctttcggca tataaccgag ataaccatac tcacggtcca gttgattgtt 540  
 gagtttgcta aaggtctacc agcgtttaca aagatacccc aggaggacca gatcacgtta 600  
 ctaaaggcct gctcgtcgga ggtgatgatg ctgctgatgg cacgacgcta tgaccacagc 660  
 tcggactcaa tattcttcgc gaataataga tcatatacgc gggattctta caaaatggcc 720  
 ggaatggctg ataacattga agacctgctg catctctgcc gccaaatgtt ctcgatgaag 780  
 gtggacaacg tcgaatacgc gcttctcact gccattgtga tcttctcgga ccggccgggc 840  
 ctggagaagg cccaactagt cgaagcgatc cagagctact acatcgacac gctacgcatt 900  
 tatatactca accgccaact cggcgactca atgagcctcg tcttctacgc aaagtgcctc 960  
 tcgatcctca ccgagctgag tacgctgggc aaccagaacg ccgagatgtg tttctcacta 1020  
 aagctcaaaa accgcaaact gcccaagttc ctcgaggaga tetgggacgt tcatgccatc 1080  
 ccgccatcgg tccagtcgca cttcagatt acccaggagg agaacgagcg tctcgagcgg 1140  
 gctgagccta tgcgggcata ggttgggggc gccattaccg ccggcattga ttgcgactct 1200  
 gcctccactt cggcggcggc agccgcggcc cagcatcagc ctcagcctca gccccagccc 1260  
 caaccctcct cctgaccca gaacgattcc cagcaccaga cacagccgca gctacaacct 1320  
 cagctaccac ctcagctgca aggtcaactg caaccccagc tccaaccaca gcttcagacg 1380

WO 01/70816

PCT/US01/09050

RH0020.ST25

caactccagc cacagattca accacagcca cagctccttc ccgtctccgc tcccgtgccc 1440  
gcctccgtaa ccgcacctgg ttccttgtec gcggtcagta cgagcagcga atacatgggc 1500  
ggaagtgcgg ccataggacc catcacgccg gcaaccacca gcagtatcac ggctgccgtt 1560  
accgctagct ccaccacatc agcggtagcg atgggcaacg gagttggagt cgggtgtggg 1620  
gtggggcgga acgtcagcat gtatgcgaac gccagacgg cgatggcctt gatgggtgta 1680  
gcctgcatt cgcaccaaga gcagcttata gggggagtgg cggttaagtc ggagcactcg 1740  
acgactgcat ag 1752

<210> 8  
<211> 1650  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> misc feature  
<223> Novel Sequence

<400> 8  
cggccggaat cgtcgtcccc ggagaaccaa tgtgcgatga agcggcgcca aaagaaggcc 60  
cagaaggaga aggacaaaat gaccacttcg ccgagctctc agcatggcgg caatggcagc 120  
ttggcctctg gtggcgccca agactttgtt aagaaggaga ttcttgacct tatgacatgc 180  
gagccgcccc agcatgccac tattccgcta ctacctgatg aaatattggc caagtgtcaa 240  
gcgcgaata taccttccctt aacgtacaat cagttggcgg ttatatacaa gttaatttgg 300  
taccaggatg gctatgagca gccatctgaa gaggatctca ggcgtataat gagtcaaccc 360  
gatgagaacg agagccaaac ggacgtcagc tttcggcata taaccgagat aaccatactc 420  
acgggtccagt tgattgttga gtttgctaaa ggtctaccag cgtttacaaa gataccccag 480  
gaggaccaga tcacgttact aaaggcctgc tcgtcggagg tgatgatgct gcgtatggca 540  
cgacgctatg accacagctc ggactcaata ttcttcgcca ataatagata atatacgcg 600  
gattcttaca aaatggccgg aatggctgat aacattgaag acctgctgca tttctgccgc 660  
caaatgttct cgtatgaagg ggacaacgtc gaatacgcgc ttctcaactgc cattgtgata 720  
ttctcggacc ggccgggctt ggagaaggcc caactagtcg aagcgatcca gagctactac 780  
atcgacacgc tacgcattta tatactcaac cgccactgcg gcgactcaat gagcctcgtc 840  
ttctacgcaa agctgctctc gatcctcacc gagctgcgta cgctgggcaa ccagaacgcc 900  
gagatgtgtt tctcactaaa gtcacaaaac cgcaactgc ccaagttcct cgaggagata 960  
tgggacgttc atgccatccc gccatcggtc cagtcgcacc ttcagattac ccaggaggag 1020  
aacgagcgtc tcgagcgggc tgagcgtatg cgggcatcgg ttggggcgcc cattaccgcc 1080  
ggcattgatt gcgactctgc ctccacttcg gcggcgccag ccgcggccca gcatcagcct 1140  
cagcctcagc ccagacccca accctcctcc ctgaccaga acgattccca gcaccagaca 1200

WO 01/70816

PCT/US01/09050

RH0020.ST25

cagccgcagc tacaacctca gctaccacct cagctgcaag gtcaactgca accccagctc 1260  
 caaccacagc ttcagacgca actccagcca cagattcaac cacagccaca gtccttccc 1320  
 gtctccgctc ccgtgccgcg ctccgtaacc gcacctggtt ccttgctccg ggtcagtagc 1380  
 agcagcgaat acatggggcg aagtgcggcc ataggaccca tcacgccggc aaccaccagc 1440  
 agtatcacgg ctgccgttac cgctagctcc accacatcag cggtagcgat gggcaacgga 1500  
 gttggagtcg gtgttggggg gggcggcaac gtcagcatgt atgcgaacgc ccagacggcg 1560  
 atggccttga tgggtgtagc cctgcattcg caccaagagc agcttatcgg gggagtggcg 1620  
 gttaagtcgg agcactcgac gactgcatag 1650

<210> 9  
 <211> 1338  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> misc feature  
 <223> Novel Sequence

<400> 9  
 tatgagcagc catctgaaga ggatctcagg cgtataatga gtcaaccoga tgagaacgag 60  
 agccaaacgg acgtcagctt tcggcatata accgagataa ccatactcac ggtccagttg 120  
 attgttgagt ttgctaaagg tctaccagcg tttaaaaga taccacagga ggaccagatc 180  
 acgttactaa aggcctgctc gtcggagggt atgatgctgc gtatggcacg acgctatgac 240  
 cacagctcgg actcaatatt ctccgcaat aatagatcat atacgcggga ttcttataaa 300  
 atggccggaa tggctgataa cattgaagac ctgctgcatt tctgccgcca aatgttctcg 360  
 atgaagggtg acaacgtcga atacgcgctt ctactgcca ttgtgatctt ctccgaccgg 420  
 ccgggccttg agaagcccca actagtcgaa gcgatccaga gctactacat cgacacgcta 480  
 cgcatttata tactcaacgg ccaactgcgg gactcaatga gcctcgtctt ctacgcaaag 540  
 ctgctctcga tcctcacga gctgcgtacg ctgggcaacc agaacgccga gatgtgtttc 600  
 tcactaaagc tcaaaaacgg caaactgccc aagttcctcg aggagatctg ggacgttcat 660  
 gccatcccg cctcgggtcca gtcgcacctt cagattaccc aggaggagaa cgagcgtctc 720  
 gagcgggctg agcgtatcgg ggcacgtgtt gggggcgcca ttaccgccgg cattgattgc 780  
 gactctgcct ccacttcggc ggcggcagcc gcggcccgag atcagcctca gcctcagccc 840  
 cagccccaac cctctccctt gaccagaaac gattcccgag accagacaca gccgcagcta 900  
 caacctcagc taccacctca gtgcaagggt caactgcaac ccagctcca accacagctt 960  
 cagacgcaac tccagccaca gattcaacca cagccacagc tccttccgt ctccgtccc 1020  
 gtgccgcct ccgtaaccgc acctgggttc ttgtcccggt tcagtacgag cagcgaatac 1080

WO 01/70816

PCT/US01/09050

RH0020.ST25

atgggcgga	gtgcgccat	aggacccatc	acgccggcaa	ccaccagcag	tatcacggct	1140
gccgttaccg	ctagctccac	cacatcagcg	gtaccgatgg	gcaacggagt	tgagatcggt	1200
gttggggtgg	gcggcaacgt	cagcatgtat	gcgaacgccc	agacggcgat	ggccttgatg	1260
ggtgtagccc	tgcatctgca	ccaagagcag	cttatcgggg	gagtggcggt	taagtcggag	1320
cactcgacga	ctgcatag					1338

<210> 10  
 <211> 969  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> misc feature  
 <223> Novel Sequence

<400> 10	
cgcccggaat	gcgtcgctccc
ggagaaccaa	tgtcgcatga
agcggcgcg	aaagaaggcc
60	
cagaaggaga	aggacaaaat
gaccacttcg	ccgagctctc
agcatggcgg	caatggcagc
120	
ttggcctctg	gtggcgccca
agactttgtt	aagaaggaga
ttcttgacct	tatgacatgc
180	
gagccgcccc	agcatgccac
tattccgcta	ctacctgatg
aaatattggc	caagtgtcaa
240	
gcgcgcaata	taccttcctt
aacgtacaat	cagttggccg
ttatatacaa	gttaatttgg
300	
taccaggatg	gctatgagca
gccatctgaa	gaggatctca
ggcgtataat	gagtcaaccc
360	
gatgagaacg	agagccaaac
ggacgtcagc	tttcggcata
taaccgagat	aaccatactc
420	
acgggtccagt	tgattgttga
gtttgctaaa	ggtctaccag
cgtttacaaa	gataccccag
480	
gaggaccaga	tcacgttact
aaaggcctgc	tcgtcggagg
tgatgatgct	gcgtatggca
540	
cgacgctatg	accacagctc
ggactcaata	ttcttcgcga
ataatagatc	atatacgcg
600	
gattcttaca	aaatggccgg
aatggctgat	aacattgaag
acctgctgca	tttctgccgc
660	
caaatgttct	cgatgaaggt
ggacaacgtc	gaatacgcgc
ttctcactgc	cattgtgatc
720	
ttctcggacc	ggccgggcct
ggagaaggcc	caactagtcg
aagcgatcca	gagctactac
780	
atcgacacgc	tacgcattta
tatactcaac	cgccactcgc
gcgactcaat	gagcctcgtc
840	
ttctacgcaa	agctgctctc
gatcctcacc	gagctgcgta
cgctgggcaa	ccagaacgcc
900	
gagatgtgtt	tctcactaaa
gctcaaaaac	cgcaaactgc
ccaagttcct	cgaggagatc
960	
tgggacgtt	
969	

<210> 11  
 <211> 412  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <221> misc feature  
 <223> Novel Sequence



WO 01/70816

PCT/US01/09050

RH0020.ST25

<400> 11

```

Lys Gly Pro Ala Pro Arg Gln Gln Glu Glu Leu Cys Leu Val Cys Gly
1          5          10          15
Asp Arg Ala Ser Gly Tyr His Tyr Asn Ala Leu Thr Cys Glu Gly Cys
20          25          30
Lys Gly Phe Phe Arg Arg Ser Val Thr Lys Asn Ala Val Tyr Ile Cys
35          40          45
Lys Phe Gly His Ala Cys Glu Met Asp Met Tyr Met Arg Arg Lys Cys
50          55          60
Gln Glu Cys Arg Leu Lys Lys Cys Leu Ala Val Gly Met Arg Pro Glu
65          70          75          80
Cys Val Val Pro Glu Thr Gln Cys Ala Met Lys Arg Lys Glu Lys Lys
85          90          95
Ala Gln Lys Glu Lys Asp Lys Leu Pro Val Ser Thr Thr Thr Val Asp
100         105         110
Asp His Met Pro Pro Ile Met Gln Cys Glu Pro Pro Pro Pro Glu Ala
115         120         125
Ala Arg Ile His Glu Val Val Pro Arg Phe Leu Ser Asp Lys Leu Leu
130         135         140
Glu Thr Asn Arg Gln Lys Asn Ile Pro Gln Leu Thr Ala Asn Gln Gln
145         150         155         160
Phe Leu Ile Ala Arg Leu Ile Trp Tyr Gln Asp Gly Tyr Glu Gln Pro
165         170         175
Ser Asp Glu Asp Leu Lys Arg Ile Thr Gln Thr Trp Gln Gln Ala Asp
180         185         190
Asp Glu Asn Glu Glu Ser Asp Thr Pro Phe Arg Gln Ile Thr Glu Met
195         200         205
Thr Ile Leu Thr Val Gln Leu Ile Val Glu Phe Ala Lys Gly Leu Pro
210         215         220
Gly Phe Ala Lys Ile Ser Gln Pro Asp Gln Ile Thr Leu Leu Lys Ala
225         230         235         240
Cys Ser Ser Glu Val Met Met Leu Arg Val Ala Arg Arg Tyr Asp Ala
245         250         255
Ala Ser Asp Ser Val Leu Phe Ala Asn Asn Gln Ala Tyr Thr Arg Asp
260         265         270
Asn Tyr Arg Lys Ala Gly Met Ala Tyr Val Ile Glu Asp Leu Leu His
275         280         285
Phe Cys Arg Cys Met Tyr Ser Met Ala Leu Asp Asn Ile His Tyr Ala
290         295         300
Leu Leu Thr Ala Val Val Ile Phe Ser Asp Arg Pro Gly Leu Glu Gln
305         310         315         320
Pro Gln Leu Val Glu Glu Ile Gln Arg Tyr Tyr Leu Asn Thr Leu Arg

```

WO 01/70816

PCT/US01/09050

RH0020.ST25

325 330 335  
 Ile Tyr Ile Leu Asn Gln Leu Ser Gly Ser Ala Arg Ser Ser Val Ile  
 340 345 350  
 Tyr Gly Lys Ile Leu Ser Ile Leu Ser Glu Leu Arg Thr Leu Gly Met  
 355 360 365  
 Gln Asn Ser Asn Met Cys Ile Ser Leu Lys Leu Lys Asn Arg Lys Leu  
 370 375 380  
 Pro Pro Phe Leu Glu Glu Ile Trp Asp Val Ala Asp Met Ser His Thr  
 385 390 395 400  
 Gln Pro Pro Pro Ile Leu Glu Ser Pro Thr Asn Leu  
 405 410

<210> 12  
 <211> 412  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <221> misc\_feature  
 <223> Novel Sequence

<400> 12

Lys Gly Pro Ala Pro Arg Gln Gln Glu Glu Leu Cys Leu Val Cys Gly  
 1 5 10 15  
 Asp Arg Ala Ser Gly Tyr His Tyr Asn Ala Leu Thr Cys Glu Gly Cys  
 20 25 30  
 Lys Gly Phe Phe Arg Arg Ser Val Thr Lys Asn Ala Val Tyr Ile Cys  
 35 40 45  
 Lys Phe Gly His Ala Cys Glu Met Asp Met Tyr Met Arg Arg Lys Cys  
 50 55 60  
 Gln Glu Cys Arg Leu Lys Lys Cys Leu Ala Val Gly Met Arg Pro Glu  
 65 70 75 80  
 Cys Val Val Pro Glu Thr Gln Cys Ala Met Lys Arg Lys Glu Lys Lys  
 85 90 95  
 Ala Gln Lys Glu Lys Asp Lys Leu Pro Val Ser Thr Thr Thr Val Asp  
 100 105 110  
 Asp His Met Pro Pro Ile Met Gln Cys Glu Pro Pro Pro Glu Ala  
 115 120 125  
 Ala Arg Ile His Glu Val Val Pro Arg Phe Leu Ser Asp Lys Leu Leu  
 130 135 140  
 Glu Thr Asn Arg Gln Lys Asn Ile Pro Gln Leu Thr Ala Asn Gln Gln  
 145 150 155 160  
 Phe Leu Ile Ala Arg Leu Ile Trp Tyr Gln Asp Gly Tyr Glu Gln Pro  
 165 170 175  
 Ser Asp Glu Asp Leu Lys Arg Ile Thr Gln Thr Trp Gln Ala Asp  
 180 185 190

WO 01/70816

PCT/US01/09050

RH0020.ST25

Asp Glu Asn Glu Glu Ser Asp Thr Pro Phe Arg Gln Ile Thr Glu Met  
195 200 205

Thr Ile Leu Thr Val Gln Leu Ile Val Glu Phe Ala Lys Gly Leu Pro  
210 215 220

Gly Phe Ala Lys Ile Ser Gln Pro Asp Gln Ile Thr Leu Leu Lys Ala  
225 230 235 240

Cys Ser Ser Glu Val Met Met Leu Arg Val Ala Arg Arg Tyr Asp Ala  
245 250 255

Ala Ser Asp Ser Val Leu Phe Ala Asn Asn Gln Ala Tyr Thr Arg Asp  
260 265 270

Asn Tyr Arg Lys Ala Gly Met Ala Tyr Val Ile Glu Asp Leu Leu His  
275 280 285

Phe Cys Arg Cys Met Tyr Ser Met Ala Leu Asp Asn Ile His Tyr Ala  
290 295 300

Leu Leu Thr Ala Val Val Ile Phe Ser Asp Arg Pro Gly Leu Glu Gln  
305 310 315 320

Pro Gln Leu Val Glu Glu Ile Gln Arg Tyr Tyr Leu Asn Thr Leu Arg  
325 330 335

Ile Tyr Ile Leu Asn Gln Leu Ser Gly Ser Ala Arg Ser Ser Val Ile  
340 345 350

Tyr Gly Lys Ile Leu Ser Ile Leu Ser Glu Leu Arg Thr Leu Gly Met  
355 360 365

Gln Asn Ser Asn Met Cys Ile Ser Leu Lys Leu Lys Asn Arg Lys Leu  
370 375 380

Pro Pro Phe Leu Glu Glu Ile Trp Asp Val Ala Asp Met Ser His Thr  
385 390 395 400

Gln Pro Pro Pro Ile Leu Glu Ser Pro Thr Asn Leu  
405 410

<210> 13  
<211> 334  
<212> PRT  
<213> Artificial Sequence

<220>  
<221> misc\_feature  
<223> Novel Sequence

<400> 13

Pro Glu Cys Val Val Pro Glu Thr Gln Cys Ala Met Lys Arg Lys Glu  
1 5 10 15

Lys Lys Ala Gln Lys Glu Lys Asp Lys Leu Pro Val Ser Thr Thr Thr  
20 25 30

Val Asp Asp His Met Pro Pro Ile Met Gln Cys Glu Pro Pro Pro Pro  
35 40 45

Glu Ala Ala Arg Ile His Glu Val Val Pro Arg Phe Leu Ser Asp Lys  
50 55 60

WO 01/70816

PCT/US01/09050

RH0020.ST25

```

Ieu Leu Glu Thr Asn Arg Gln Lys Asn Ile Pro Gln Leu Thr Ala Asn
65      70      75      80

Gln Gln Phe Leu Ile Ala Arg Leu Ile Trp Tyr Gln Asp Gly Tyr Glu
85      90      95

Gln Pro Ser Asp Glu Asp Leu Lys Arg Ile Thr Gln Thr Trp Gln Gln
100     105     110

Ala Asp Asp Glu Asn Glu Glu Ser Asp Thr Pro Phe Arg Gln Ile Thr
115     120     125

Glu Met Thr Ile Leu Thr Val Gln Leu Ile Val Glu Phe Ala Lys Gly
130     135     140

Leu Pro Gly Phe Ala Lys Ile Ser Gln Pro Asp Gln Ile Thr Leu Leu
145     150     155     160

Lys Ala Cys Ser Ser Glu Val Met Met Leu Arg Val Ala Arg Arg Tyr
165     170     175

Asp Ala Ala Ser Asp Ser Val Leu Phe Ala Asn Asn Gln Ala Tyr Thr
180     185     190

Arg Asp Asn Tyr Arg Lys Ala Gly Met Ala Tyr Val Ile Glu Asp Leu
195     200     205

Leu His Phe Cys Arg Cys Met Tyr Ser Met Ala Leu Asp Asn Ile His
210     215     220

Tyr Ala Leu Leu Thr Ala Val Val Ile Phe Ser Asp Arg Pro Gly Leu
225     230     235     240

Glu Gln Pro Gln Leu Val Glu Glu Ile Gln Arg Tyr Tyr Leu Asn Thr
245     250     255

Leu Arg Ile Tyr Ile Leu Asn Gln Leu Ser Gly Ser Ala Arg Ser Ser
260     265     270

Val Ile Tyr Gly Lys Ile Leu Ser Ile Leu Ser Glu Leu Arg Thr Leu
275     280     285

Gly Met Gln Asn Ser Asn Met Cys Ile Ser Leu Lys Leu Lys Asn Arg
290     295     300

Lys Leu Pro Pro Phe Leu Glu Glu Ile Trp Asp Val Ala Asp Met Ser
305     310     315     320

His Thr Gln Pro Pro Ile Leu Glu Ser Pro Thr Asn Leu
325     330

```

<210> 14  
 <211> 244  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <221> misc feature  
 <223> Novel Sequence

<400> 14

Tyr Gln Asp Gly Tyr Glu Gln Pro Ser Asp Glu Asp Leu Lys Arg Ile

WO 01/70816

PCT/US01/09050

RH0020.ST25

```

1           5           10           15
Thr Gln Thr Trp Gln Gln Ala Asp Asp Glu Asn Glu Glu Ser Asp Thr
      20      25      30
Pro Phe Arg Gln Ile Thr Glu Met Thr Ile Leu Thr Val Gln Leu Ile
      35      40      45
Val Glu Phe Ala Lys Gly Leu Pro Gly Phe Ala Lys Ile Ser Gln Pro
      50      55      60
Asp Gln Ile Thr Leu Leu Lys Ala Cys Ser Ser Glu Val Met Met Leu
      65      70      75      80
Arg Val Ala Arg Arg Tyr Asp Ala Ala Ser Asp Ser Val Leu Phe Ala
      85      90      95
Asn Asn Gln Ala Tyr Thr Arg Asp Asn Tyr Arg Lys Ala Gly Met Ala
      100     105     110
Tyr Val Ile Glu Asp Leu Leu His Phe Cys Arg Cys Met Tyr Ser Met
      115     120     125
Ala Leu Asp Asn Ile His Tyr Ala Leu Leu Thr Ala Val Val Ile Phe
      130     135     140
Ser Asp Arg Pro Gly Leu Glu Gln Pro Gln Leu Val Glu Glu Ile Gln
      145     150     155     160
Arg Tyr Tyr Leu Asn Thr Leu Arg Ile Tyr Ile Leu Asn Gln Leu Ser
      165     170     175
Gly Ser Ala Arg Ser Ser Val Ile Tyr Gly Lys Ile Leu Ser Ile Leu
      180     185     190
Ser Glu Leu Arg Thr Leu Gly Met Gln Asn Ser Asn Met Cys Ile Ser
      195     200     205
Leu Lys Leu Lys Asn Arg Lys Leu Pro Pro Phe Leu Glu Glu Ile Trp
      210     215     220
Asp Val Ala Asp Met Ser His Thr Gln Pro Pro Pro Ile Leu Glu Ser
      225     230     235     240
Pro Thr Asn Leu

```

```

<210> 15
<211> 320
<212> PRT
<213> Artificial Sequence

<220>
<221> misc feature
<223> Novel Sequence

```

<400> 15

```

Pro Glu Cys Val Val Pro Glu Thr Gln Cys Ala Met Lys Arg Lys Glu
1           5           10           15
Lys Lys Ala Gln Lys Glu Lys Asp Lys Leu Pro Val Ser Thr Thr Thr
      20      25      30

```

WO 01/70816

PCT/US01/09050

RH0020.ST25

Val Asp Asp His Met Pro Pro Ile Met Gln Cys Glu Pro Pro Pro  
35 40 45

Glu Ala Ala Arg Ile His Glu Val Val Pro Arg Phe Leu Ser Asp Lys  
50 55 60

Leu Leu Glu Thr Asn Arg Gln Lys Asn Ile Pro Gln Leu Thr Ala Asn  
65 70 75 80

Gln Gln Phe Leu Ile Ala Arg Leu Ile Trp Tyr Gln Asp Gly Tyr Glu  
85 90 95

Gln Pro Ser Asp Glu Asp Leu Lys Arg Ile Thr Gln Thr Trp Gln Gln  
100 105 110

Ala Asp Asp Glu Asn Glu Glu Ser Asp Thr Pro Phe Arg Gln Ile Thr  
115 120 125

Glu Met Thr Ile Leu Thr Val Gln Leu Ile Val Glu Phe Ala Lys Gly  
130 135 140

Leu Pro Gly Phe Ala Lys Ile Ser Gln Pro Asp Gln Ile Thr Leu Leu  
145 150 155 160

Lys Ala Cys Ser Ser Glu Val Met Met Leu Arg Val Ala Arg Arg Tyr  
165 170 175

Asp Ala Ala Ser Asp Ser Val Leu Phe Ala Asn Asn Gln Ala Tyr Thr  
180 185 190

Arg Asp Asn Tyr Arg Lys Ala Gly Met Ala Tyr Val Ile Glu Asp Leu  
195 200 205

Leu His Phe Cys Arg Cys Met Tyr Ser Met Ala Leu Asp Asn Ile His  
210 215 220

Tyr Ala Leu Leu Thr Ala Val Val Ile Phe Ser Asp Arg Pro Gly Leu  
225 230 235 240

Glu Gln Pro Gln Leu Val Glu Glu Ile Gln Arg Tyr Tyr Leu Asn Thr  
245 250 255

Leu Arg Ile Tyr Ile Leu Asn Gln Leu Ser Gly Ser Ala Arg Ser Ser  
260 265 270

Val Ile Tyr Gly Lys Ile Leu Ser Ile Leu Ser Glu Leu Arg Thr Leu  
275 280 285

Gly Met Gln Asn Ser Asn Met Cys Ile Ser Leu Lys Leu Lys Asn Arg  
290 295 300

Lys Leu Pro Pro Phe Leu Glu Glu Ile Trp Asp Val Ala Asp Met Ser  
305 310 315 320

<210> 16

<211> 625

<212> PRT

<213> Artificial Sequence

<220>

<221> misc feature

<223> Novel Sequence

<400> 16

WO 01/70816

PCT/US01/09050

RH0020.ST25

Gly Pro Ala Pro Arg Val Gln Glu Glu Leu Cys Leu Val Cys Gly Asp  
1 5 10 15

Arg Ala Ser Gly Tyr His Tyr Asn Ala Leu Thr Cys Glu Gly Cys Lys  
20 25 30

Gly Phe Phe Arg Arg Ser Val Thr Lys Ser Ala Val Tyr Cys Cys Lys  
35 40 45

Phe Gly Arg Ala Cys Glu Met Asp Met Tyr Met Arg Arg Lys Cys Gln  
50 55 60

Glu Cys Arg Leu Lys Lys Cys Leu Ala Val Gly Met Arg Pro Glu Cys  
65 70 75 80

Val Val Pro Glu Asn Gln Cys Ala Met Lys Arg Arg Glu Lys Lys Ala  
85 90 95

Gln Lys Glu Lys Asp Lys Met Thr Thr Ser Pro Ser Ser Gln His Gly  
100 105 110

Gly Asn Gly Ser Leu Ala Ser Gly Gly Gly Gln Asp Phe Val Lys Lys  
115 120 125

Glu Ile Leu Asp Leu Met Thr Cys Glu Pro Pro Gln His Ala Thr Ile  
130 135 140

Pro Leu Leu Pro Asp Glu Ile Leu Ala Lys Cys Gln Ala Arg Asn Ile  
145 150 155 160

Pro Ser Leu Thr Tyr Asn Gln Leu Ala Val Ile Tyr Lys Leu Ile Trp  
165 170 175

Tyr Gln Asp Gly Tyr Glu Gln Pro Ser Glu Glu Asp Leu Arg Arg Ile  
180 185 190

Met Ser Gln Pro Asp Glu Asn Glu Ser Gln Thr Asp Val Ser Phe Arg  
195 200 205

His Ile Thr Glu Ile Thr Ile Leu Thr Val Gln Leu Ile Val Glu Phe  
210 215 220

Ala Lys Gly Leu Pro Ala Phe Thr Lys Ile Pro Gln Glu Asp Gln Ile  
225 230 235 240

Thr Leu Leu Lys Ala Cys Ser Ser Glu Val Met Met Leu Arg Met Ala  
245 250 255

Arg Arg Tyr Asp His Ser Ser Asp Ser Ile Phe Phe Ala Asn Asn Arg  
260 265 270

Ser Tyr Thr Arg Asp Ser Tyr Lys Met Ala Gly Met Ala Asp Asn Ile  
275 280 285

Glu Asp Leu Leu His Phe Cys Arg Gln Met Phe Ser Met Lys Val Asp  
290 295 300

Asn Val Glu Tyr Ala Leu Leu Thr Ala Ile Val Ile Phe Ser Asp Arg  
305 310 315 320

Pro Gly Leu Glu Lys Ala Gln Leu Val Glu Ala Ile Gln Ser Tyr Tyr  
325 330 335

Ile Asp Thr Leu Arg Ile Tyr Ile Leu Asn Arg His Cys Gly Asp Ser

WO 01/70816

PCT/US01/09050

RH0020.ST25

340	345	350
Met Ser Leu Val Phe Tyr Ala Lys Leu Leu Ser Ile Leu Thr Glu Leu		
355	360	365
Arg Thr Leu Gly Asn Gln Asn Ala Glu Met Cys Phe Ser Leu Lys Leu		
370	375	380
Lys Asn Arg Lys Leu Pro Lys Phe Leu Glu Glu Ile Trp Asp Val His		
385	390	400
Ala Ile Pro Pro Ser Val Gln Ser His Leu Gln Ile Thr Gln Glu Glu		
405	410	415
Asn Glu Arg Leu Glu Arg Ala Glu Arg Met Arg Ala Ser Val Gly Gly		
420	425	430
Ala Ile Thr Ala Gly Ile Asp Cys Asp Ser Ala Ser Thr Ser Ala Ala		
435	440	445
Ala Ala Ala Ala Gln His Gln Pro Gln Pro Gln Pro Gln Pro Gln Pro		
450	455	460
Ser Ser Leu Thr Gln Asn Asp Ser Gln His Gln Thr Gln Pro Gln Leu		
465	470	475
Gln Pro Gln Leu Pro Pro Gln Leu Gln Gly Gln Leu Gln Pro Gln Leu		
485	490	495
Gln Pro Gln Leu Gln Thr Gln Leu Gln Pro Gln Ile Gln Pro Gln Pro		
500	505	510
Gln Leu Leu Pro Val Ser Ala Pro Val Pro Ala Ser Val Thr Ala Pro		
515	520	525
Gly Ser Leu Ser Ala Val Ser Thr Ser Ser Glu Tyr Met Gly Gly Ser		
530	535	540
Ala Ala Ile Gly Pro Ile Thr Pro Ala Thr Thr Ser Ser Ile Thr Ala		
545	550	555
Ala Val Thr Ala Ser Ser Thr Thr Ser Ala Val Pro Met Gly Asn Gly		
565	570	575
Val Gly Val Gly Val Gly Val Gly Gly Asn Val Ser Met Tyr Ala Asn		
580	585	590
Ala Gln Thr Ala Met Ala Leu Met Gly Val Ala Leu His Ser His Gln		
595	600	605
Glu Gln Leu Ile Gly Gly Val Ala Val Lys Ser Glu His Ser Thr Thr		
610	615	620

Ala  
625

<210> 17

<211> 583

<212> PRT

<213> Artificial Sequence

<220>

<221> misc feature

<223> Novel Sequence



WO 01/70816

PCT/US01/09050

RH0020.ST25

<400> 17

Ala Val Tyr Cys Cys Lys Phe Gly Arg Ala Cys Glu Met Asp Met Tyr  
1 5 10 15  
Met Arg Arg Lys Cys Gln Glu Cys Arg Leu Lys Lys Cys Leu Ala Val  
20 25 30  
Gly Met Arg Pro Glu Cys Val Val Pro Glu Asn Gln Cys Ala Met Lys  
35 40 45  
Arg Arg Glu Lys Lys Ala Gln Lys Glu Lys Asp Lys Met Thr Thr Ser  
50 55 60  
Pro Ser Ser Gln His Gly Gly Asn Gly Ser Leu Ala Ser Gly Gly Gly  
65 70 75 80  
Gln Asp Phe Val Lys Lys Glu Ile Leu Asp Leu Met Thr Cys Glu Pro  
85 90 95  
Pro Gln His Ala Thr Ile Pro Leu Leu Pro Asp Glu Ile Leu Ala Lys  
100 105 110  
Cys Gln Ala Arg Asn Ile Pro Ser Leu Thr Tyr Asn Gln Leu Ala Val  
115 120 125  
Ile Tyr Lys Leu Ile Trp Tyr Gln Asp Gly Tyr Glu Gln Pro Ser Glu  
130 135 140  
Glu Asp Leu Arg Arg Ile Met Ser Gln Pro Asp Glu Asn Glu Ser Gln  
145 150 155 160  
Thr Asp Val Ser Phe Arg His Ile Thr Glu Ile Thr Ile Leu Thr Val  
165 170 175  
Gln Leu Ile Val Glu Phe Ala Lys Gly Leu Pro Ala Phe Thr Lys Ile  
180 185 190  
Pro Gln Glu Asp Gln Ile Thr Leu Leu Lys Ala Cys Ser Ser Glu Val  
195 200 205  
Met Met Leu Arg Met Ala Arg Arg Tyr Asp His Ser Ser Asp Ser Ile  
210 215 220  
Phe Phe Ala Asn Asn Arg Ser Tyr Thr Arg Asp Ser Tyr Lys Met Ala  
225 230 235 240  
Gly Met Ala Asp Asn Ile Glu Asp Leu Leu His Phe Cys Arg Gln Met  
245 250 255  
Phe Ser Met Lys Val Asp Asn Val Glu Tyr Ala Leu Leu Thr Ala Ile  
260 265 270  
Val Ile Phe Ser Asp Arg Pro Gly Leu Glu Lys Ala Gln Leu Val Glu  
275 280 285  
Ala Ile Gln Ser Tyr Tyr Ile Asp Thr Leu Arg Ile Tyr Ile Leu Asn  
290 295 300  
Arg His Cys Gly Asp Ser Met Ser Leu Val Phe Tyr Ala Lys Leu Leu  
305 310 315 320  
Ser Ile Leu Thr Glu Leu Arg Thr Leu Gly Asn Gln Asn Ala Glu Met  
325 330 335

WO 01/70816

PCT/US01/09050

RH0020.ST25

Cys Phe Ser Leu Lys Leu Lys Asn Arg Lys Leu Pro Lys Phe Leu Glu  
340 345 350  
Glu Ile Trp Asp Val His Ala Ile Pro Pro Ser Val Gln Ser His Leu  
355 360 365  
Gln Ile Thr Gln Glu Glu Asn Glu Arg Leu Glu Arg Ala Glu Arg Met  
370 375 380  
Arg Ala Ser Val Gly Gly Ala Ile Thr Ala Gly Ile Asp Cys Asp Ser  
385 390 395 400  
Ala Ser Thr Ser Ala Ala Ala Ala Ala Ala Gln His Gln Pro Gln Pro  
405 410 415  
Gln Pro Gln Pro Gln Pro Ser Ser Leu Thr Gln Asn Asp Ser Gln His  
420 425 430  
Gln Thr Gln Pro Gln Leu Gln Pro Gln Leu Pro Pro Gln Leu Gln Gly  
435 440 445  
Gln Leu Gln Pro Gln Leu Gln Pro Gln Leu Gln Thr Gln Leu Gln Pro  
450 455 460  
Gln Ile Gln Pro Gln Pro Gln Leu Leu Pro Val Ser Ala Pro Val Pro  
465 470 475 480  
Ala Ser Val Thr Ala Pro Gly Ser Leu Ser Ala Val Ser Thr Ser Ser  
485 490 495  
Glu Tyr Met Gly Gly Ser Ala Ala Ile Gly Pro Ile Thr Pro Ala Thr  
500 505 510  
Thr Ser Ser Ile Thr Ala Ala Val Thr Ala Ser Ser Thr Thr Ser Ala  
515 520 525  
Val Pro Met Gly Asn Gly Val Gly Val Gly Val Gly Val Gly Gly Asn  
530 535 540  
Val Ser Met Tyr Ala Asn Ala Gln Thr Ala Met Ala Leu Met Gly Val  
545 550 555 560  
Ala Leu His Ser His Gln Glu Gln Leu Ile Gly Gly Val Ala Val Lys  
565 570 575  
Ser Glu His Ser Thr Thr Ala  
580

<210> 18  
<211> 549  
<212> PRT  
<213> Artificial Sequence

<400> 18

Arg Pro Glu Cys Val Val Pro Glu Asn Gln Cys Ala Met Lys Arg Arg  
1 5 10 15  
Glu Lys Lys Ala Gln Lys Glu Lys Asp Lys Met Thr Thr Ser Pro Ser  
20 25 30  
Ser Gln His Gly Gly Asn Gly Ser Leu Ala Ser Gly Gly Gly Gln Asp  
35 40 45

WO 01/70816

PCT/US01/09050

RH0020.ST25

Phe Val Lys Lys Glu Ile Leu Asp Leu Met Thr Cys Glu Pro Pro Gln  
50 55 60

His Ala Thr Ile Pro Leu Leu Pro Asp Glu Ile Leu Ala Lys Cys Gln  
65 70 75 80

Ala Arg Asn Ile Pro Ser Leu Thr Tyr Asn Gln Leu Ala Val Ile Tyr  
85 90 95

Lys Leu Ile Trp Tyr Gln Asp Gly Tyr Glu Gln Pro Ser Glu Glu Asp  
100 105 110

Leu Arg Arg Ile Met Ser Gln Pro Asp Glu Asn Glu Ser Gln Thr Asp  
115 120 125

Val Ser Phe Arg His Ile Thr Glu Ile Thr Ile Leu Thr Val Gln Leu  
130 135 140

Ile Val Glu Phe Ala Lys Gly Leu Pro Ala Phe Thr Lys Ile Pro Gln  
145 150 155 160

Glu Asp Gln Ile Thr Leu Leu Lys Ala Cys Ser Ser Glu Val Met Met  
165 170 175

Leu Arg Met Ala Arg Arg Tyr Asp His Ser Ser Asp Ser Ile Phe Phe  
180 185 190

Ala Asn Asn Arg Ser Tyr Thr Arg Asp Ser Tyr Lys Met Ala Gly Met  
195 200 205

Ala Asp Asn Ile Glu Asp Leu Leu His Phe Cys Arg Gln Met Phe Ser  
210 215 220

Met Lys Val Asp Asn Val Glu Tyr Ala Leu Leu Thr Ala Ile Val Ile  
225 230 235 240

Phe Ser Asp Arg Pro Gly Leu Glu Lys Ala Gln Leu Val Glu Ala Ile  
245 250 255

Gln Ser Tyr Tyr Ile Asp Thr Leu Arg Ile Tyr Ile Leu Asn Arg His  
260 265 270

Cys Gly Asp Ser Met Ser Leu Val Phe Tyr Ala Lys Leu Leu Ser Ile  
275 280 285

Leu Thr Glu Leu Arg Thr Leu Gly Asn Gln Asn Ala Glu Met Cys Phe  
290 295 300

Ser Leu Lys Leu Lys Asn Arg Lys Leu Pro Lys Phe Leu Glu Glu Ile  
305 310 315 320

Trp Asp Val His Ala Ile Pro Pro Ser Val Gln Ser His Leu Gln Ile  
325 330 335

Thr Gln Glu Glu Asn Glu Arg Leu Glu Arg Ala Glu Arg Met Arg Ala  
340 345 350

Ser Val Gly Gly Ala Ile Thr Ala Gly Ile Asp Cys Asp Ser Ala Ser  
355 360 365

Thr Ser Ala Ala Ala Ala Ala Gln His Gln Pro Gln Pro Gln Pro  
370 375 380

Gln Pro Gln Pro Ser Ser Leu Thr Gln Asn Asp Ser Gln His Gln Thr  
385 390 395 400

WO 01/70816

PCT/US01/09050

RH0020.ST25

Gln Pro Gln Leu Gln Pro Gln Leu Pro Pro Gln Leu Gln Gly Gln Leu  
405 410 415

Gln Pro Gln Leu Gln Pro Gln Leu Gln Thr Gln Leu Gln Pro Gln Ile  
420 425 430

Gln Pro Gln Pro Gln Leu Leu Pro Val Ser Ala Pro Val Pro Ala Ser  
435 440 445

Val Thr Ala Pro Gly Ser Leu Ser Ala Val Ser Thr Ser Ser Glu Tyr  
450 455 460

Met Gly Gly Ser Ala Ala Ile Gly Pro Ile Thr Pro Ala Thr Thr Ser  
465 470 475 480

Ser Ile Thr Ala Ala Val Thr Ala Ser Ser Thr Thr Ser Ala Val Pro  
485 490 495

Met Gly Asn Gly Val Gly Val Gly Val Gly Val Gly Gly Asn Val Ser  
500 505 510

Met Tyr Ala Asn Ala Gln Thr Ala Met Ala Leu Met Gly Val Ala Leu  
515 520 525

His Ser His Gln Glu Gln Leu Ile Gly Gly Val Ala Val Lys Ser Glu  
530 535 540

His Ser Thr Thr Ala  
545

<210> 19  
<211> 445  
<212> PRT  
<213> Artificial Sequence

<400> 19

Tyr Glu Gln Pro Ser Glu Glu Asp Leu Arg Arg Ile Met Ser Gln Pro  
1 5 10 15

Asp Glu Asn Glu Ser Gln Thr Asp Val Ser Phe Arg His Ile Thr Glu  
20 25 30

Ile Thr Ile Leu Thr Val Gln Leu Ile Val Glu Phe Ala Lys Gly Leu  
35 40 45

Pro Ala Phe Thr Lys Ile Pro Gln Glu Asp Gln Ile Thr Leu Leu Lys  
50 55 60

Ala Cys Ser Ser Glu Val Met Met Leu Arg Met Ala Arg Arg Tyr Asp  
65 70 75 80

His Ser Ser Asp Ser Ile Phe Phe Ala Asn Asn Arg Ser Tyr Thr Arg  
85 90 95

Asp Ser Tyr Lys Met Ala Gly Met Ala Asp Asn Ile Glu Asp Leu Leu  
100 105 110

His Phe Cys Arg Gln Met Phe Ser Met Lys Val Asp Asn Val Glu Tyr  
115 120 125

Ala Leu Leu Thr Ala Ile Val Ile Phe Ser Asp Arg Pro Gly Leu Glu  
130 135 140

WO 01/70816

PCT/US01/09050

RH0020.ST25

Lys Ala Gln Leu Val Glu Ala Ile Gln Ser Tyr Tyr Ile Asp Thr Leu  
145 150 155 160

Arg Ile Tyr Ile Leu Asn Arg His Cys Gly Asp Ser Met Ser Leu Val  
165 170 175

Phe Tyr Ala Lys Leu Leu Ser Ile Leu Thr Glu Leu Arg Thr Leu Gly  
180 185 190

Asn Gln Asn Ala Glu Met Cys Phe Ser Leu Lys Leu Lys Asn Arg Lys  
195 200 205

Leu Pro Lys Phe Leu Glu Glu Ile Trp Asp Val His Ala Ile Pro Pro  
210 215 220

Ser Val Gln Ser His Leu Gln Ile Thr Gln Glu Glu Asn Glu Arg Leu  
225 230 235 240

Glu Arg Ala Glu Arg Met Arg Ala Ser Val Gly Gly Ala Ile Thr Ala  
245 250 255

Gly Ile Asp Cys Asp Ser Ala Ser Thr Ser Ala Ala Ala Ala Ala Ala  
260 265 270

Gln His Gln Pro Gln Pro Gln Pro Gln Pro Gln Pro Ser Ser Leu Thr  
275 280 285

Gln Asn Asp Ser Gln His Gln Thr Gln Pro Gln Leu Gln Pro Gln Leu  
290 295 300

Pro Pro Gln Leu Gln Gly Gln Leu Gln Pro Gln Leu Gln Pro Gln Leu  
305 310 315 320

Gln Thr Gln Leu Gln Pro Gln Ile Gln Pro Gln Pro Gln Leu Leu Pro  
325 330 335

Val Ser Ala Pro Val Pro Ala Ser Val Thr Ala Pro Gly Ser Leu Ser  
340 345 350

Ala Val Ser Thr Ser Ser Glu Tyr Met Gly Gly Ser Ala Ala Ile Gly  
355 360 365

Pro Ile Thr Pro Ala Thr Thr Ser Ser Ile Thr Ala Ala Val Thr Ala  
370 375 380

Ser Ser Thr Thr Ser Ala Val Pro Met Gly Asn Gly Val Gly Val Gly  
385 390 395 400

Val Gly Val Gly Gly Asn Val Ser Met Tyr Ala Asn Ala Gln Thr Ala  
405 410 415

Met Ala Leu Met Gly Val Ala Leu His Ser His Gln Glu Gln Leu Ile  
420 425 430

Gly Gly Val Ala Val Lys Ser Glu His Ser Thr Thr Ala  
435 440 445

<210> 20

<211> 323

<212> PRT

<213> Artificial Sequence

<400> 20

Arg Pro Glu Cys Val Val Pro Glu Asn Gln Cys Ala Met Lys Arg Arg

WO 01/70816

PCT/US01/09050

RH0020.ST25

1 5 10 15  
 Leu Lys Lys Ala Gln Lys Glu Lys Asp Lys Met Thr Thr Ser Pro Ser  
 20 25 30  
 Ser Gln His Gly Gly Asn Gly Ser Leu Ala Ser Gly Gly Gly Gln Asp  
 35 40 45  
 Phe Val Lys Lys Glu Ile Leu Asp Leu Met Thr Cys Glu Pro Pro Gln  
 50 55 60  
 His Ala Thr Ile Pro Leu Leu Pro Asp Glu Ile Leu Ala Lys Cys Gln  
 65 70 75 80  
 Ala Arg Asn Ile Pro Ser Leu Thr Tyr Asn Gln Leu Ala Val Ile Tyr  
 85 90 95  
 Lys Leu Ile Trp Tyr Gln Asp Gly Tyr Glu Gln Pro Ser Glu Glu Asp  
 100 105 110  
 Leu Arg Arg Ile Met Ser Gln Pro Asp Glu Asn Glu Ser Gln Thr Asp  
 115 120 125  
 Val Ser Phe Arg His Ile Thr Glu Ile Thr Ile Leu Thr Val Gln Leu  
 130 135 140  
 Ile Val Glu Phe Ala Lys Gly Leu Pro Ala Phe Thr Lys Ile Pro Gln  
 145 150 155 160  
 Glu Asp Gln Ile Thr Leu Leu Lys Ala Cys Ser Ser Glu Val Met Met  
 165 170 175  
 Leu Arg Met Ala Arg Arg Tyr Asp His Ser Ser Asp Ser Ile Phe Phe  
 180 185 190  
 Ala Asn Asn Arg Ser Tyr Thr Arg Asp Ser Tyr Lys Met Ala Gly Met  
 195 200 205  
 Ala Asp Asn Ile Glu Asp Leu Leu His Phe Cys Arg Gln Met Phe Ser  
 210 215 220  
 Met Lys Val Asp Asn Val Glu Tyr Ala Leu Leu Thr Ala Ile Val Ile  
 225 230 235 240  
 Phe Ser Asp Arg Pro Gly Leu Glu Lys Ala Gln Leu Val Glu Ala Ile  
 245 250 255  
 Gln Ser Tyr Tyr Ile Asp Thr Leu Arg Ile Tyr Ile Leu Asn Arg His  
 260 265 270  
 Cys Gly Asp Ser Met Ser Leu Val Phe Tyr Ala Lys Leu Leu Ser Ile  
 275 280 285  
 Leu Thr Glu Leu Arg Thr Leu Gly Asn Gln Asn Ala Glu Met Cys Phe  
 290 295 300  
 Ser Leu Lys Leu Lys Asn Arg Lys Leu Pro Lys Phe Leu Glu Glu Ile  
 305 310 315 320

Trp Asp Val

<210> 21  
 <211> 987  
 <212> DNA

RH0020.ST25

## &lt;213&gt; Artificial Sequence

&lt;400&gt; 21

```

tgtgctatct gtggggaccg ctctcagggc aaacactatg gggatacag ttgtgagggc   60
tgcaagggct tcttcaagag gacagtacgc aaagacctga cctacacctg ccgagacaac   120
aaggactgcc tgatcgacaa gagacagcgg aaccgggtgc agtactgccg ctaccagaag   180
tgcttgccca tgggcatgaa gcgggaagct gtgcaggagg agcggcagcg gggcaaggac   240
cggaatgaga acgaggtgga gtccaccagc agtgccaacg aggacatgcc tgtagagaag   300
attctggaag ccgagcttgc tgcgagccc aagactgaga catacgtgga ggcaaacaatg   360
gggctgaacc ccagctcacc aaatgaccct gttaccaaca tctgtcaagc agcagacaag   420
cagctcttca ctcttgtgga gtgggccaag aggatccac acttttctga gctgcccta   480
gacgaccagg tcctctgct acgggcaggc tggaaacgagc tgcgtatcgc ctcttctcc   540
caccgctcca tagctgtgaa agatgggatt ctctggcca ccggcctgca cgtacaccgg   600
aacagcgctc acagtgttg ggtggcgcc atctttgaca ggggtgtaac agagctggtg   660
tctaagatgc gtgacatgca gatggacaag acggagctgg gctgcctgag agccattgtc   720
ctgttcaacc ctgactctaa ggggtctca aacctgctg aggtggaggc gttgagggag   780
aaggtgtatg cgtactaga agcgtactgc aaacacaagt accctgagca gccgggcagg   840
tttgccaagc tgcgtccg cctgcctgca ctgcgttcca tcgggctcaa gtgcctggag   900
cacctgttct tcttcaagct catcggggac acgcccacg acaccttct catggagatg   960
ctggaggcac cacatcaagc cacctag   987

```

&lt;210&gt; 22

&lt;211&gt; 789

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;400&gt; 22

```

aagcgggaag ctgtgcagga ggagcggcag cggggcaagg accggaatga gaacgaggtg   60
gagtccacca gcagtccaa cgaggacatg cctgtagaga agattctgga agccgagctt   120
gctgtcgagc ccaagactga gacatactg gaggcaaaaca tggggctgaa cccagctca   180
ccaaatgacc ctgttaccaa catctgtcaa gcagcagaca agcagctctt cactcttgtg   240
gagtgggcca agaggatccc acactttct gagctgccc tagacgacca ggtcatcctg   300
ctacgggcag gctggaacga gctgctgac gccctcttct cccaccgctc catagctgtg   360
aaagatggga ttctcctggc caccggcctg cactacacc ggaacagcgc tcacagtgtc   420
gggggtggcg ccattcttga cagggtgcta acagagctgg tgtetaagat gcgtgacatg   480
cagatggaca agacggagct gggctgcctg cgagccattg tcctgttcaa cctgactct   540
aaggggtct caaacctgc tgaggtggag gcgttgagg agaaggtgta tgcgtacta   600
gaagcgtact gcaaacacaa gtaccctgag cagccgggca ggtttgcaa gctgctgctc   660

```

WO 01/70816

PCT/US01/09050

RH0020.ST25

cgctgcctg cactgcgttc catcgggctc aagtgcctgg agcacctgtt cttcttcaag 720  
ctcatcgggg acacgcccac cgacaccttc ctcatggaga tgctggaggc accacatcaa 780  
gccacctag 789

<210> 23  
<211> 714  
<212> DNA  
<213> Artificial Sequence

<400> 23  
gccaacgagg acatgcctgt agagaagatt ctggaagccg agcttgctgt cgagcccaag 60  
actgagacat acgtggaggc aaacatgggg ctgaacccca gctcaccaaa tgaccctgtt 120  
accaacatct gtcaagcagc agacaagcag ctcttcactc ttgtggagtg ggccaagagg 180  
atcccacact tttctgagct gcccttagac gaccaggcca tcctgctacg ggcaggctgg 240  
aacgagctgc tgategcctc cttctccac cgctccatag ctgtgaaaga tgggattctc 300  
ctggccaccg gctgcacgt acaccggaac agcgctcaca gtgctggggg gggcgccatc 360  
tttgacaggg tgctaacaga gctgggtgtc aagatgcgtg acatgcagat ggacaagacg 420  
gagctgggct gcctgcgagc cattgtcctg ttcaaccctg actctaagg gctctcaaac 480  
cctgctgagg tggaggcgtt gaggggagaag gtgtatgctg cactagaagc gtactgcaaa 540  
cacaagtacc ctgagcagcc gggcagggtt gccaaagctg tgctccgcct gcctgcactg 600  
cgttccatcg ggtcctaagt cctggagcac ctgttcttct tcaagctcat cggggacacg 660  
cccatcgaca cttctctcat ggagatgctg gaggcaccac atcaagccac ctag 714

<210> 24  
<211> 536  
<212> DNA  
<213> Artificial Sequence

<400> 24  
ggatcccaca cttttctgag ctgcccctag acgaccaggt catcctgcta cgggcaggct 60  
ggaacgagct gctgatcgcc tccttctccc accgctccat agctgtgaaa gatgggattc 120  
tcctggccac cggcctgcac gtacaccgga acagcgtcga cagtgtctgg gtggcgcca 180  
tctttgacag ggtgctaaca gagctggtgt ctaagatgag tgacatgcag atggacaaga 240  
cggagctggg ctgcctgcga gccattgtcc tgttcaaccc tgactctaag gggctctcaa 300  
accctgctga ggtggaggcg ttgagggaga aggtgtatgc gtcactagaa gcgtactgca 360  
aacacaagta ccttgagcag cggggcagggt ttgccaagct gctgctccgc ctgcctgcac 420  
tgcgttccat cgggctcaag tgcttgagc acctgttctt ctcaagctc atcggggaca 480  
cgcccatcga caccttctc atggagatgc tggaggcacc acatcaagcc acctag 536

<210> 25



RH0020.ST25

&lt;211&gt; 672

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;400&gt; 25

```

gccaacgagg acatgcctgt agagaagatt ctggaagccg agcttgctgt cgagcccaag    60
actgagacat acgtggaggg aaacatgggg ctgaacccca gctcaccaaa tgaccctgtt    120
accaacatct gtcaagcagc agacaagcag ctcttcactc ttgtggagtg ggccaagagg    180
atccacactc tttctgagct gccctagac gaccagggtca tcctgctacg ggcaggctgg    240
aacgagctgc tgatcgctc cttctccac cgctccatag ctgtgaaaga tgggattctc    300
ctggccaccg gctgcacgt acaccgaac agcgctcaca gtgctggggg gggcgccatc    360
tttgacaggg tgctaacaga gctggtgtct aagatgcgtg acatgcagat ggacaagacg    420
gagctgggct gctgcgagc cattgtctc ttcaaccctg actctaagg gctctcaaac    480
cctgctgagg tggaggcggt gagggagaag gtgtatgct cactagaagc gtactgcaaa    540
cacaagtacc ctgagcagcc gggcagggtt gccaaagctg tgctccgcct gcctgactg    600
cgttccatcg ggctcaagtg cctggagcac ctgttcttct tcaagctcat cggggacacg    660
cccatcgaca cc                                     672

```

&lt;210&gt; 26

&lt;211&gt; 1123

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;400&gt; 26

```

tgcgccatct gcggggaccg ctcctcaggg aagcactatg gagtgtacag ctgcgagggg    60
tgcaagggct tcttcaagcg gacggtgctg aaggacctga cctacacctg ccgcgacaac    120
aaggactgcc tgattgacaa gcggcagcgg aaccggtgcc agtactgccg ctaccagaag    180
tgcttgccca tgggcatgaa gcgggaagcc gtgcaggagg agcggcagcg tggcaaggac    240
cggaacgaga atgaggtgga gtcgaccagc agcgccaacg aggacatgcc ggtggagagg    300
atcctggagg ctgagctggc cgtggagccc aagaccgaga cctacgtgga ggcaaacatg    360
gggtgaacc ccagctcgcc gaacgacctc gtcaccaaca ttgtccaagc agccgacaaa    420
cagcttttca ccctggtgga gtgggccaag cggatccac acttctcaga gctgccctg    480
gacgaccagg tcactctgct gcgggcaggc tggaatgagc tgctcatcgc ctcttctcc    540
caccgctcca tcgccgtgaa ggacgggatc ctcctggcca ccgggctgca cgtccaccgg    600
aacagcgccc acagcgagg ggtggcgccc atctttgaca ggtgctgac ggagcttggt    660
tccaagatgc gggacatgca gatggacaag acggagctgg gctgcctgcg cgccatcgtc    720
ctctttaacc ctgactccaa ggggtctctg aaccggccg aggtggaggc gctgaggagg    780
aaggctctat cgtccttgga ggcctactgc aagcacaagt acccagagca gccgggaagg    840
ttcgctaagc tcttgctcgg cctgccggct ctgcgctcca tcgggctcaa atgcctggaa    900

```

## RH0020.ST25

catctcttct tcttcaagct catcggggac acaccattg acaccttct tatggagatg 960  
 ctggaggcgc cgcaccaa at gacttaggcc tgcgggcca tctttgtgc ccaccgttc 1020  
 tggccaccct gcctggagc cagctgttct tctcagcctg agccctgtcc ctgcccttct 1080  
 ctgcctggcc tgtttggact ttggggcaca gcctgtcact gct 1123

<210> 27  
 <211> 925  
 <212> DNA  
 <213> Artificial Sequence

<400> 27  
 aagcgggaag ccgtgcagga ggagcggcag cgtggcaagg accggaacga gaatgaggatg 60  
 gagtgcacca gcagcgccaa cgaggacatg ccggtggaga ggatcctgga ggctgagctg 120  
 gccgtggagc ccaagaccga gacctacgtg gaggcaaaca tggggctgaa cccagctcg 180  
 ccgaacgacc ctgtcaccaa catttgccaa gcagccgaca aacagctttt caccctgggtg 240  
 gagtgggcca agcggatccc acatttctca gagctgcccc tggacgacca ggtcatcctg 300  
 ctgcgggcag gctggaatga gctgctcatc gcctccttct cccacgcctc catcgccgtg 360  
 aaggacggga tctctctggc caccgggtg cactgccacc ggaacagcgc ccacagcgca 420  
 ggggtgggcg ccattcttga caggggtgtg acggagcttg tgtccaagat gcgggacatg 480  
 cagatggaca agacggagct gggctgcctg cgcgccatcg tctcttttaa ccctgaactcc 540  
 aaggggtctt cgaacccggc cgaggtggag gcgctgaggg agaaggtcta tgcgtccttg 600  
 gaggcctact gcaagcacia gtaccagag cagccgggaa ggttcgctaa gctcttgctc 660  
 cgctgcccgc ctctgcgtc catcgggctc aaatgcctgg aacatctctt cttcttcaag 720  
 ctcatcgggg acacacccat tgacacctc cttatggaga tgcctggagg gccgcaccaa 780  
 atgacttagg cctgcgggcc catcctttgt gcccacccgt tctggccacc ctgcctggac 840  
 gccagctgtt cttctcagcc tgagccctgt ccctgccctt ctctgcctg cctgtttgga 900  
 ctttggggca cagcctgtca ctgct 925

<210> 28  
 <211> 850  
 <212> DNA  
 <213> Artificial Sequence

<400> 28  
 gccaacgagg acatgccggt ggagaggatc ctggaggctg agctggcctg ggagcccaag 60  
 accgagacct acgtggaggc aaacatgggg ctgaacccca gctcgccgaa cgaccctgtc 120  
 accaacattt gccaaagcag cgacaacag cttttcacc tgggtgagtg ggccaagcgg 180  
 atccccact tctcagagct gccctggac gaccaggtca tctgctgctg ggcaggctgg 240  
 aatgagctgc tcatgcctc cttctccac cgctccatcg ccgtgaagga cgggacctc 300

RH0020.ST25

ctggccaccg ggctgcacgt ccaccggaac agcgcccaca gcgcaggggt gggcgccatc 360  
 ttgacaggg tgctgacgga gcttgtgtcc aagatgcggg acatgcagat ggacaagacg 420  
 gagctgggct gcctgcgcgc catcgctcctc ttttaaccctg actccaaggg gctctcgaac 480  
 ccggccgagg tggaggcgct gagggagaag gtctatgcgt ccttgagggc ctactgcaag 540  
 cacaagtacc cagagcagcc ggaaggttc gctaagctct tgcctcgccct gccggtctg 600  
 cgctccatcg ggctcaaagt cctggaacat ctcttcttct tcaagctcat cggggacaca 660  
 cccattgaca ccttccttat ggagatgctg gaggcgcgc accaaatgac ttaggcctgc 720  
 gggcccatcc tttgtgcca cccgttctg ccaccctgcc tggacgccag ctgttcttct 780  
 cagcctgagc cctgtccctg cccttctctg cctggcctgt ttggactttg gggcacagcc 840  
 tgtcactgct 850

<210> 29  
 <211> 670  
 <212> DNA  
 <213> Artificial Sequence

<400> 29  
 atccacact tctcagagct gccctggac gaccaggtca tcctgctgcg ggcaggctgg 60  
 aatgagctgc tcatgcctc ctctccac cgctccatcg ccgtgaagga cgggacctc 120  
 ctggccaccg ggctgcacgt ccaccggaac agcgcccaca gcgcaggggt gggcgccatc 180  
 ttgacaggg tgctgacgga gcttgtgtcc aagatgcggg acatgcagat ggacaagacg 240  
 gagctgggct gcctgcgcgc catcgctcctc ttttaaccctg actccaaggg gctctcgaac 300  
 ccggccgagg tggaggcgct gagggagaag gtctatgcgt ccttgagggc ctactgcaag 360  
 cacaagtacc cagagcagcc ggaaggttc gctaagctct tgcctcgccct gccggtctg 420  
 cgctccatcg ggctcaaagt cctggaacat ctcttcttct tcaagctcat cggggacaca 480  
 cccattgaca ccttccttat ggagatgctg gaggcgcgc accaaatgac ttaggcctgc 540  
 gggcccatcc tttgtgcca cccgttctg ccaccctgcc tggacgccag ctgttcttct 600  
 cagcctgagc cctgtccctg cccttctctg cctggcctgt ttggactttg gggcacagcc 660  
 tgtcactgct 670

<210> 30  
 <211> 672  
 <212> DNA  
 <213> Artificial Sequence

<400> 30  
 gccacgagg acatgccgtt ggagaggtc ctggaggctg agctggcctt ggagcccaag 60  
 accgagacct acgtggaggc aaacatgggg ctgaaccca gctcgccgaa cgaccctgtc 120  
 accaacattt gccaaagcgc cgacaaacag cttttcacc ttgtggagtg ggccaagcgg 180  
 atccacact tctcagagct gccctggac gaccaggtca tcctgctgcg ggcaggctgg 240

WO 01/70816

PCT/US01/09050

RH0020.ST25

```

aatgagctgc tcatcgccctc cttctccac cgctccatcg ccgtgaagga cgggatcctc 300
ctggccaccg ggctgcacgt ccaccggaac agcgcccaca gcgcaggggt gggcgccatc 360
tttgacaggg tgctgacgga gcttgtgtcc aagatgcggg acatgcagat ggacaagacg 420
gagctgggct gcctgcgcgc catcgctctc tttaaccctg actccaaggg gctctcgaac 480
ccggccgagg tggaggcgct gagggagaag gtctatgcgt ccttgagggc ctactgcaag 540
cacaagtacc cagagcagcc gggaagggtc gctaagctct tgctccgcct gccggctctg 600
cgctccatcg ggctcaaatg cctggaacat ctctctctct tcaagctcat cggggacaca 660
ccattgaca cc 672

```

<210> 31  
 <211> 328  
 <212> PRT  
 <213> Artificial Sequence

&lt;400&gt; 31

```

Cys Ala Ile Cys Gly Asp Arg Ser Ser Gly Lys His Tyr Gly Val Tyr
1      5      10      15
Ser Cys Glu Gly Cys Lys Gly Phe Phe Lys Arg Thr Val Arg Lys Asp
20     25     30
Leu Thr Tyr Thr Cys Arg Asp Asn Lys Asp Cys Leu Ile Asp Lys Arg
35     40     45
Gln Arg Asn Arg Cys Gln Tyr Cys Arg Tyr Gln Lys Cys Leu Ala Met
50     55     60
Gly Met Lys Arg Glu Ala Val Gln Glu Glu Arg Gln Arg Gly Lys Asp
65     70     75     80
Arg Asn Glu Asn Glu Val Glu Ser Thr Ser Ser Ala Asn Glu Asp Met
85     90     95
Pro Val Glu Lys Ile Leu Glu Ala Glu Leu Ala Val Glu Pro Lys Thr
100    105    110
Glu Thr Tyr Val Glu Ala Asn Met Gly Leu Asn Pro Ser Ser Pro Asn
115    120    125
Asp Pro Val Thr Asn Ile Cys Gln Ala Ala Asp Lys Gln Leu Phe Thr
130    135    140
Leu Val Glu Trp Ala Lys Arg Ile Pro His Phe Ser Glu Leu Pro Leu
145    150    155    160
Asp Asp Gln Val Ile Leu Leu Arg Ala Gly Trp Asn Glu Leu Leu Ile
165    170    175
Ala Ser Phe Ser His Arg Ser Ile Ala Val Lys Asp Gly Ile Leu Leu
180    185    190
Ala Thr Gly Leu His Val His Arg Asn Ser Ala His Ser Ala Gly Val
195    200    205
Gly Ala Ile Phe Asp Arg Val Leu Thr Glu Leu Val Ser Lys Met Arg

```

WO 01/70816

PCT/US01/09050

RH0020.ST25  
220

210 215  
Asp Met Gln Met Asp Lys Thr Glu Leu Gly Cys Leu Arg Ala Ile Val  
225 230 235 240  
Leu Phe Asn Pro Asp Ser Lys Gly Leu Ser Asn Pro Ala Glu Val Glu  
245 250 255  
Ala Leu Arg Glu Lys Val Tyr Ala Ser Leu Glu Ala Tyr Cys Lys His  
260 265 270  
Lys Tyr Pro Glu Gln Pro Gly Arg Phe Ala Lys Leu Leu Leu Arg Leu  
275 280 285  
Pro Ala Leu Arg Ser Ile Gly Leu Lys Cys Leu Glu His Leu Phe Phe  
290 295 300  
Phe Lys Leu Ile Gly Asp Thr Pro Ile Asp Thr Phe Leu Met Glu Met  
305 310 315 320  
Leu Glu Ala Pro His Gln Ala Thr  
325

<210> 32  
<211> 262  
<212> PRT  
<213> Artificial Sequence

<400> 32

Lys Arg Glu Ala Val Gln Glu Glu Arg Gln Arg Gly Lys Asp Arg Asn  
1 5 10 15  
Glu Asn Glu Val Glu Ser Thr Ser Ser Ala Asn Glu Asp Met Pro Val  
20 25 30  
Glu Lys Ile Leu Glu Ala Glu Leu Ala Val Glu Pro Lys Thr Glu Thr  
35 40 45  
Tyr Val Glu Ala Asn Met Gly Leu Asn Pro Ser Ser Pro Asn Asp Pro  
50 55 60  
Val Thr Asn Ile Cys Gln Ala Ala Asp Lys Gln Leu Phe Thr Leu Val  
65 70 75 80  
Glu Trp Ala Lys Arg Ile Pro His Phe Ser Glu Leu Pro Leu Asp Asp  
85 90 95  
Gln Val Ile Leu Leu Arg Ala Gly Trp Asn Glu Leu Leu Ile Ala Ser  
100 105 110  
Phe Ser His Arg Ser Ile Ala Val Lys Asp Gly Ile Leu Leu Ala Thr  
115 120 125  
Gly Leu His Val His Arg Asn Ser Ala His Ser Ala Gly Val Gly Ala  
130 135 140  
Ile Phe Asp Arg Val Leu Thr Glu Leu Val Ser Lys Met Arg Asp Met  
145 150 155 160  
Gln Met Asp Lys Thr Glu Leu Gly Cys Leu Arg Ala Ile Val Leu Phe  
165 170 175  
Asn Pro Asp Ser Lys Gly Leu Ser Asn Pro Ala Glu Val Glu Ala Leu  
180 185 190

WO 01/70816

PCT/US01/09050

RH0020.ST25

Arg Glu Lys Val Tyr Ala Ser Leu Glu Ala Tyr Cys Lys His Lys Tyr  
195 200 205  
Pro Glu Gln Pro Gly Arg Phe Ala Lys Leu Leu Leu Arg Leu Pro Ala  
210 215 220  
Leu Arg Ser Ile Gly Leu Lys Cys Leu Glu His Leu Phe Phe Phe Lys  
225 230 235 240  
Leu Ile Gly Asp Thr Pro Ile Asp Thr Phe Leu Met Glu Met Leu Glu  
245 250 255

Ala Pro His Gln Ala Thr  
260

<210> 33  
<211> 237  
<212> PRT  
<213> Artificial Sequence

<400> 33

Ala Asn Glu Asp Met Pro Val Glu Lys Ile Leu Glu Ala Glu Leu Ala  
1 5 10 15  
Val Glu Pro Lys Thr Glu Thr Tyr Val Glu Ala Asn Met Gly Leu Asn  
20 25 30  
Pro Ser Ser Pro Asn Asp Pro Val Thr Asn Ile Cys Gln Ala Ala Asp  
35 40 45  
Lys Gln Leu Phe Thr Leu Val Glu Trp Ala Lys Arg Ile Pro His Phe  
50 55 60  
Ser Glu Leu Pro Leu Asp Asp Gln Val Ile Leu Leu Arg Ala Gly Trp  
65 70 75 80  
Asn Glu Leu Leu Ile Ala Ser Phe Ser His Arg Ser Ile Ala Val Lys  
85 90 95  
Asp Gly Ile Leu Leu Ala Thr Gly Leu His Val His Arg Asn Ser Ala  
100 105 110  
His Ser Ala Gly Val Gly Ala Ile Phe Asp Arg Val Leu Thr Glu Leu  
115 120 125  
Val Ser Lys Met Arg Asp Met Gln Met Asp Lys Thr Glu Leu Gly Cys  
130 135 140  
Leu Arg Ala Ile Val Leu Phe Asn Pro Asp Ser Lys Gly Leu Ser Asn  
145 150 155 160  
Pro Ala Glu Val Glu Ala Leu Arg Glu Lys Val Tyr Ala Ser Leu Glu  
165 170 175  
Ala Tyr Cys Lys His Lys Tyr Pro Glu Gln Pro Gly Arg Phe Ala Lys  
180 185 190  
Leu Leu Leu Arg Leu Pro Ala Leu Arg Ser Ile Gly Leu Lys Cys Leu  
195 200 205  
Glu His Leu Phe Phe Phe Lys Leu Ile Gly Asp Thr Pro Ile Asp Thr  
210 215 220

WO 01/70816

PCT/US01/09050

RH0020.ST25

Phe Leu Met Glu Met Leu Glu Ala Pro His Gln Ala Thr  
 225 230 235

<210> 34  
 <211> 177  
 <212> PRT  
 <213> Artificial Sequence

<400> 34

Ile Pro His Phe Ser Glu Leu Pro Leu Asp Asp Gln Val Ile Leu Leu  
 1 5 10 15

Arg Ala Gly Trp Asn Glu Leu Leu Ile Ala Ser Phe Ser His Arg Ser  
 20 25 30

Ile Ala Val Lys Asp Gly Ile Leu Leu Ala Thr Gly Leu His Val His  
 35 40 45

Arg Asn Ser Ala His Ser Ala Gly Val Gly Ala Ile Phe Asp Arg Val  
 50 55 60

Leu Thr Glu Leu Val Ser Lys Met Arg Asp Met Gln Met Asp Lys Thr  
 65 70 75 80

Glu Leu Gly Cys Leu Arg Ala Ile Val Leu Phe Asn Pro Asp Ser Lys  
 85 90 95

Gly Leu Ser Asn Pro Ala Glu Val Glu Ala Leu Arg Glu Lys Val Tyr  
 100 105 110

Ala Ser Leu Glu Ala Tyr Cys Lys His Lys Tyr Pro Glu Gln Pro Gly  
 115 120 125

Arg Phe Ala Lys Leu Leu Leu Arg Leu Pro Ala Leu Arg Ser Ile Gly  
 130 135 140

Leu Lys Cys Leu Glu His Leu Phe Phe Phe Lys Leu Ile Gly Asp Thr  
 145 150 155 160

Pro Ile Asp Thr Phe Leu Met Glu Met Leu Glu Ala Pro His Gln Ala  
 165 170 175

Thr

<210> 35  
 <211> 224  
 <212> PRT  
 <213> Artificial Sequence

<400> 35

Ala Asn Glu Asp Met Pro Val Glu Lys Ile Leu Glu Ala Glu Leu Ala  
 1 5 10 15

Val Glu Pro Lys Thr Glu Thr Tyr Val Glu Ala Asn Met Gly Leu Asn  
 20 25 30

Pro Ser Ser Pro Asn Asp Pro Val Thr Asn Ile Cys Gln Ala Ala Asp  
 35 40 45

Lys Gln Leu Phe Thr Leu Val Glu Trp Ala Lys Arg Ile Pro His Phe  
 50 55 60

WO 01/70816

PCT/US01/09050

RH0020.ST25

Ser Glu Leu Pro Leu Asp Asp Gln Val Ile Leu Leu Arg Ala Gly Trp  
65 70 75 80

Asn Glu Leu Leu Ile Ala Ser Phe Ser His Arg Ser Ile Ala Val Lys  
85 90 95

Asp Gly Ile Leu Leu Ala Thr Gly Leu His Val His Arg Asn Ser Ala  
100 105 110

His Ser Ala Gly Val Gly Ala Ile Phe Asp Arg Val Leu Thr Glu Leu  
115 120 125

Val Ser Lys Met Arg Asp Met Gln Met Asp Lys Thr Glu Leu Gly Cys  
130 135 140

Leu Arg Ala Ile Val Leu Phe Asn Pro Asp Ser Lys Gly Leu Ser Asn  
145 150 155 160

Pro Ala Glu Val Glu Ala Leu Arg Glu Lys Val Tyr Ala Ser Leu Glu  
165 170 175

Ala Tyr Cys Lys His Lys Tyr Pro Glu Gln Pro Gly Arg Phe Ala Lys  
180 185 190

Leu Leu Leu Arg Leu Pro Ala Leu Arg Ser Ile Gly Leu Lys Cys Leu  
195 200 205

Glu His Leu Phe Phe Phe Lys Leu Ile Gly Asp Thr Pro Ile Asp Thr  
210 215 220

<210> 36

<211> 328

<212> PRT

<213> Artificial Sequence

<220>

<221> misc feature

<223> Novel Sequence

<400> 36

Cys Ala Ile Cys Gly Asp Arg Ser Ser Gly Lys His Tyr Gly Val Tyr  
1 5 10 15

Ser Cys Glu Gly Cys Lys Gly Phe Phe Lys Arg Thr Val Arg Lys Asp  
20 25 30

Leu Thr Tyr Thr Cys Arg Asp Asn Lys Asp Cys Leu Ile Asp Lys Arg  
35 40 45

Gln Arg Asn Arg Cys Gln Tyr Cys Arg Tyr Gln Lys Cys Leu Ala Met  
50 55 60

Gly Met Lys Arg Glu Ala Val Gln Glu Glu Arg Gln Arg Gly Lys Asp  
65 70 75 80

Arg Asn Glu Asn Glu Val Glu Ser Thr Ser Ser Ala Asn Glu Asp Met  
85 90 95

Pro Val Glu Arg Ile Leu Glu Ala Glu Leu Ala Val Glu Pro Lys Thr  
100 105 110

Glu Thr Tyr Val Glu Ala Asn Met Gly Leu Asn Pro Ser Ser Pro Asn  
115 120 125



WO 01/70816

PCT/US01/09050

RH0020.ST25

Asp Pro Val Thr Asn Ile Cys Gln Ala Ala Asp Lys Gln Leu Phe Thr  
 130 135 140  
 Leu Val Glu Trp Ala Lys Arg Ile Pro His Phe Ser Glu Leu Pro Leu  
 145 150 155 160  
 Asp Asp Gln Val Ile Leu Leu Arg Ala Gly Trp Asn Glu Leu Leu Ile  
 165 170 175  
 Ala Ser Phe Ser His Arg Ser Ile Ala Val Lys Asp Gly Ile Leu Leu  
 180 185 190  
 Ala Thr Gly Leu His Val His Arg Asn Ser Ala His Ser Ala Gly Val  
 195 200 205  
 Gly Ala Ile Phe Asp Arg Val Leu Thr Glu Leu Val Ser Lys Met Arg  
 210 215 220  
 Asp Met Gln Met Asp Lys Thr Glu Leu Gly Cys Leu Arg Ala Ile Val  
 225 230 235 240  
 Leu Phe Asn Pro Asp Ser Lys Gly Leu Ser Asn Pro Ala Glu Val Glu  
 245 250 255  
 Ala Leu Arg Glu Lys Val Tyr Ala Ser Leu Glu Ala Tyr Cys Lys His  
 260 265 270  
 Lys Tyr Pro Glu Gln Pro Gly Arg Phe Ala Lys Leu Leu Leu Arg Leu  
 275 280 285  
 Pro Ala Leu Arg Ser Ile Gly Leu Lys Cys Leu Glu His Leu Phe Phe  
 290 295 300  
 Phe Lys Leu Ile Gly Asp Thr Pro Ile Asp Thr Phe Leu Met Glu Met  
 305 310 315 320  
 Leu Glu Ala Pro His Gln Met Thr  
 325

<210> 37  
 <211> 262  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <221> misc\_feature  
 <223> Novel Sequence

<400> 37

Lys Arg Glu Ala Val Gln Glu Glu Arg Gln Arg Gly Lys Asp Arg Asn  
 1 5 10 15  
 Glu Asn Glu Val Glu Ser Thr Ser Ser Ala Asn Glu Asp Met Pro Val  
 20 25 30

Glu Arg Ile Leu Glu Ala Glu Leu Ala Val Glu Pro Lys Thr Glu Thr  
 35 40 45

Tyr Val Glu Ala Asn Met Gly Leu Asn Pro Ser Ser Pro Asn Asp Pro  
 50 55 60

Val Thr Asn Ile Cys Gln Ala Ala Asp Lys Gln Leu Phe Thr Leu Val

WO 01/70816

PCT/US01/09050

RH0020.ST25

65	70	75	80
Glu Trp Ala Lys Arg Ile Pro His Phe Ser Glu Leu Pro Leu Asp Asp	85	90	95
Gln Val Ile Leu Leu Arg Ala Gly Trp Asn Glu Leu Leu Ile Ala Ser	100	105	110
Phe Ser His Arg Ser Ile Ala Val Lys Asp Gly Ile Leu Leu Ala Thr	115	120	125
Gly Leu His Val His Arg Asn Ser Ala His Ser Ala Gly Val Gly Ala	130	135	140
Ile Phe Asp Arg Val Leu Thr Glu Leu Val Ser Lys Met Arg Asp Met	145	150	155
Gln Met Asp Lys Thr Glu Leu Gly Cys Leu Arg Ala Ile Val Leu Phe	165	170	175
Asn Pro Asp Ser Lys Gly Leu Ser Asn Pro Ala Glu Val Glu Ala Leu	180	185	190
Arg Glu Lys Val Tyr Ala Ser Leu Glu Ala Tyr Cys Lys His Lys Tyr	195	200	205
Pro Glu Gln Pro Gly Arg Phe Ala Lys Leu Leu Leu Arg Leu Pro Ala	210	215	220
Leu Arg Ser Ile Gly Leu Lys Cys Leu Glu His Leu Phe Phe Phe Lys	225	230	235
Leu Ile Gly Asp Thr Pro Ile Asp Thr Phe Leu Met Glu Met Leu Glu	245	250	255
Ala Pro His Gln Met Thr	260		

<210> 38  
 <211> 237  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <221> misc\_feature  
 <223> Novel Sequence

<400> 38

Ala Asn Glu Asp Met Pro Val Glu Arg Ile Leu Glu Ala Glu Leu Ala	1	5	10	15
Val Glu Pro Lys Thr Glu Thr Tyr Val Glu Ala Asn Met Gly Leu Asn	20	25	30	
Pro Ser Ser Pro Asn Asp Pro Val Thr Asn Ile Cys Gln Ala Ala Asp	35	40	45	
Lys Gln Leu Phe Thr Leu Val Glu Trp Ala Lys Arg Ile Pro His Phe	50	55	60	
Ser Glu Leu Pro Leu Asp Asp Gln Val Ile Leu Leu Arg Ala Gly Trp	65	70	75	80

WO 01/70816

PCT/US01/09050

RH0020.ST25

Asn Glu Leu Leu Ile Ala Ser Phe Ser His Arg Ser Ile Ala Val Lys  
85 90 95  
Asp Gly Ile Leu Leu Ala Thr Gly Leu His Val His Arg Asn Ser Ala  
100 105 110  
His Ser Ala Gly Val Gly Ala Ile Phe Asp Arg Val Leu Thr Glu Leu  
115 120 125  
Val Ser Lys Met Arg Asp Met Gln Met Asp Lys Thr Glu Leu Gly Cys  
130 135 140  
Leu Arg Ala Ile Val Leu Phe Asn Pro Asp Ser Lys Gly Leu Ser Asn  
145 150 155 160  
Pro Ala Glu Val Glu Ala Leu Arg Glu Lys Val Tyr Ala Ser Leu Glu  
165 170 175  
Ala Tyr Cys Lys His Lys Tyr Pro Glu Gln Pro Gly Arg Phe Ala Lys  
180 185 190  
Leu Leu Leu Arg Leu Pro Ala Leu Arg Ser Ile Gly Leu Lys Cys Leu  
195 200 205  
Glu His Leu Phe Phe Phe Lys Leu Ile Gly Asp Thr Pro Ile Asp Thr  
210 215 220  
Phe Leu Met Glu Met Leu Glu Ala Pro His Gln Met Thr  
225 230 235

<210> 39  
<211> 177  
<212> PRT  
<213> Artificial Sequence

<220>  
<221> misc feature  
<223> Novel Sequence

<400> 39

Ile Pro His Phe Ser Glu Leu Pro Leu Asp Asp Gln Val Ile Leu Leu  
1 5 10 15  
Arg Ala Gly Trp Asn Glu Leu Leu Ile Ala Ser Phe Ser His Arg Ser  
20 25 30  
Ile Ala Val Lys Asp Gly Ile Leu Leu Ala Thr Gly Leu His Val His  
35 40 45  
Arg Asn Ser Ala His Ser Ala Gly Val Gly Ala Ile Phe Asp Arg Val  
50 55 60  
Leu Thr Glu Leu Val Ser Lys Met Arg Asp Met Gln Met Asp Lys Thr  
65 70 75 80  
Glu Leu Gly Cys Leu Arg Ala Ile Val Leu Phe Asn Pro Asp Ser Lys  
85 90 95  
Gly Leu Ser Asn Pro Ala Glu Val Glu Ala Leu Arg Glu Lys Val Tyr  
100 105 110  
Ala Ser Leu Glu Ala Tyr Cys Lys His Lys Tyr Pro Glu Gln Pro Gly  
115 120 125

WO 01/70816

PCT/US01/09050

RH0020.ST25

Arg Phe Ala Lys Leu Leu Leu Arg Leu Pro Ala Leu Arg Ser Ile Gly  
130 135 140  
Leu Lys Cys Leu Glu His Leu Phe Phe Phe Lys Leu Ile Gly Asp Thr  
145 150 155 160  
Pro Ile Asp Thr Phe Leu Met Glu Met Leu Glu Ala Pro His Gln Met  
165 170 175

Thr

<210> 40  
<211> 224  
<212> PRT  
<213> Artificial Sequence

<220>  
<221> misc feature  
<223> Novel Sequence

<400> 40

Ala Asn Glu Asp Met Pro Val Glu Arg Ile Leu Glu Ala Glu Leu Ala  
1 5 10 15  
Val Glu Pro Lys Thr Glu Thr Tyr Val Glu Ala Asn Met Gly Leu Asn  
20 25 30  
Pro Ser Ser Pro Asn Asp Pro Val Thr Asn Ile Cys Gln Ala Ala Asp  
35 40 45  
Lys Gln Leu Phe Thr Leu Val Glu Trp Ala Lys Arg Ile Pro His Phe  
50 55 60  
Ser Glu Leu Pro Leu Asp Asp Gln Val Ile Leu Leu Arg Ala Gly Trp  
65 70 75 80  
Asn Glu Leu Leu Ile Ala Ser Phe Ser His Arg Ser Ile Ala Val Lys  
85 90 95  
Asp Gly Ile Leu Leu Ala Thr Gly Leu His Val His Arg Asn Ser Ala  
100 105 110  
His Ser Ala Gly Val Gly Ala Ile Phe Asp Arg Val Leu Thr Glu Leu  
115 120 125  
Val Ser Lys Met Arg Asp Met Gln Met Asp Lys Thr Glu Leu Gly Cys  
130 135 140  
Leu Arg Ala Ile Val Leu Phe Asn Pro Asp Ser Lys Gly Leu Ser Asn  
145 150 155 160  
Pro Ala Glu Val Glu Ala Leu Arg Glu Lys Val Tyr Ala Ser Leu Glu  
165 170 175  
~~Ala-Tyr-Cys-Lys-His-Lys-Tyr-Pro-Glu-Gln-Pro-Gly-Arg-Phe-Ala-Lys~~  
180 185 190  
Leu Leu Leu Arg Leu Pro Ala Leu Arg Ser Ile Gly Leu Lys Cys Leu  
195 200 205  
Glu His Leu Phe Phe Phe Lys Leu Ile Gly Asp Thr Pro Ile Asp Thr

WO 01/70816

PCT/US01/09050

RH0020.ST25  
220

210

215

<210> 41  
<211> 441  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> misc\_feature  
<223> Novel Sequence

<400> 41  
atgaagctac tgtcttctat cgaacaagca tgcgatattt gccgacttaa aaagctcaag 60  
tgctccaaag aaaaaccgaa gtgcgccaag tgtctgaaga acaactggga gtgtcgctac 120  
tctcccaaaa caaaagggtc tccgctgact agggcacatc tgacagaagt ggaatcaagg 180  
ctagaagac tggaacagct atttctactg atttttcctc gagaagacct tgacatgatt 240  
ttgaaaatgg attctttaca ggatataaaa gcattgttaa caggattatt tgtacaagat 300  
aatgtgaata aagatgccgt cacagataga ttggcttcag tggagactga tatgcctcta 360  
acattgagac agcatagaat aagtgcgaca tcatcatcgg aagagagtag taacaaaggt 420  
caaagacagt tgactgtatc g 441

<210> 42  
<211> 147  
<212> PRT  
<213> Artificial Sequence

<220>  
<221> misc\_feature  
<223> Novel Sequence

<400> 42

Met Lys Leu Leu Ser Ser Ile Glu Gln Ala Cys Asp Ile Cys Arg Leu  
1 5 10 15  
Lys Lys Leu Lys Cys Ser Lys Glu Lys Pro Lys Cys Ala Lys Cys Leu  
20 25 30  
Lys Asn Asn Trp Glu Cys Arg Tyr Ser Pro Lys Thr Lys Arg Ser Pro  
35 40 45  
Leu Thr Arg Ala His Leu Thr Glu Val Glu Ser Arg Leu Glu Arg Leu  
50 55 60  
Glu Gln Leu Phe Leu Leu Ile Phe Pro Arg Glu Asp Leu Asp Met Ile  
65 70 75 80  
Leu Lys Met Asp Ser Leu Gln Asp Ile Lys Ala Leu Leu Thr Gly Leu  
85 90 95  
Phe Val Gln Asp Asn Val Asn Lys Asp Ala Val Thr Asp Arg Leu Ala  
100 105 110  
Ser Val Glu Thr Asp Met Pro Leu Thr Leu Arg Gln His Arg Ile Ser  
115 120 125

WO 01/70816

PCT/US01/09050

RH0020.ST25

Ala Thr Ser Ser Ser Glu Glu Ser Ser Asn Lys Gly Gln Arg Gln Leu  
130 135 140

Thr Val Ser  
145

<210> 43  
<211> 606  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> misc\_feature  
<223> Novel Sequence

<400> 43  
atgaaagcgt taacggccag gcaacaagag gtgtttgatc tcattccgtga tcacatcagc 60  
cagacaggta tgccgccgac gcgtgcggaa atcgcgagc gtttggggtt ccgttcccca 120  
aacgcggctg aagaacatct gaaggcgtg gcacgcaaag gcgttattga aattgtttcc 180  
ggcgcatcac ggggattcgt tctgttcag gaagaggaag aagggttgcc gctgtaggt 240  
cgtgtggctg ccgtgaacc acttctggcg caacagcata ttgaaggta ttatcaggtc 300  
gatccttctt tattcaagcc gaatgctgat ttcctgctgc gcgtcagcgg gatgctgatg 360  
aaagatatcg gcattatgga tgggtgacttg ctggcagtg ataaaaactca ggatgtacgt 420  
aacggtcagg tcgtgtgcgc acgtattgat gacgaagta ccgttaagcg cctgaaaaaa 480  
cagggcaata aagtcgaact gttgccagaa aatagcgagt ttaaaccaat tgcgtagat 540  
cttcgtcagc agagcttcac cattgaaggg ctggcggttg gggttattcg caacggcgac 600  
tggtctg 606

<210> 44  
<211> 202  
<212> PRT  
<213> Artificial Sequence

<220>  
<221> misc\_feature  
<223> Novel Sequence

<400> 44

Met Lys Ala Leu Thr Ala Arg Gln Gln Glu Val Phe Asp Leu Ile Arg  
1 5 10 15

Asp His Ile Ser Gln Thr Gly Met Pro Pro Thr Arg Ala Glu Ile Ala  
20 25 30

Gln Arg Leu Gly Phe Arg Ser Pro Asn Ala Ala Glu Glu His Leu Lys  
35 40 45

Ala Leu Ala Arg Lys Gly Val Ile Glu Ile Val Ser Gly Ala Ser Arg  
50 55 60

WO 01/70816

PCT/US01/09050

RH0020.ST25

Gly Ile Arg Leu Leu Gln Glu Glu Glu Glu Gly Leu Pro Leu Val Gly  
65 70 75 80

Arg Val Ala Ala Gly Glu Pro Leu Leu Ala Gln Gln His Ile Glu Gly  
85 90 95

His Tyr Gln Val Asp Pro Ser Leu Phe Lys Pro Asn Ala Asp Phe Leu  
100 105 110

Leu Arg Val Ser Gly Met Ser Met Lys Asp Ile Gly Ile Met Asp Gly  
115 120 125

Asp Leu Leu Ala Val His Lys Thr Gln Asp Val Arg Asn Gly Gln Val  
130 135 140

Val Val Ala Arg Ile Asp Asp Glu Val Thr Val Lys Arg Leu Lys Lys  
145 150 155 160

Gln Gly Asn Lys Val Glu Leu Leu Pro Glu Asn Ser Glu Phe Lys Pro  
165 170 175

Ile Val Val Asp Leu Arg Gln Gln Ser Phe Thr Ile Glu Gly Leu Ala  
180 185 190

Val Gly Val Ile Arg Asn Gly Asp Trp Leu  
195 200

<210> 45  
<211> 271  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> misc feature  
<223> Novel Sequence

<400> 45  
atgggcccta-aaaagaagcg taaagtcgcc ccccgaccg atgtcagcct gggggacgag 60  
ctccacttag acggcgagga cgtggcgatg gcgcatgccg acgcgctaga cgatttcgat 120  
ctggacatgt tgggggacgg ggattcccg gggccgggat ttacccccca cgactccgcc 180  
ccctacggcg ctctggatat ggccgacttc gagtttgagc agatgtttac cgatgccott 240  
ggaattgacg agtacggtgg ggaattcccg g 271

<210> 46  
<211> 90  
<212> PRT  
<213> Artificial Sequence

<220>  
<221> misc feature  
<223> Novel Sequence

<400> 46

Met Gly Pro Lys Lys Lys Arg Lys Val Ala Pro Pro Thr Asp Val Ser  
1 5 10 15

Leu Gly Asp Glu Leu His Leu Asp Gly Glu Asp Val Ala Met Ala His

WO 01/70816

PCT/US01/09050

RH0020.ST25

20 25 30  
 Ala Asp Ala Leu Asp Asp Phe Asp Leu Asp Met Leu Gly Asp Gly Asp  
 35 40 45  
 Ser Pro Gly Pro Gly Phe Thr Pro His Asp Ser Ala Pro Tyr Gly Ala  
 50 55 60  
 Leu Asp Met Ala Asp Phe Glu Phe Glu Gln Met Phe Thr Asp Ala Leu  
 65 70 75 80  
 Gly Ile Asp Glu Tyr Gly Gly Glu Phe Pro  
 85 90

<210> 47  
 <211> 19  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> misc feature  
 <223> Novel Sequence

<400> 47  
 ggagtactgt cctccgagc

19

<210> 48  
 <211> 666  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> misc feature  
 <223> Novel Sequence

<400> 48  
 ggatccccag cttggaattc gacaggttat cagcaacaac acagtcatat ccattctcaa 60  
 ttagctctac cacagtgtgt gaaccaatgt atccagcacc acctgtaacc aaaacaattt 120  
 tagaagtact ttcactttgt aactgagctg tcatttatat tgaattttca aaaattctta 180  
 cttttttttt ggatggacgc aaagaagttt aataatcata ttacatggca ttaccaccat 240  
 atacatatcc atatacatat ccatatctaa tottacctcg actgctgtat ataaaaccag 300  
 tggttatatg tacagtactg ctgtatataa aaccagtggg tatatgtaca gtacgtcgac 360  
 tgctgtatat aaaaccagtg gttatatgta cagtactgct gtatatataa ccagtgggta 420  
 tatgtacagt acgtcgaggg atgataatgc gattagtttt ttagccttat ttctggggta 480  
 attaatcagc gaagcgatga tttttgatct attaacagat atataaatgc aaaaactgca 540  
 taaccacttt aactaatact ttcaacattt tcggtttgta ttacttctta ttcaaatgta 600  
 ataaaagtat caacaaaaaa ttgttaatat acctctatac tttaacgtca aggagaaaaa 660  
 actata 666

<210> 49



WO 01/70816

PCT/US01/09050

RH0020.ST25

<211> 1542  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> misc feature  
<223> Novel Sequence

```

<400> 49
ctggacctga aacacgaagt ggcttaccga ggggtgctcc cagggccaggt gaaggccgaa    60
ccgggggtcc acaacggcca ggtcaacggc cacgtgaggg actggatggc aggcggcgct    120
ggtgcccaatt cgccgtctcc gggagcgggt gctcaacccc agcctaacaa tgggtattcg    180
tcgccactct cctcgggaag ctacgggccc tacagtccaa atgggaaaat aggccgtgag    240
gaactgtcgc cagcttcaag tataaatggg tgcagtacag atggcgaggc acgacgtcag    300
aagaagggcc ctgcgccccg tcagcaagag gaactgtgtc tggtagcgg ggacagagcc    360
tccggatacc actacaatgc gtcacgtgt gaaggggtga aagggttctt cagacggagt    420
gttaccaaaa atgcggttta tatttgtaaa ttcggtcacg ctgcgaaat ggacatgtac    480
atgcgacgga aatgccagga gtgcgcctg aagaagtgtc tagctgtagg catgaggcct    540
gagtgcgtag taccgagac tcagtgcgc atgaagcgga aagagaagaa agcacagaag    600
gagaaggaca aactgcctgt cagcacgacg acggtggacg accacatgcc gccattatg    660
cagtgtgaac ctccacctcc tgaagcagca aggattcacg aagtgtccc aaggtttctc    720
tccgacaagc tgttgagac aaaccggcag aaaaacatcc cccagttgac agccaaccag    780
cagttcctta tcgccaggct catctggtac caggacgggt acgagcagcc ttctgatgaa    840
gatttgaaga ggattacgca gacgtggcag caagcggacg atgaaaacga agagtctgac    900
actcccttcc gccagatcac agagatgact atcctcacg tccaacttat cgtggagttc    960
gcgaagggat tgcagggtt cgccaagatc tcgcagcctg atcaaattac gctgcttaag    1020
gcttgctcaa gtgaggtaat gatgtccga gtgcgcgac gatacgatgc ggccctcagac    1080
agtgttctgt tcgcaacaa ccaagcgtac actcgcgaca actaccgcaa ggctggcatg    1140
gcctacgtca tcgaggatct actgcacttc tgccggtgca tgtactctat ggcggtggac    1200
aacatccatt acgcgctgct cagcgtgtc gtcattttt ctgaccggcc agggttggag    1260
cagccgcaac tgggtgaaga aatccagcgg tactacctga atacgtccg catctatata    1320
ctgaaccagc tgagcgggtc ggcgcgttcg tccgtcatat acggcaagat cctctcaatc    1380
ctctctgagc tacgcacgct cggcatgcaa aactccaaca tgtgcatctc cctcaagctc    1440
aagaacagaa agctgcgcgc tttcctcgag gagatctggg atgtggcgga catgtcgac    1500
acccaaccgc cgcctatcct cgagtcccc acgaatctct ag                                1542

```

<210> 50  
<211> 513

WO 01/70816

PCT/US01/09050

RH0020.ST25

<212> PRT  
<213> Artificial Sequence  
  
<220>  
<221> misc feature  
<223> Novel Sequence

<400> 50

Leu Asp Leu Lys His Glu Val Ala Tyr Arg Gly Val Leu Pro Gly Gln  
1 5 10 15  
Val Lys Ala Glu Pro Gly Val His Asn Gly Gln Val Asn Gly His Val  
20 25 30  
Arg Asp Trp Met Ala Gly Gly Ala Gly Ala Asn Ser Pro Ser Pro Gly  
35 40 45  
Ala Val Ala Gln Pro Gln Pro Asn Asn Gly Tyr Ser Ser Pro Leu Ser  
50 55 60  
Ser Gly Ser Tyr Gly Pro Tyr Ser Pro Asn Gly Lys Ile Gly Arg Glu  
65 70 75 80  
Glu Leu Ser Pro Ala Ser Ser Ile Asn Gly Cys Ser Thr Asp Gly Glu  
85 90 95  
Ala Arg Arg Gln Lys Lys Gly Pro Ala Pro Arg Gln Gln Glu Glu Leu  
100 105 110  
Cys Leu Val Cys Gly Asp Arg Ala Ser Gly Tyr His Tyr Asn Ala Leu  
115 120 125  
Thr Cys Glu Gly Cys Lys Gly Phe Phe Arg Arg Ser Val Thr Lys Asn  
130 135 140  
Ala Val Tyr Ile Cys Lys Phe Gly His Ala Cys Glu Met Asp Met Tyr  
145 150 155 160  
Met Arg Arg Lys Cys Gln Glu Cys Arg Leu Lys Lys Cys Leu Ala Val  
165 170 175  
Gly Met Arg Pro Glu Cys Val Val Pro Glu Thr Gln Cys Ala Met Lys  
180 185 190  
Arg Lys Glu Lys Lys Ala Gln Lys Glu Lys Asp Lys Leu Pro Val Ser  
195 200 205  
Thr Thr Thr Val Asp Asp His Met Pro Pro Ile Met Gln Cys Glu Pro  
210 215 220  
Pro Pro Pro Glu Ala Ala Arg Ile His Glu Val Val Pro Arg Phe Leu  
225 230 235 240  
Ser Asp Lys Leu Leu Glu Thr Asn Arg Gln Lys Asn Ile Pro Gln Leu  
245 250 255  
~~Thr Ala Asn Gln Gln Phe Leu Ile Ala Arg Leu Ile Trp Tyr Gln Asp~~  
260 265 270  
Gly Tyr Glu Gln Pro Ser Asp Glu Asp Leu Lys Arg Ile Thr Gln Thr  
275 280 285  
Trp Gln Gln Ala Asp Asp Glu Asn Glu Glu Ser Asp Thr Pro Phe Arg

WO 01/70816

PCT/US01/09050

RH0020.ST25  
300

290	295	
Gln Ile Thr Glu Met Thr Ile Leu Thr Val Gln Leu Ile Val Glu Phe 305 310 315 320		
Ala Lys Gly Leu Pro Gly Phe Ala Lys Ile Ser Gln Pro Asp Gln Ile 325 330 335		
Thr Leu Leu Lys Ala Cys Ser Ser Glu Val Met Met Leu Arg Val Ala 340 345 350		
Arg Arg Tyr Asp Ala Ala Ser Asp Ser Val Leu Phe Ala Asn Asn Gln 355 360 365		
Ala Tyr Thr Arg Asp Asn Tyr Arg Lys Ala Gly Met Ala Tyr Val Ile 370 375 380		
Glu Asp Leu Leu His Phe Cys Arg Cys Met Tyr Ser Met Ala Leu Asp 385 390 395 400		
Asn Ile His Tyr Ala Leu Leu Thr Ala Val Val Ile Phe Ser Asp Arg 405 410 415		
Pro Gly Leu Glu Gln Pro Gln Leu Val Glu Glu Ile Gln Arg Tyr Tyr 420 425 430		
Leu Asn Thr Leu Arg Ile Tyr Ile Leu Asn Gln Leu Ser Gly Ser Ala 435 440 445		
Arg Ser Ser Val Ile Tyr Gly Lys Ile Leu Ser Ile Leu Ser Glu Leu 450 455 460		
Arg Thr Leu Gly Met Gln Asn Ser Asn Met Cys Ile Ser Leu Lys Leu 465 470 475 480		
Lys Asn Arg Lys Leu Pro Pro Phe Leu Glu Glu Ile Trp Asp Val Ala 485 490 495		
Asp Met Ser His Thr Gln Pro Pro Pro Ile Leu Glu Ser Pro Thr Asn 500 505 510		

Leu

<210> 51  
<211> 4375  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> misc\_feature  
<223> Novel Sequence

<400> 51  
tgtaattttg atgggcgcgcg tgcgcaccg tgtgccatat tgccatccag tcgaatagaa 60  
aaaaaaaaa aaaaaaaaaat atcagttggt ttgtccctcg ctgcctttcg agtgatttcg 120  
gaatattaga cgtcataatt cagcagtgtc ttttaaattt atatagcgat tagcggggcc 180  
gtttgttgga cgtgcgcttg cgttttagtg agtgcaggga tagtgaggcg agtatggtag 240  
ttcgtggtca tgtcaagtgt ggcgaagaaa gacaagccga cgatgtcggg gacggcgctg 300

## RH0020.ST25

atcaactggg	cgcgccggc	gccgccaggc	ccgccgcagc	cgagtcagc	gtcgctgcg	360
ccggcagcca	tgctgcagca	gtccccgacg	cagtcaatgc	agtcgttaaa	ccacatccca	420
actgtcgatt	gtcgctcga	tatgcagtgg	cttaatttag	aacctggatt	catgtcgct	480
atgtcacctc	ctgagatgaa	accagacacc	gccatgcttg	atgggctacg	agacgacgcc	540
acttcgccgc	ctaacttcaa	gaactaccgc	cctaatacacc	ccctgagtgg	ctccaaacac	600
ctatgctcta	tatgcggcga	cagggcgctc	gggaagcact	atgggggtga	cagttgcgaa	660
ggatgcaagg	gtttcttcaa	gcggaccgtc	cggaaggacc	tgctgtacgc	ttgccgggag	720
gagcggaaact	gcatcataga	caagcgacaa	aggaaccgat	gccagtactg	ccgctatcaa	780
aagtgtttgg	cttgcggtat	gaagcgagag	gcggtgcaag	aggagcgcca	gaggaatgct	840
cgcgggcgcg	aggatgcgca	cccagtagc	tcggtgcagg	taagcgatga	gctgtcaatc	900
gagcgcctaa	cggagatgga	gtctttggtg	gcagatccca	gcgaggagtt	ccagttcctc	960
cgcggtgggg	ctgacagcaa	cgtgcctcca	cgttaccgcg	cgcccgctctc	ctccctctgc	1020
caaataggca	acaagcaa	agcggcgttg	gtggtatggg	cgcgcgacat	ccctcatttc	1080
gggcagctgg	agctggacga	tcaagtggta	ctcatcaagg	cctcctggaa	tgagctgcta	1140
ctcttcgcca	tcgctggcg	ctctatggag	tatttggaag	atgagaggga	gaacggggac	1200
ggaacgcgga	gcaccactca	gccacaactg	atgtgtctca	tgccctggcat	gacgttgcac	1260
cgcaactcgg	cgacgcaggc	gggcgtgggc	gccatcttcg	accgcgtgct	gtccgagctc	1320
agtctgaaga	tgcgacacct	gcgcattggc	caggccgagt	acgtcgcgct	caaagccatc	1380
gtgctgctca	accctgatgt	gaaaggactg	aagaatcggc	aagaagtga	cgttttgcga	1440
gaaaaaatgt	tctcttgct	ggacgactac	tgccggcggt	cggaagcaa	cgagggaaggc	1500
cggtttgctg	ccttgctgct	gcggctgcca	gtctccgct	ccatctcgct	caagagcttc	1560
gaacacctct	acttcttcca	cctcgtggcc	gaaggctcca	tcagcgata	catacgagag	1620
gcgctccgaa	accacgcgcc	tccgatcgac	gtcaatgcca	tgatgtaaag	tgcgatacac	1680
gccctgccga	tgtagaaga	actatggcta	atagaagcga	aactgaatac	atctagggtg	1740
ggacttaact	tggaactatc	attaaagtat	cacgcaaatt	atgcgtagtc	agaaagtcgc	1800
gtcgatcaaa	cttttttata	aacgaattga	gtttctaacg	actgcaacac	agcggagttt	1860
tgcttctgat	agtttttatt	ctaattggtta	agatgcttta	cacgggcatt	attgacattc	1920
aagtgttaagt	ggaagttagc	aaccttgaca	tttatatcac	gtttgttaatt	ggttaaataa	1980
attaattaat	cacaagtaag	actaacatca	acgtcacgat	actaacgcca	tttagtgata	2040
ttttcatgt	caagaaactc	attgttttga	taaaatattt	ttctaattac	tcagtgaa	2100
tcacccaaat	gtgaccagct	ttcccgaga	gttcccgtg	taaaatcatc	tttagggaca	2160
tatccccgc	tatctcatga	aattccaagg	atcagtaggg	gccaattccc	ccgatgtgtt	2220
gggaggcaga	attttcgata	atctacgact	attgttagcc	tacgaattag	ttgaattttt	2280

## RH0020.ST25

tgaaattatt tttattaagt cgccactttc caaacacatc agcagggtat atgtgcaatt	2340
ttgtaacgat aactctattc atttctgata tttatcgaaa ttttatctta cataacatgc	2400
tggctggtcc aggtgtttgg tagttacata tgtatctacg gtttgtttta aattatagct	2460
tttttattgt aatctgtata aaattgagtt atcttacttc acactacgat cgagtaaacc	2520
catcgtcagc tacgaaaaac taatcgata aggcgtaaga gtaaataact aattgacaac	2580
cagcaacgag gaccacctca gtccctcgtgc ttacattgtg ccgtagctta atatgatgga	2640
agctgtcgtc gttacgacat tagataaagt gcatgaatac caaaaatgta ccatcccgt	2700
ctgatctctc atgctctcgc tgcgtgggac ccgtgtcgag tgcgtaagg actgactaat	2760
attttagact aggcgtctat gcttcagtaa ttccttatac atattataag tcatccaaat	2820
aacgagtaag gcggcatgtt gagatcagca ttccgagagt caaagagccc ctaacgtgac	2880
tgagaagtag agacaatata ctgattttct gagatgaacg caaccgagat tgacactaaa	2940
aatctattta tggatttcaa aatggcgatg cttgattgtc tgcggtgtgg atagactgaa	3000
atgggtttgc ttaacactgg atattgtttt tattagttaa tagtcttaca ttgcaagttg	3060
gtaattcggg gctaataatcg accggtttgt taactatcta acggttccca gtgtcaggca	3120
cacatctttc ccaagcagac aacgcaagag tgtacaaaat gtacatgtta caaataaagg	3180
aacattcgtc ggataagtgt aacagttgat aggtaaagaa aatggggccg cctctttatt	3240
attacgtagc cgtaaaatta ttaacgtatt tagtttagat gttcagctaa ttaggataat	3300
tctattttgc gagtacctag atgtccatag tgaattaata taataattag actgttacgc	3360
gtaggtaatt ataaagtta ccaaatctct cttcaaagca aaaactttgt acacttccgt	3420
actgagacgt cgtagcttat tctgattcac gaaatatttg gatcacattg ttacaaggcg	3480
accgtcacgt agtatatgat tatttacaaa tgacacgtat gtatcaatgc tataagtgtt	3540
ttcgttacat atgtcgggtc ttaacgtgc atttcgatgt gcagattaaa aatagcaaga	3600
aatcttgaaa ttgttttaga aaatatttga tttccttatt gaaagtatt tttaaatgta	3660
aatatttcgt aatcataata attatgtatt gtgtagttat ttcacctta cgttgggat	3720
attatttaat ggtggcctac gaaagtgatt ataaccatcc gcgtcctcaa aaaggccagt	3780
ttatttttgt acctcatata tactaattac gtaagtaata tcaggcgaat ggttgactaa	3840
caactaacca gtattaaaaa ttaaaagact tcgtcctaataaaaatgtaat atctatgtat	3900
aaaaatgaaa aatctggcgt ataataggta aaattaaact agattgttaa tgaatgtgat	3960
gtctcataaa cgtttagttt ttaatgagaa acatgttttag tcgcctacta taagacgaga	4020
cggaagctc accgagttaa ctgcgtaaca ggaatgttga aaaagatgac acaatttata	4080
tttggtattg aaattatgac taacctgcg ctctatcgtt tgttatggat gcatagtatt	4140
gctgttgaaa ataatggaat taggtaatta ctgcattaat gttgaaaact tgatattatt	4200

WO 01/70816

PCT/US01/09050

RH0020.ST25

ctatggttgg gtatgaattc tatgttgga gttgtgcagc gttgttaaag atgatttata 4260  
atgatgttca ctaaatactct gactaaatgt aagttatttt tttttgtata gacatagctt 4320  
taagatgaag gtgattaaac tttatcctta tcacaataaa aaaaaaaaaa aaaaa 4375

<210> 52  
<211> 472  
<212> PRT  
<213> Artificial Sequence

<220>  
<221> misc\_feature  
<223> Novel Sequence

<400> 52

Met Ser Ser Val Ala Lys Lys Asp Lys Pro Thr Met Ser Val Thr Ala  
1 5 10 15  
Leu Ile Asn Trp Ala Arg Pro Ala Pro Pro Gly Pro Pro Gln Pro Gln  
20 25 30  
Ser Ala Ser Pro Ala Pro Ala Ala Met Leu Gln Gln Leu Pro Thr Gln  
35 40 45  
Ser Met Gln Ser Leu Asn His Ile Pro Thr Val Asp Cys Ser Leu Asp  
50 55 60  
Met Gln Trp Leu Asn Leu Glu Pro Gly Phe Met Ser Pro Met Ser Pro  
65 70 75 80  
Pro Glu Met Lys Pro Asp Thr Ala Met Leu Asp Gly Leu Arg Asp Asp  
85 90 95  
Ala Thr Ser Pro Pro Asn Phe Lys Asn Tyr Pro Pro Asn His Pro Leu  
100 105 110  
Ser Gly Ser Lys His Leu Cys Ser Ile Cys Gly Asp Arg Ala Ser Gly  
115 120 125  
Lys His Tyr Gly Val Tyr Ser Cys Glu Gly Cys Lys Gly Phe Phe Lys  
130 135 140  
Arg Thr Val Arg Lys Asp Leu Ser Tyr Ala Cys Arg Glu Glu Arg Asn  
145 150 155 160  
Cys Ile Ile Asp Lys Arg Gln Arg Asn Arg Cys Gln Tyr Cys Arg Tyr  
165 170 175  
Gln Lys Cys Leu Ala Cys Gly Met Lys Arg Glu Ala Val Gln Glu Glu  
180 185 190  
Arg Gln Arg Asn Ala Arg Gly Ala Glu Asp Ala His Pro Ser Ser Ser  
195 200 205  
~~Val Gln Val Ser Asp Glu Leu Ser Ile Glu Arg Leu Thr Glu Met Glu~~  
210 215 220  
Ser Leu Val Ala Asp Pro Ser Glu Glu Phe Gln Phe Leu Arg Val Gly  
225 230 235 240  
Pro Asp Ser Asn Val Pro Pro Arg Tyr Arg Ala Pro Val Ser Ser Leu

WO 01/70816

PCT/US01/09050

RH0020.ST25

245 250 255  
 Cys Gln Ile Gly Asn Lys Gln Ile Ala Ala Leu Val Val Trp Ala Arg  
 260 265 270  
 Asp Ile Pro His Phe Gly Gln Leu Glu Leu Asp Asp Gln Val Val Leu  
 275 280 285  
 Ile Lys Ala Ser Trp Asn Glu Leu Leu Leu Phe Ala Ile Ala Trp Arg  
 290 295 300  
 Ser Met Glu Tyr Leu Glu Asp Glu Arg Glu Asn Gly Asp Gly Thr Arg  
 305 310 315 320  
 Ser Thr Thr Gln Pro Gln Leu Met Cys Leu Met Pro Gly Met Thr Leu  
 325 330 335  
 His Arg Asn Ser Ala Gln Gln Ala Gly Val Gly Ala Ile Phe Asp Arg  
 340 345 350  
 Val Leu Ser Glu Leu Ser Leu Lys Met Arg Thr Leu Arg Met Asp Gln  
 355 360 365  
 Ala Glu Tyr Val Ala Leu Lys Ala Ile Val Leu Leu Asn Pro Asp Val  
 370 375 380  
 Lys Gly Leu Lys Asn Arg Gln Glu Val Asp Val Leu Arg Glu Lys Met  
 385 390 395 400  
 Phe Ser Cys Leu Asp Asp Tyr Cys Arg Arg Ser Arg Ser Asn Glu Glu  
 405 410 415  
 Gly Arg Phe Ala Ser Leu Leu Leu Arg Leu Pro Ala Leu Arg Ser Ile  
 420 425 430  
 Ser Leu Lys Ser Phe Glu His Leu Tyr Phe Phe His Leu Val Ala Glu  
 435 440 445  
 Gly Ser Ile Ser Gly Tyr Ile Arg Glu Ala Leu Arg Asn His Ala Pro  
 450 455 460  
 Pro Ile Asp Val Asn Ala Met Met  
 465 470

<210> 53  
 <211> 1404  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> misc feature  
 <223> Novel Sequence

<400> 53  
 atggacacca aacatttcct gccgctcgac ttctctaccc aggtgaactc ttcgtccctc 60  
 aactctccaa cgggtcgagg ctccatggct gtcccctcgc tgcacccctc cttgggtcgg 120  
 ggaatcggtc ctccactggg ctgcctggg cagctgcact ctctatcag caccctgagc 180  
 tccccatca atggcatggg tccgcccttc tctgtcatca gctccccat gggcccgcac 240  
 tccatgtcgg taccacaccac accacattg ggcttcggga ctggtagccc ccagctcaat 300

WO 01/70816

PCT/US01/09050

RH0020.ST25

```

tcacccatga accctgtgag cagcactgag gatatcaagc cgccactagg cctcaatggc 360
gtcctcaagg ttcctgcccc tccctcagga aatatggcct ccttcaccaa gcacatctgt 420
gctatctgtg gggaccgctc ctcaggcaaa cactatgggg tatacagttg tgagggtgc 480
aagggcttct tcaagaggac agtacgcaaa gacctgacct acacctgccg agacaacaag 540
gactgcctga tcgacaagag acagcggaac cgggtgtcagt actgccgcta ccagaagtgc 600
ctggccatgg gcatgaagcg ggaagctgtg caggaggagc ggcagcgggg caaggaccgg 660
aatgagaacg aggtggagtc caccagcagt gccaacgagg acatgcctgt agagaagatt 720
ctggaagcgg agcttgctgt cgagcccaag actgagacat acgtggaggc aaacatgggg 780
ctgaacccca gctcaccaaa tgacctgtt accaacaatct gtcaagcagc agacaagcag 840
ctcttcactc ttgtggagtg ggccaagagg atcccacact tttctgagct gcccctagac 900
gaccaggtca tcctgctacg ggcaggctgg aacgagctgc tgatcgctc cttctccac 960
cgctccatag ctgtgaaaga tgggattctc ctggccaccg gctgcacgt acaccggaac 1020
agcgctcaca gtgctggggt gggcgccatc ttgacaggg tgctaacaga gctggtgtct 1080
aagatgcgtg acatgcagat ggacaagacg gagctgggct gcctgcgagc cattgtcctg 1140
ttcaaccctg actctaaggg gctctcaaac cctgctgagg tggaggcgtt gagggagaag 1200
gtgtatgcgt cactagaagc gtactgcaaa cacaagtacc ctgagcagcc gggcaggttt 1260
gccaagctgc tgctccgct gctgcactg cgttccatcg ggctcaagt cctggagcac 1320
ctgttcttct tcaagctcat cggggacacg cccatcgaca ccttctcat ggagatgctg 1380
gaggcaccac atcaagccac ctg 1404

```

<210> 54  
 <211> 467  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <221> misc feature  
 <223> Novel Sequence

<400> 54

Met Asp Thr Lys His Phe Leu Pro Leu Asp Phe Ser Thr Gln Val Asn  
1 5 10 15

Ser Ser Ser Leu Asn Ser Pro Thr Gly Arg Gly Ser Met Ala Val Pro  
20 25 30

Ser Leu His Pro Ser Leu Gly Pro Gly Ile Gly Ser Pro Leu Gly Ser  
35 40 45

Pro Gly Gln Leu His Ser Pro Ile Ser Thr Leu Ser Ser Pro Ile Asn  
50 55 60

Gly Met Gly Pro Pro Phe Ser Val Ile Ser Ser Pro Met Gly Pro His  
65 70 75 80



WO 01/70816

PCT/US01/09050

RH0020.ST25

Ser Met Ser Val Pro Thr Thr Pro Thr Leu Gly Phe Gly Thr Gly Ser  
85 90 95

Pro Gln Leu Asn Ser Pro Met Asn Pro Val Ser Ser Thr Glu Asp Ile  
100 105 110

Lys Pro Pro Leu Gly Leu Asn Gly Val Leu Lys Val Pro Ala His Pro  
115 120 125

Ser Gly Asn Met Ala Ser Phe Thr Lys His Ile Cys Ala Ile Cys Gly  
130 135 140

Asp Arg Ser Ser Gly Lys His Tyr Gly Val Tyr Ser Cys Glu Gly Cys  
145 150 155 160

Lys Gly Phe Phe Lys Arg Thr Val Arg Lys Asp Leu Thr Tyr Thr Cys  
165 170 175

Arg Asp Asn Lys Asp Cys Leu Ile Asp Lys Arg Gln Arg Asn Arg Cys  
180 185 190

Gln Tyr Cys Arg Tyr Gln Lys Cys Leu Ala Met Gly Met Lys Arg Glu  
195 200 205

Ala Val Gln Glu Glu Arg Gln Arg Gly Lys Asp Arg Asn Glu Asn Glu  
210 215 220

Val Glu Ser Thr Ser Ser Ala Asn Glu Asp Met Pro Val Glu Lys Ile  
225 230 235 240

Leu Glu Ala Glu Leu Ala Val Glu Pro Lys Thr Glu Thr Tyr Val Glu  
245 250 255

Ala Asn Met Gly Leu Asn Pro Ser Ser Pro Asn Asp Pro Val Thr Asn  
260 265 270

Ile Cys Gln Ala Ala Asp Lys Gln Leu Phe Thr Leu Val Glu Trp Ala  
275 280 285

Lys Arg Ile Pro His Phe Ser Glu Leu Pro Leu Asp Asp Gln Val Ile  
290 295 300

Leu Leu Arg Ala Gly Trp Asn Glu Leu Leu Ile Ala Ser Phe Ser His  
305 310 315 320

Arg Ser Ile Ala Val Lys Asp Gly Ile Leu Leu Ala Thr Gly Leu His  
325 330 335

Val His Arg Asn Ser Ala His Ser Ala Gly Val Gly Ala Ile Phe Asp  
340 345 350

Arg Val Leu Thr Glu Leu Val Ser Lys Met Arg Asp Met Gln Met Asp  
355 360 365

Lys Thr Glu Leu Gly Cys Leu Arg Ala Ile Val Leu Phe Asn Pro Asp  
370 375 380

~~Ser Lys Gly Leu Ser Asn Pro Ala Glu Val Glu Ala Leu Arg Glu Lys~~  
385 390 395 400

Val Tyr Ala Ser Leu Glu Ala Tyr Cys Lys His Lys Tyr Pro Glu Gln  
405 410 415

Pro Gly Arg Phe Ala Lys Leu Leu Leu Arg Leu Pro Ala Leu Arg Ser

WO 01/70816

PCT/US01/09050

RH0020.ST25

420 425 430  
Ile Gly Leu Lys Cys Leu Glu His Leu Phe Phe Phe Lys Leu Ile Gly  
435 440 445  
Asp Thr Pro Ile Asp Thr Phe Leu Met Glu Met Leu Glu Ala Pro His  
450 455 460

Gln Ala Thr  
465

<210> 55  
<211> 309  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> misc feature  
<223> Novel Sequence

<400> 55  
ggtgtggaat gtccccaggc tccccagcag gcagaagtat gcaaagcatg catctcaatt 60  
agtcagcaac caggtgtgga aagtcgccag gctccccagc aggcagaagt atgcaaagca 120  
tgcatctcaa ttagtcagca accatagtcc cgcccctaac tccgcccatc ccgccctaa 180  
ctccgccag ttccgccat tctccgcc atgggtgact aattttttt atttatgcag 240  
aggccgaggc cgcccgcc tctgagctat tccagaagta gtgaggaggc ttttttggag 300  
gcctaggct 309

<210> 56  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> misc feature  
<223> Novel Sequence

<400> 56  
tatataatgg atccccgggt accg 24

<210> 57  
<211> 1653  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> misc feature  
<223> Novel Sequence

<400> 57  
atggaagacg ccaaaaacat aaagaaaggc cggcgccat tctatcctct agaggatgga 60  
accgctggag agcaactgca taaggctatg aagagatagc cctgtgttcc tggacaatt 120  
gcttttacag atgcacatat cgaggatgaac atcacgtacg cggaataactt cgaaatgtcc 180

## RH0020.ST25

```

gttcggttg cagaagctat gaaacgatat gggctgaata caaatcacag aatcgtcgta 240
tgcagtga aa actctcttca attctttatg ccggtgttg gcgcgttatt tatcggagtt 300
gcagttgcgc ccgcgaacga cttttataat gaacgtgaat tgctcaacag tatgaacatt 360
tcgcagccta ccgtagtggt tgtttccaaa aaggggttgc aaaaaatttt gaacgtgcaa 420
aaaaaattac caataatcca gaaaattatt atcatggatt ctaaaacgga ttaccagga 480
tttcagtcga tgtacacggt cgtcacatct catctacctc ccggttttaa tgaatacgat 540
tttgtaccag agtcttttga tcgtgacaaa acaattgcac tgataatgaa ttcctctgga 600
tctactgggt tacctaaggg tgtggccctt ccgcatacaa ctgcctgcgt cagattctcg 660
catgccagag atcctatttt tggcaatcaa atcattccgg atactgcgat ttaagtgtt 720
gttccattcc atcacggttt tggaatgttt actacactcg gatatttgat atgtggattt 780
cgagtcgtct taatgtatag atttgaagaa gagctgtttt tacgatccct tcaggattac 840
aaaattcaaa gtgcgttgct agtaccaacc ctattttcat tcttcgcaa aagcactctg 900
attgacaaat acgatttata taatttacac gaaattgctt ctgggggcgc acctcttctg 960
aaagaagtcg ggaagcgggt tgcaaacgc ttccatcttc cagggatcgc acaaggatat 1020
gggctcactg agactacatc agctattctg attacaccgc agggggatga taaaccgggc 1080
gcggtcggta aagttgttcc attttttgaa gcgaaggttg tggatctgga taccgggaaa 1140
acgctgggcy ttaatcagag aggcgaatta tgtgtcagag gacctatgat tatgtccggt 1200
tatgtaaaca atccggaagc gaccaacgcc ttgattgaca aggatggatg gctacattct 1260
ggagacatag cttactggga cgaagacgaa cacttcttca tagttgaccg cttgaagtct 1320
ttaattaaat acaaggata tcaggtggcc ccgctgaat tggatcgat attgttaca 1380
caccacaaca tcttcgacgc gggcgtggca ggtcttcccg acgatgacgc cggtgaactt 1440
ccgcgcgcgc ttgttgtttt ggagcacgga aagacgatga cggaaaaaga gatcgtggat 1500
tacgtcgcca gtcaagtaac aaccgcgaaa aagttgcgcg gagagttgt gtttgtggac 1560
gaagtaccga aaggtcttac cggaaaactc gacgcaagaa aaatcagaga gatcctcata 1620
aaggccaaga agggcgga aa gtccaaattg taa 1653

```

<210> 58  
 <211> 867  
 <212> DNA  
 <213> Artificial Sequence

<220>

<221> misc feature

<223> Novel Sequence

<400> 58  
 aagcgagagg cggtgcaaga ggagcgccag aggaatgctc gcggcgcgga ggatgcgcac 60

WO 01/70816

PCT/US01/09050

RH0020.ST25

ccgagtagct cggtgcaggt aagcgatgag ctgtcaatcg agcgccctaac ggagatggag	120
tctttggtgg cagatcccag cgaggagtgc cagttcctcc gcgtggggcc tgacagcaac	180
gtgcctccac gttaccgcgc gcccgctctcc tccctctgcc aaataggcaa caagcaaata	240
gcggcggttg tggtatgggc gcgcgacatc cctcatttcg ggcagctgga gctggacgat	300
caagtggtag tcatcaaggc ctccctggaat gagctgctac tcttcgcat cgccctggcg	360
tctatggagt atttgaaga tgagaggag aacggggacg gaacgcggag caccactcag	420
ccacaactga tgtgtctcat gcctggcatg acgttgacc gcaactcggc gcagcaggcg	480
ggcgtgggag ccactcttga ccgcgtgctg tccgagctca gtctgaagat gcgcaccttg	540
cgcatggacc aggcggagta cgtcgcgtc aaagccatcg tgctgctcaa ccctgatgtg	600
aaaggactga agaatcggca agaagtgac gttttgcgag aaaaaatgtt ctcttgccctg	660
gacgactact gccggcggtc gcgaagcaac gaggaaggcc gggttgctgc cttgctgctg	720
cggtgcccag ctctccgctc catctcgtc aagagcttcg aacacctcta cttctccac	780
ctcgtggccg aaggctccat cagcggatac atacgagagg cgctccgaaa ccacgcgcct	840
ccgatcgacg tcaatgccat gatgtaa	867

<210> 59  
 <211> 225  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> misc feature  
 <223> Novel Sequence

<400> 59	
tcgacattgg acaagtgcac tgaacccttg tctctcgaga gacaaggggg ttcaatgcac	60
ttgtccaatg tcgagagaca aggggggttc atgcacttgt ccaatgtcga gagacaagg	120
ggttcaatgc acttggtcaa tgcgagaga caaggggggt caatgcactt gtccaatgtc	180
gagagacaag ggggttcaat gcacttgctc aatgtcgact ctaga	225

<210> 60  
 <211> 619  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> misc feature  
 <223> Novel Sequence

<400> 60	
cgttacataa cttacggtaa atggcccgcc tggtgaccg cccaacgacc cccgccatt	60
gacgtcaata atgacgtatg ttccatagt aacgccaata gggactttcc attgacgtca	120
atgggtyggag tatttacggg aaactgccca cttggcagta catcaagtgt atcatatgcc	180

## RH0020.ST25

aagtacgccc cctattgacg tcaatgacgg taaatggccc gcctggcatt atgccagta 240  
 catgacctta tgggactttc ctacttggca gtacatctac gtattagtca tcgctattac 300  
 catggtgatg cggtttttggc agtacatcaa tgggcgtgga tagcggtttg actcacgggg 360  
 atttccaagt ctccacccca ttgacgtcaa tgggagtttg ttttggcacc aaaatcaacg 420  
 ggactttcca aaatgtcgta acaactccgc cccattgacg caaatgggcg gtaggcgtgt 480  
 acggtgggag gtctatataa gcagagctcg tttagtgaac cgtcagatcg cctggagacg 540  
 ccatccacgc tgttttgacc tccatagaag acaccgggac cgatccagcc tccgcggccg 600  
 ggaacggtgc attggaacg 619

<210> 61  
 <211> 262  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> misc\_feature  
 <223> Novel Sequence

<400> 61  
 atgtagtctt atgcaactact cttgtagtct tgcaacatgg taacgatgag ttagcaacat 60  
 gccttacaag gagagaaaaa gcaccgtgca tgccgatagg tggaagtaag gtggtacgat 120  
 cgtgccttat taggaaggca acagacgggt ctgacatgga ttggacgaac cactgaattc 180  
 cgcattgcag agatattgta ttttaagtgc tagctcgata caataaacgc catttgacca 240  
 ttcaccacat tggagtgcac ct 262

<210> 62  
 <211> 1247  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> misc\_feature  
 <223> Novel Sequence

<400> 62  
 tctatttcct caggccgtga ggaactgtcg ccagcttcaa gtataaatgg gtgcagtaca 60  
 gatggcgagg cagcagctca gaagaagggc cctgcgcccc gtcagcaaga ggaactgtgt 120  
 ctggtatgcg gggacagagc ctccggatac cactacaatg cgctcacgtg tgaaggtgt 180  
 aaagggttct tcagacggag tgttaccaaa aatgcggttt atattttaa attcggtcac 240  
 gcttgcgaaa tggacatgta catgcgacgg aaatgccagg agtgccgcct gaagaagtgc 300  
 ttagctgtag gcatgagggc tgagtgcgta gtacccgaga ctacgtgcgc catgaagcgg 360  
 aaagagaaga aagcacagaa ggagaaggac aaactgcctg tcagcacgac gacggtggac 420

WO 01/70816

PCT/US01/09050

RH0020.ST25

gaccacatgc cgccattat gcagtgtgaa cctccacctc ctgaagcagc aaggattcac 480  
 gaagtgggtcc caaggtttct ctccgacaag ctgttgaggaga caaaccggca gaaaaacatc 540  
 ccccagttga cagccaacca gcagttcctt atcgccaggc tcattctggtta ccaggacggg 600  
 tacgagcagc cttctgatga agatttgaag aggattacgc agacgtggca gcaagcggac 660  
 gatgaaaacg aagagtctga cactcccttc cgccagatca cagagatgac tatectcacg 720  
 gtccaactta tcgtggagtt cgcaaggga ttgccagggt tcgccaagat ctgcgagcct 780  
 gatcaaatga cgctgcttaa ggcttgctca agtgaggtaa tgatgctccg agtcgcgcga 840  
 cgatacagtg cggcctcaga cagtgttctg ttgcggaaca accaagcgta cactcgcgac 900  
 aactaccgca aggttggtcat ggcttacgtc atcgaggatc tactgcactt ctgccggtgc 960  
 atgtactcta tggcggttga caacatccat tacgcgctgc tcacggctgt cgtcatcttt 1020  
 tctgaccggc cagggttga gacgacgaa ctggtggaag aaatccagcg gtactacctg 1080  
 aatacgctcc gcatttatat cctgaaccag ctgagcgggt cggcgcgctc gtccgtcata 1140  
 tacggcaaga tcctctcaat cctctctgag ctacgcacgc tcggcatgca aaactccaac 1200  
 atgtgcatct ccctcaagct caagaacaga aagctgccgc ctttcct 1247

<210> 63  
 <211> 440  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <221> misc feature  
 <223> Novel Sequence

<400> 63

Ser Ile Ser Ser Gly Arg Glu Glu Leu Ser Pro Ala Ser Ser Ile Asn  
 1 5 10 15  
 Gly Cys Ser Thr Asp Gly Glu Ala Arg Arg Gln Lys Lys Gly Pro Ala  
 20 25 30  
 Pro Arg Gln Gln Glu Glu Leu Cys Leu Val Cys Gly Asp Arg Ala Ser  
 35 40 45  
 Gly Tyr His Tyr Asn Ala Leu Thr Cys Glu Gly Cys Lys Gly Phe Phe  
 50 55 60  
 Arg Arg Ser Val Thr Lys Asn Ala Val Tyr Ile Cys Lys Phe Gly His  
 65 70 75 80  
 Ala Cys Glu Met Asp Met Tyr Met Arg Arg Lys Cys Gln Glu Cys Arg  
 85 90 95  
 Leu Lys Lys Cys Leu Ala Val Gly Met Arg Pro Glu Cys Val Val Pro  
 100 105 110  
 Glu Thr Gln Cys Ala Met Lys Arg Lys Glu Lys Lys Ala Gln Lys Glu  
 115 120 125

WO 01/70816

PCT/US01/09050

RH0020.ST25

Lys Asp Lys Leu Pro Val Ser Thr Thr Thr Val Asp Asp His Met Pro  
 130 135 140  
 Pro Ile Met Gln Cys Glu Pro Pro Pro Pro Glu Ala Ala Arg Ile His  
 145 150 155 160  
 Glu Val Val Pro Arg Phe Leu Ser Asp Lys Leu Leu Glu Thr Asn Arg  
 165 170 175  
 Gln Lys Asn Ile Pro Gln Leu Thr Ala Asn Gln Gln Phe Leu Ile Ala  
 180 185 190  
 Arg Leu Ile Trp Tyr Gln Asp Gly Tyr Glu Gln Pro Ser Asp Glu Asp  
 195 200 205  
 Leu Lys Arg Ile Thr Gln Thr Trp Gln Gln Ala Asp Asp Glu Asn Glu  
 210 215 220  
 Glu Ser Asp Thr Pro Phe Arg Gln Ile Thr Glu Met Thr Ile Leu Thr  
 225 230 235 240  
 Val Gln Leu Ile Val Glu Phe Ala Lys Gly Leu Pro Gly Phe Ala Lys  
 245 250 255  
 Ile Ser Gln Pro Asp Gln Ile Thr Leu Leu Lys Ala Cys Ser Ser Glu  
 260 265 270  
 Val Met Met Leu Arg Val Ala Arg Arg Tyr Asp Ala Ala Ser Asp Ser  
 275 280 285  
 Val Leu Phe Ala Asn Asn Gln Ala Tyr Thr Arg Asp Asn Tyr Arg Lys  
 290 295 300  
 Ala Gly Met Ala Tyr Val Ile Glu Asp Leu Leu His Phe Cys Arg Cys  
 305 310 315 320  
 Met Tyr Ser Met Ala Leu Asp Asn Ile His Tyr Ala Leu Leu Thr Ala  
 325 330 335  
 Val Val Ile Phe Ser Asp Arg Pro Gly Leu Glu Gln Pro Gln Leu Val  
 340 345 350  
 Glu Glu Ile Gln Arg Tyr Tyr Leu Asn Thr Leu Arg Ile Tyr Ile Leu  
 355 360 365  
 Asn Gln Leu Ser Gly Ser Ala Arg Ser Ser Val Ile Tyr Gly Lys Ile  
 370 375 380  
 Leu Ser Ile Leu Ser Glu Leu Arg Thr Leu Gly Met Gln Asn Ser Asn  
 385 390 395 400  
 Met Cys Ile Ser Leu Lys Leu Lys Asn Arg Lys Leu Pro Pro Phe Leu  
 405 410 415  
 Glu Glu Ile Trp Asp Val Ala Asp Met Ser His Thr Gln Pro Pro Pro  
 420 425 430

Ile Leu Glu Ser Pro Thr Asn Leu

435 440

<210> 64  
 <211> 943  
 <212> DNA  
 <213> Artificial Sequence

WO 01/70816

PCT/US01/09050

RH0020.ST25

<220>  
<221> misc feature  
<223> Novel Sequence

<400> 64  
atgacttcga aagtttatga tccagaacaa aggaaacgga tgataactgg tccgcagtgg 60  
tgggccagat gtaaacaaat gaatgttctt gattcattta ttaattatta tgattcagaa 120  
aaacatgcag aaaatgctgt tatTTTTTTT catggtaacg cggcctcttc ttatttatgg 180  
cgacatgttg tgccacatat tgagccagta gcgcggtgta ttataccaga cttatttgg 240  
atgggcaa at caggcaa atc tggtaatggt tcttataggt tacttgatca ttacaaatat 300  
cttactgcat gggttgaa ct tcttaattta ccaaagaaga tcatttttgt cggccatgat 360  
tggggtgctt gtttggcatt tcattatagc tatgagcatc aagataagat caaagcaata 420  
gttcacgctg aaagtgtagt agatgtgatt gaatcatggg atgaatggcc tgatattgaa 480  
gaagatattg cgttgatcaa atctgaagaa ggagaaaaaa tggttttgga gaataacttc 540  
ttcgtgga ccatgttgcc atcaaaaatc atgagaaagt tagaaccaga agaatttgca 600  
gcatactctg aaccattcaa agagaaaggt gaagtctgct gtccaacatt atcatggcct 660  
cgtgaaatcc cgttagtaaa aggtggtaaa cctgacgttg taaaaattgt taggaattat 720  
aatgcttatc tacgtgcaag tgatgattta ccaaaaatgt ttattgaatc ggaccagga 780  
ttcttttcca atgctattgt tgaagggtgc aagaagtttc ctaatactga atttgcataa 840  
gtaaaaggtc ttcatttttc gcaagaagat gcacctgatg aaatgggaaa atatatcaaa 900  
tcgttcgttg agcgagttct caaaaatgaa caataattct aga 943